

EXHIBIT 18

DISCUSSION DOCUMENT

SEROQUEL™

**DIABETES MELLITUS, DIABETIC KETOACIDOSIS, NON-KETOTIC
HYPEROSMOLAR COMA, AND HYPERGLYCAEMIA**

***ALL FINDINGS PRESENTED IN THIS DOCUMENT ARE TO BE SUBJECT
TO FURTHER CONSIDERATION AT SERM***

SERM NO 32 - 22 JUNE 2000 - MINUTES

Participants : M Brecher

After review of the available data SERM considered that no amendment to the CDS was justified

ACTION - WG create a position paper.

MB keep this under review and bring back to SERM the expected additional clinical trial data

AUTHOR(S):

**Wayne K. Geller MD
Medical Director, Drug Safety
Wilmington, DE**

SIGNATURE:

DATE:

'SEROQUEL' is a trademark, the property of AstraZeneca Limited

SUMMARY AND CONCLUSIONS:

Presently, the SEROQUEL Core Data Sheet (CDS) does not include any references to diabetes mellitus, diabetic ketoacidosis, or hyperglycaemia associated with SEROQUEL therapy. Safety data derived from clinical trials and spontaneous reports, despite often containing limited information, suggest the possibility of an association between SEROQUEL use and impaired glucose regulation including occasional reports of new onset diabetes mellitus. While none of these reports are absolutely steadfast, the number of reports is fairly sizeable. Currently, no such signals exist for the complications of diabetes such as non-ketotic hyperosmolar coma or diabetic ketoacidosis. While there were no reports of positive dechallenges and rechallenges, consideration should be given to the suggestion that SEROQUEL therapy may cause impaired glucose regulation including diabetes mellitus in certain individuals.

1 INTRODUCTION

In May 2000 FDA notified AstraZeneca that, based upon review of postmarketing safety data for SEROQUEL and other atypical antipsychotics, they were further investigating a possible signal for new onset diabetes mellitus (NODM), non-ketotic hyperosmolar coma (NKHOC), and diabetic ketoacidosis (DKA). FDA expressed concern that increased market exposure could result in an increased number of reports of these events as has been observed with similar agents. In their correspondence (see attachment), they have requested "more extensive safety information" from all phases of clinical development to the present for SEROQUEL for their review. This discussion document will specifically address FDA's third item on their list of requests, "A review of spontaneous postmarketing reports for new-onset diabetes mellitus, hyperglycaemia, hyperosmolar coma, diabetic ketoacidosis, and weight gain".

Reviewing the labeling for other atypical antipsychotic medications (risperidone, olanzapine, clozapine, and sertindole)¹, several of these products have in their labels statements regarding diabetes mellitus, diabetic ketoacidosis, and/or severe hyperglycaemia.

Risperidone: No mention of diabetes or hyperglycaemia.

Olanzapine: Warnings and Precautions: Hyperglycaemia or exacerbation of pre-existing diabetes has been reported in very rare cases during Zyprexa treatment. In some cases, a prior increase in body weight has been reported, which may be a predisposing factor. Appropriate clinical monitoring is advisable in diabetic patients and in patients with risk factors for the development of diabetes mellitus.

Frequent (>10%): Hyperglycaemia or exacerbation of pre-existing diabetes has been reported in very rare cases (see also section 4.4, 'Special warnings and special precautions for use').

Clozapine: Warnings and Precautions:**Hyperglycaemia:**

Severe hyperglycaemia, sometimes leading to ketoacidosis, has been reported during CLOZARIL® (clozapine) treatment in patients with no prior history of hyperglycaemia. While a causal relationship to CLOZARIL® (clozapine) use has not been definitively established, glucose levels normalized in most patients after discontinuation of CLOZARIL® (clozapine), and a rechallenge in one patient produced a recurrence of hyperglycaemia. The effect of CLOZARIL® (clozapine) on glucose metabolism in patients with diabetes mellitus has not been studied. The possibility of impaired glucose tolerance should be considered in patients receiving CLOZARIL® (clozapine) who develop symptoms of hyperglycaemia, such as polydipsia, polyuria, polyphagia, and weakness. In patients with significant treatment-emergent hyperglycaemia, the discontinuation of CLOZARIL® (clozapine) should be considered.

Sertindole: Warnings and Precautions: *Diabetic patients:* serdolect may modify insulin and glucose responses in diabetic patients calling for adjustment of antidiabetic therapy.

Side-effects: Convulsions, hyperglycaemia and syncope have been reported rarely.

2 BACKGROUND

The SEROQUEL core data sheet (CDS) last revised in March 2000 does not include listings for NODM, hyperglycaemia, NKHOC, or DKA. The following statement addresses the issue of weight gain with SEROQUEL:

“As with other antipsychotics, SEROQUEL may also be associated with limited weight gain, predominantly during the early weeks of treatment”.

The SEROQUEL US package insert (PI) includes hyperglycaemia and diabetes mellitus as labeled events occurring infrequently (in 1/100 to 1/1000 patients) according to premarketing clinical trial safety data. The aforementioned complications of diabetes mellitus are not in the PI.

Patients with either impaired glucose tolerance (IGT) or frank diabetes mellitus have hyperglycaemia². The term IGT represents a metabolic condition between normal glucose homeostasis and diabetes mellitus. This includes individuals with fasting glucose levels ≥ 110 mg/dl (6.1 mmol/l) but < 126 mg/dl (7.0 mmol/l).

Diabetes mellitus is a group of metabolic diseases characterized by hyperglycaemia resulting from defects in insulin secretion, insulin action, or both. The criteria for the diagnosis of DM are as follows:

- (1) Symptoms of DM (polyuria, polydipsia, and unexplained weight loss) plus random plasma glucose concentration ≥ 200 mg/dl (11.1 mmol/l); or
- (2) Minimum 8 hour fasting plasma glucose ≥ 126 mg/dl (7.0 mmol/l); or
- (3) Two hour plasma glucose ≥ 200 mg/dl (11.1 mmol/l) during an oral glucose tolerance test using a glucose equivalent of 75 g anhydrous glucose dissolved in water

Patients with diabetes mellitus are classified as having Type I or Type 2 disease. Patients with Type 1 disease are absolute insulin deficient with β -cell destruction and are at risk for developing DKA. Patients who develop Type 2 disease have both abnormal insulin secretion and insulin resistance in target tissues and are not at risk for developing DKA. It is felt that insulin resistance in these patients is the primary event and that obesity contributes to insulin resistance³. Type 2 diabetes mellitus is most prevalent and is thought to be a polygenic disease. The majority of patients with Type 2 disease are obese ($\geq 120\%$ desirable body weight or a BMI ≥ 27 kg/m²), but this is not thought to be the only factor that contributes to insulin resistance. Individuals with dyslipidemia and/or hypertension are at increased risk. There is a strong genetic predisposition to Type 2 disease. It is well known that a modest weight reduction in an obese individual with

Type 2 DM frequently results in significant reduction in blood glucose levels. This is the cornerstone of therapy in patients with Type 2 diabetes mellitus, prior to and during treatment with pharmacologic agents.

Diseases and conditions that have been associated with diabetes mellitus include pancreatic diseases, acromegaly, Cushing's syndrome, glucagonoma, pheochromocytoma, hyperthyroidism, somatostatinoma, aldosteronoma, congenital rubella, cytomegalovirus, pregnancy, and others. Drugs that are known to cause or contribute to hyperglycaemia include: pentamidine, nicotinic acid, glucocorticoids, thyroxine, diazoxide, β -agonists, thiazide diuretics, phenytoin, α -interferon, and others.

Criteria for testing for DM in otherwise asymptomatic, undiagnosed individuals are as follows:

Individuals ≥ 45 years of age, or younger in patients who:

- Are obese ($\geq 120\%$ desirable body weight or a BMI ≥ 27 kg/m²)
- Have a first degree relative with DM
- Belong to high-risk population
- Delivered a ≥ 9 pound baby or have been diagnosed with gestational DM
- Are hypertensive ($\geq 140/90$ mmHg)

- Have hyperlipidemia
- Have had abnormal IGT or IFG

3 THE LITERATURE

Wilson et al⁴ presented a poster entitled, *New-Onset Diabetes and Ketoacidosis with Atypical Antipsychotics* at the American College of Neuropsychopharmacology Annual Meeting, December 12 to 16, 1999, in Acapulco, Mexico. They evaluated the risk of patients using atypical antipsychotics for developing new-onset diabetes mellitus and ketoacidosis. Their interest evolved from case reports in the literature describing altered glucose metabolism in patients receiving atypical antipsychotic agents (notably clozapine, olanzapine, and quetiapine). They conducted a retrospective analysis of the Ohio Department of Mental Health database searching for patients treated with an atypical antipsychotic agent who had also been evaluated or treated for diabetes mellitus. In 11 of 126 (8.7%) of patients receiving clozapine, olanzapine, or quetiapine were diagnosed with new-onset, acute, or market glucose intolerance. Six of these patients required insulin (4 short-term) and five developed DKA. Confounding these results are that only 21/126 patients studied had baseline fasting glucose and that only 14 patients had follow-up studies. Their findings were that:

- (1) The mean and median time to onset of diabetic ketoacidosis after starting treatment with atypical antipsychotic medications were 81 and 33 days, respectively (N=5).
- (2) Changes in glucose tolerance were not related to significant weight gain and often occurred during the first 6 weeks of treatment. Mean and median weight gains in patients with new-onset DM were 16 and 8 pounds, respectively.

Reviewing the labeling for other atypical antipsychotic medications (risperidone, olanzapine, clozapine, and sertindole)⁴, several of these products have in their label statements regarding diabetes mellitus, diabetic ketoacidosis, and/or severe hyperglycaemia.

Risperidone: No mention of diabetes or hyperglycaemia.

Olanzapine: Warnings and Precautions: Hyperglycaemia or exacerbation of pre-existing diabetes has been reported in very rare cases during Zyprexa treatment. In some cases, a prior increase in body weight has been reported, which may be a predisposing factor. Appropriate clinical monitoring is advisable in diabetic patients and in patients with risk factors for the development of diabetes mellitus.

Frequent (>10%): Hyperglycaemia or exacerbation of pre-existing diabetes has been reported in very rare cases (see also section 4.4, 'Special warnings and special precautions for use').

Clozapine: Warnings and Precautions:

Undesirable effects: On rare occasions, hyperglycaemia has been reported in patients on clozaril treatment.

Sertindole: Warnings and Precautions: *Diabetic patients:* serdolect may modify insulin and glucose responses in diabetic patients calling for adjustment of antidiabetic therapy.

Side-effects: *Convulsions, hyperglycaemia and syncope have been reported rarely.*

4 CLINTRACE DATABASE (IN HOUSE SAFETY DATA)

A search was conducted for all cases in which diabetes mellitus, hyperglycaemia, diabetic ketoacidosis, and non-ketotic hyperosmolar coma were reported with SEROQUEL. The following are narratives for these 28 cases.

Case Number: 2000UW01164**KETOACIDOSIS, DIABETES MELLITUS, POLYURIA, POLYDIPSIA, WEIGHT LOSS, ELEVATED GLUCOSE LEVEL**

A report has been received from a registered pharmacist, via sales rep, concerning a 43 year old male with a history of mental illness who started SEROQUEL 200 mg HS in December 1999. Over a period of a few weeks he developed polyuria, polydipsia, and an unexplained weight loss of over 30 lbs. Fasting blood sugar showed glucose level over 700. Patient developed ketoacidosis and was hospitalized where a diagnosis of new onset diabetes mellitus was made. (Patient has no family history of diabetes). Patient is also receiving "fenlaxin" (venlafaxine) and continues on SEROQUEL. More information will be sought.

Case Number: 2000UW01047**COUGH, ELEVATED CHOLESTEROL, WEIGHT GAIN, CONSTIPATION, ASTHMA, WORSENING FIBROMYALGIA, MUSCLE SPASMS, TENSENESS IN NECK, URINE ODOR, WORSENING ARTHRITIS, WORSENING ENDOMETRIOSIS, ELEVATED BLOOD SUGAR, EXCITABLE, DIFFICULTY IN WAKING, NEGATIVE MOOD, DECREASED SEX DRIVE, INABILITY TO HAVE ORGASMS**

A report has been received from a nutritionist, who is also the patient, who has been receiving SEROQUEL, six 25 mg tablets at night, since September 1997 for psychotic episodes. She has been experiencing cough, elevated cholesterol, weight gain, constipation, asthma, worsening fibromyalgia, muscle spasms in her neck and back, tenseness in neck, urine odor, worsening arthritis, worsening endometriosis, elevated blood sugar, she is more excitable, has difficulty in waking, negative mood, decreased sex drive, and inability to have orgasms.

Case Number: 2000UW00266

DIABETES MELLITUS

A report has been received from a pharmacist concerning a 12 year old male patient who had been receiving SEROQUEL 300 mg daily since 06 December 1999. On 26 January 2000, the patient experienced hyperglycaemia (blood sugar level of 863) and was hospitalized. SEROQUEL was discontinued and the patient was treated with insulin. His blood sugar level has decreased to 170. Concomitant medications include zoloft, klonopin, haldol and depakote.

Follow-up 23 February 2000: Pharm D reports that "after further testing, the attending physicians did not feel that SEROQUEL was involved in the patient's hyperglycaemia. The patient was diagnosed with insulin dependent diabetes mellitus. The patient is currently being seen in the diabetic clinic."

Case Number: 1999UW03532

DIABETES MELLITUS, WEIGHT GAIN

A report has been received from a physician concerning a 45 year old female who has been receiving SEROQUEL and developed diabetes. Physician feels that SEROQUEL may possibly be responsible for the development of diabetes.

Follow-up 11 November 1999: Physician reports that the 47 year old female (not 45) had been receiving SEROQUEL 600 mg daily for schizoaffective disorder for a total of 12 months and experienced a severe 50 pound weight gain (date of onset unknown with no improvement). The patient was hospitalized in June 1999 due to the development of severe diabetes mellitus (difficult to control hyperglycemic). The patient is now receiving insulin and although the event continues, it has improved. SEROQUEL was tapered for discontinuation. Concomitant medications include klonopin and benadryl. The patient has a medical history of Hepatitis C, hypertension and arthritis. Physician states that he believes "SEROQUEL caused the weight gain which brought out diabetes in this patient likely predisposed for this condition."

Case Number: 1999UW03387

TYPE II DIABETES, DROWSINESS

A report has been received from a physician concerning a 17 year old Hispanic male patient who had been receiving SEROQUEL 100 mg every evening since January 1999 for psychotic depression and auditory hallucinations. In March 1999, due to drowsiness in the daytime, the dosage of SEROQUEL was decreased to 50 mg every evening. In July 1999, the patient was diagnosed with Type II diabetes. On 11 September 1999, SEROQUEL dosage was again decreased to 25 mg every evening. The patient had been receiving risperidol prior to

SEROQUEL. Concomitant medications include ritalin for attention disorder and serzone for depression.

Case Number: 1999UW00969

COMPLICATIONS OF DIABETES MELLITUS

A report has been received from a physician concerning a 28 year old male patient who was taking SEROQUEL and lithium (doses, duration, indications unknown). The patient presented to the emergency room following flu-like symptoms lasting for one week, as well as asthma, and weakness in the legs. There was no report of an increase or decrease in body temperature before presentation. His temperature at the emergency room was 107+ F (rectal probe 108 F), cardiac arrhythmias, liver enzymes were twice normal, blood glucose was 2240, potassium low, CPK normal, Lithium level was not elevated (0.4 or 0.6). The patient developed severe arrhythmias, ventricular fibrillation, and disseminated intravascular coagulation with low fibrinogen. He died at 4:00 a.m., on 14 March 1999. The tentative diagnosis is neuroleptic malignant syndrome. Complete autopsy reports are pending.

*Follow-up received 22 March 1999: A pharmacist reports that the patient started zithromax on 10 March 1999, to counter a suspected infection. She also reports that the patient had been bleeding from his eyes and nose.

*Follow-up received 14 March 1999: The patient presented on 14 March 1999 with focal twitching. He had increased tone, no doll's eyes or corneal responses. The pupils were mid-position and not reactive, and there was no reaction to noxious stimuli. Ativan was started, to a loading dose of 0.03 mg/kg. No further seizure activity was noted. The patient was started on Dantrium 2 mg/kg, with the dose of 150 mg. The patient developed cardiac arrhythmia requiring anti-arrhythmics. He continued to be acidotic and received bicarbonate and insulin. Split fibrins were ordered for the bleeding abnormalities, as well as whole blood. A pacemaker was placed and a rhythm obtained. The patient had been packed in ice from the onset. He died on 15 March 1999. The reporter attributed the event to a drug effect from multiple prescriptions.

*Follow-up received 05 May 1999: The county coroner reported the patient's cause of death as complications of diabetes mellitus. The patient did not have neuroleptic malignant syndrome. There had been a history of a 10 to 16 lb weight loss with flu-like symptoms, and blood glucose of 2240 on admission.

Case Number: 1999UW00967

DIABETES

A report has been received from a physician concerning a 17 year old male who is receiving SEROQUEL 200 mg twice daily for schizophrenia. The patient was initially started on 100 mg which was increased. While being hospitalized for his schizophrenia, routine lab work revealed diabetes which is being treated with glucochol 300 mg daily. Patient also receives paxil and depakote. Patient continues on SEROQUEL.

Case Number: 1999UW00288

BLOOD SUGAR RISING

A report has been received from a 58-year-old diabetic female patient who has been receiving SEROQUEL since September 1997. In 1994 she was diagnosed with diabetes mellitus. In 1997 her blood sugar readings began rising and on 20 January 1999 the reading was 321.

Case Number: 1999AP06660

LOSS OF DIABETIC CONTROL, TOOTH PAIN, INSOMNIA

A report has been received from a pharmacist concerning a 45 year old male patient who has been receiving SEROQUEL since April 1999 for treatment of schizophrenia. The patient began quetiapine therapy on 300 to 400 mg/day and increased to 750 mg/day in September/October 1999. For two years previously, the patient had a history non-insulin dependent diabetes mellitus. This was initially treated with metformin and then diet-controlled only until he started SEROQUEL in April 1999. After starting quetiapine therapy, the patient developed a loss of diabetic control, particularly on the higher dosage. Blood glucose which was previously stable at 10 (units unknown) rose to 13 or greater. He was treated with glibenclamide 7.5 mg/day. At the time of reporting the events were ongoing.

The reporter felt that the loss of diabetic control was related to quetiapine therapy due to the temporal relationship. It was noted that the patient had a history of non-insulin dependent diabetes mellitus that was previously diet controlled.

Case Number: 1999AP05757

DIABETES, KETOACIDOSIS.

A report has been received from a physician concerning a 25 year old male patient who has been receiving quetiapine fumarate 750 mg daily for psychosis since November 1997. He was receiving acamprosate, depixol and priadel concomitantly. In August 1999, 1 year 9 months after starting quetiapine fumarate, the patient was hospitalised due to the development of diabetes mellitus and ketoacidosis. It was also reported that he had experienced weight gain (date of onset and quantity of weight gained unknown). The patient is now being treated with insulin, has recovered with residual effects and quetiapine is continuing.

The reporter had no opinion regarding the causal relationship between the events and quetiapine fumarate, but commented that the weight gain may have been a contributing factor.

Case Number: 1999AP05218

DIABETES DURING PREGNANCY

Patient developed diabetes during pregnancy and started insulin on 30 September 1999. Baby due 06 December 1999, but patient's water broke 30 September 1999 and baby born in October 1999. See case 1999AP06076.

Case Number: 1999AP02989

DIABETES MELLITUS

This patient started treatment with SEROQUEL on 13 November 1998 and with fluoxetine on 12 November 1998. Urine and blood tests on 26 November 1998 indicated that she had developed diabetes mellitus. Urinary sugar was raised and blood sugar was 17.1.

Case Number: 1999AP01985

NON INSULIN DEPENDENT DIABETES

A physician reported that a 44 year male patient was given SEROQUEL 250 mg BID for resistant schizophrenia. Treatment began on 27 August 1998. Concomitant medication included clonazepam, sodium valproate and cyproterone. The patient had no history of diabetes mellitus and was being treated with cyproterone for a disorder of sexual inhibition. Five months after starting SEROQUEL, the patient developed non-insulin dependent diabetes. SEROQUEL was stopped toward the end of January 1999. No follow-up is available.

Case Number: 1998UW49554

CEREBROVASCULAR ACCIDENT, DIABETIC ACIDOSIS, TRANSIENT ISCHEMIC ATTACK, COLLAPSE.

A report has been received from a physician concerning a 58 year old male patient who received SEROQUEL 800 mg daily for schizoaffective disorder. The patient has a history of diabetes and cerebrovascular accident. He also takes gabapentin. The patient experienced a transient ischemic attack and was unresponsive except for painful stimuli. Five minutes later, the patient recovered fully. The following day, he collapsed in the shower and died. An autopsy was performed and the primary cause of death was listed as diabetic acidosis and secondary to cerebrovascular accident. Additional information will be requested.

Case Number: 1998UW49081

HYPERGLYCAEMIA

Patient is an 83 year old female who was admitted to the hospital on 27 September 1998 with a diagnosis of hyperglycaemia. Past history and medical conditions include diabetes mellitus. The first patient completed the double-blind portion of the trial on 14 September 1998. Open label medication started on 14 September 1998 and ended on 26 September 1998. This event took place on day 12 of study medication at a dose of 25 mg. In the opinion of the investigator, the elevated blood sugar was not related to the study medication.

Case Number: 1998UW48844

HYPERGLYCAEMIA, DIABETES.

A report has been received from a physician concerning a male patient in his early forties who has been receiving SEROQUEL for four weeks and is experiencing hyperglycaemia and diabetes. The patient, who has no previous history of diabetes, is now showing blood sugars of over 600 mg/dl.

NEW ONSET DIABETES MELLITUS ASSOCIATED WITH THE INITIATION OF QUETIAPINE TREATMENT, J OF CLINICAL PSYCHIATRY, 60: 556-557, AUG 99, USA, SOBEL, M., JAGGERS, ED, FRANZ, MA

Case Number: 1998UW48512

DIABETES MELLITUS

A report has been received from a pharmacist concerning a 42-year-old male patient who has taken SEROQUEL since July 1998. On 31 August 1998 he was diagnosed with diabetes mellitus. Further information is being requested.

*Follow up 12 October 1999: Further information reveals that the patient was receiving SEROQUEL 200 mg for a bipolar disorder since July 1998. On 31 August 1999, patient was admitted to the hospital with a new-onset diabetes mellitus. Patient had no prior history of glucose intolerance or hyperglycaemia. Four months prior to admission blood glucose was 126 mg/dL and 107 mg/dL. At admission blood glucose was 607 mg/dL. SEROQUEL was tapered, then discontinued. Insulin requirements decreased markedly and insulin was eventually discontinued in January 1999. Patient was also receiving lithium carbonate, gabapentin, clonazepam, and venlafaxine.

SOBEL M, JAGGERS ED, FRANZ MA: NEW-ONSET DIABETES MELLITUS... J OF CLIN PSYCHIATRY; 1999;60(8):556-557.

Case Number: 1998AP50408

HYPERGLYCAEMIA (NON-SERIOUS)

A pharmacist and a nurse reported that a male patient taking SEROQUEL developed hyperlycaemia. The pharmacist considered the event unrelated to SEROQUEL; the nurse considered the event related to SEROQUEL. The patient was also taking stelazine.

Case Number: 1998AP45979

LOSS OF DIABETIC CONTROL, AGGRESSIVE BEHAVIOUR, STROPPY BEHAVIOUR

Patient had actually been messing about with his insulin injections that weekend, the event therefore had nothing to do with SEROQUEL, SEROQUEL dosage has been reduced from 400 to 200mg. The physician is thinking of stopping SEROQUEL altogether.

Case Number: 1998AP18089

HYPERGLYCAEMIA.

A report has been received from a physician concerning a 32 year old male patient who has been receiving SEROQUEL from 21 May 1995 for psychosis as part of a clinical trial. The patient has a medical history of obesity, abdominal pain, indigestion, constipation, muscle stiffness, restlessness, depression, and hypertension. He was also taking valproate semisodium, benztropine mesylate and propranolol.

On 26 January 1998, 2 years 36 weeks after starting study medication, the patient was found to have hyperglycaemia and was hospitalised. At the time this report was received, the event was still ongoing. The study drug was stopped on 01 February 1998 due to the potential effect of unstable glucose levels on the patient's mood. The reporter considered that there was not a reasonable possibility that this event was related to the study therapy.

Case Number: 1997AP36803

DIABETIC KETOACIDOSIS

A report has been received from a physician concerning a 36 year old male who has been receiving SEROQUEL in a dose up to 500 mg daily for schizophrenia as part of a clinical trial. SEROQUEL started on 06 September 1996. The patient had recently been diagnosed with diabetes mellitus which was controlled on glucotrol. On 18 March 1997, 28 weeks after starting SEROQUEL, he was admitted to hospital with decreased level of consciousness. He had not been taking his glucotrol or SEROQUEL for 3 to 4 days prior to admission. He was given IV fluids and insulin but later developed severe acidosis and an increased lipase of 1819 u/l(25-229)and amylase of 135u/l(27-92). Other abnormal laboratory findings were:

sodium 130 mmol/l (135-146), chloride 99 mmol/l (100-107), bicarbonate 5mmol/l (22-32), creatinine 1.9 mg/dl (0.4-1.4), glucose 413mg/dl (70-160), uric acid 12.3mg/dl (2.2-7.2), White blood count 17,000 (4,000-11000), beta-hydroxy butyrate 182mg/dl(0.4-4). The patient was started on subcutaneous insulin and food was started once amylase and lipase were within normal range. At the time of reporting the patient had not restarted SEROQUEL. The event resolved on 01 April 1997. The investigator felt that there was not a reasonable possibility that the event was related to SEROQUEL.

Case Number: 1997AP36246

UNCONTROLLED DIABETES

A report has been received from a physician concerning a 29-year old male who has been receiving SEROQUEL since 22 January 1997 in a clinical trial for schizophrenia. After 8 months treatment, the patient was attending a hospital trial visit on 23 September 1997 when he felt faint and collapsed. He was found to have elevated blood glucose, decreased blood pressure (70/50) and an abnormal ECG with cardiac enzymes raised. SEROQUEL treatment was put on hold and the patients diabetes treated with humulin in Hospital. The event was ongoing at the time of the report. The physician felt that there was not a reasonable possibility that this event was related to the SEROQUEL therapy.

Case number: 1997AP35710

UNCONTROLLED DIABETES MELLITUS

A report has been received regarding a 45 year old male who has been receiving SEROQUEL as part of a clinical trial. He has a medical history of diabetes mellitus, insomnia, gonorrhoea, genital herpes, alcohol and heroin abuse. His concomitant medications were clonazepam, amitriptyline, famotidine and lisinopril. On 10 August 1997, 163 days after starting SEROQUEL, he had a moderately severe episode of uncontrolled diabetes mellitus requiring hospital treatment. He recovered after IV fluids and a 2200 calorie diabetic diet. He remains in the trial.

The investigator considered the event not related to trial therapy.

Case Number: 1996AP19874

PNEUMONIA, DIABETES, HYPERTENSION

This 65-year old male patient with Parkinsons disease, anaemia of chronic disease, obsessive compulsive disorder, penile implant, and peptic ulcer disease was being treated with SEROQUEL as part of a clinical trial. The patient was receiving gastric tube nutrition secondary to poor gag reflex. Treatment began on 21 September 1995. Earlier in the year the patient had been hospitalised suffering from pneumonia. On 28 March 1995, the patient complained of chest congestion. X-ray confirmed that he had pneumonia. He was treated with antibiotic in his nursing home but was later admitted to hospital for further antibiotic treatment. During his admission, he was noted to have elevated blood sugar and blood pressure. Discharge diagnoses were right lower lobe pneumonia, possible nasotracheal aspiration, new onset diabetes and hypertension. The diabetes and hypertension were considered to be not regulatory serious and not related to trial therapy.

The investigator considered the pneumonia was not related to trial therapy.

Case Number: 1995AP10737

DIABETES MELLITUS

This 52 year-old-female with schizophrenia was taking SEROQUEL 400 mg from 28 January 1995 as part of a clinical trial. On 31 January 1995 this patient was hospitalised with diabetes mellitus. She was not withdrawn from the trial. When first reported 3rd April 1995, this event was considered probably not related. However, further information now reveals that elevated sugar levels have been detected in this patient for two years. Therefore it is considered that her diabetes was definitely not related to the study medication.

This event is now regarded as non-serious by the investigator as it was symptoms of the patient's schizophrenia which led to prolonged hospitalisation and not the diabetes.

Case Number: 1994AP04544

AGITATION, UNREST, INCOMMUNICATIVE, DISINHIBITION, PARANOIA, DIABETES, INCREASED TRIGLYCERIDES

Patient with impaired glucose metabolism pre-trial. Entered in SEROQUEL trial on 26 September. On study day 8 this patient developed an acute psychosis, suggesting lack of efficacy, which led to withdrawal from the trial. On 4 November, the patient developed

symptoms of diabetes. Physician assessment is that there is no reason to suspect that development of diabetes is related to treatment with SEROQUEL.

Case Number: 1994AP03286

HYPERGLYCAEMIA

An investigator reported that a 53 year old female patient started taking SEROQUEL on 22 July 1994. The patient had a history of insulin-treated diabetes and had been taking several concomitant medications. On 8 August 1994, the patient was noted to be hyperglycaemic. The investigator reported that the patient had the same level of hyperglycaemia that she had prior to study entry.

Case Number: 1994AP00893

HYPERGLYCAEMIA

An investigator reported that a 45 year old male was treated with SEROQUEL beginning on 4 March 1994. Concomitant medications included zantac and haldol. The patient had no history of diabetes mellitus. He had recently stopped taking an unblinded SEROQUEL study drug. On 3 March, the fasting blood sugar was 393. The following day, it rose slightly before increasing to 1104 on 13 March. SEROQUEL was stopped that day. No treatment was reported but the blood glucose on 14 March was 200.

5. DISCUSSION

There were 27 reports of diabetes mellitus and 2 reports of hyperglycaemia received by AstraZeneca to date. New onset diabetes mellitus was described in 19 of these 27 reports and exacerbation of pre-existing diabetes mellitus accounted for 8 reports. Four reports described patients who developed diabetic ketoacidosis (2000UW01164, 1999AP05757, 1998UW49554, and 1997AP36803). Two of these were new onset reports and the other two involved worsening of pre-existing diabetes mellitus. There have been no reported cases of non-ketotic hyperosmolar coma received to date. Of these total 28 reports, 16 were spontaneous reports, 10 were from clinical trials, and 2 were literature reports. The investigator attributed none of the cases reported from clinical trials to SEROQUEL.

New onset diabetes mellitus: There have been 19 cases of new onset diabetes mellitus reported to date. The age range for patients with new onset diabetes mellitus is 12 to 65 with an average age at onset of 37.5 years (median = 41 years). There is a male predominance with males constituting 74% of all reports. Daily SEROQUEL dosages ranged from 50 mg to 800 mg, with an average daily dose of 419 mg (median = 400 mg). The average time interval between initial therapy and the date of the reported event was 6.2 months with a range of 3 days to 27 months

(median = 2.5 months). Two patients with new onset diabetes mellitus also developed diabetic ketoacidosis (2000UW01164 and 1999AP05757). The former patient reportedly lost 30 pounds. Blood glucose concentrations at clinical presentation averaged 833 mg/dl and ranged from 311 to 2240 mg/dl.

Two patients who developed new onset diabetes mellitus also experienced weight gain (1999UW03532 and 1999AP05757). The latter patient also had diabetic ketoacidosis. One patient who developed hyperglycaemia also experienced weight gain (2000UW01047). Weight gain was not reported in any other cases.

Two patients with new onset diabetes mellitus experienced dose related loss of glycemic control as reported by their physicians (1999UW00969 and 1998UW48512).

Diabetic ketoacidosis: There have been 4 cases of diabetic ketoacidosis reported to date all involving males. The age range for patients with diabetic ketoacidosis is 25 to 58 with an average age at onset of 40.5 years. Daily SEROQUEL dosages ranged from 200 mg to 800 mg, with an average daily dose of 562 mg (median = 625 mg). The average time interval between initial therapy and the date of the reported event was 9.7 months with a range of 1 to 21 months. Two patients with new onset diabetes mellitus also developed diabetic ketoacidosis (2000UW01164 and 1999AP05757). The former patient reportedly lost 30 pounds. Only one case (1997AP36803) reported the blood glucose concentration at clinical presentation, which was 413 mg/dl. One patient died of complications of diabetes mellitus (1998UW49554). A second patient (1997AP36803) recently diagnosed with diabetes mellitus, discontinued taking his oral hypoglycemic agent three days before being hospitalized with DKA. A third patient (1999AP05757) with new onset diabetes mellitus also experienced weight gain (unspecified) and at last word required insulin therapy.

Non-ketotic hyperosmolar coma: There have been no reported cases of non-ketotic hyperosmolar coma.

Hyperglycaemia: There have been two reports of hyperglycaemia reported to date (2000UW01047 and 1998AP50408). Blood glucose concentrations were not provided for either patient. The former report involved a 47-year-old female who developed weight gain and hyperglycaemia after taking SEROQUEL 150 mg daily for 30 months. The latter report contains scant information, except the daily SEROQUEL dose which was 750 mg.

Confounding factors: Few, if any, of these patients had baseline fasting glucose levels. Seven patients with new onset diabetes mellitus were using concomitant medications known to impair glucose tolerance and cause diabetes mellitus including risperidone, fluoxetine, albuterol, and venlafaxine (2000UW01164, 1999UW03387, 1999AP02989, 1999AP05218, 1998UW48512, 1999UW00969, and 1998AP18089). This last patient was also reportedly obese. One patient

developed Type 1 diabetes mellitus (2000UW00266). Several reports contained only scant information which precluded detailed analysis of these cases.

While there were no reports of positive dechallenges and rechallenges, consideration should be given to the suggestion that SEROQUEL therapy may cause impaired glucose regulation including diabetes mellitus in certain individuals.

6 REFERENCES

- (1) Electronic Medicines Compendium: <http://emc.vhn.net>: accessed June 5, 2000.
- (2) American Diabetes Association: Clinical Practice Recommendations 2000, Volume 23 Supplement 1, Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus
- (3) Foster D. Diabetes mellitus. In: Fauci AS et al, editors. Harrison's Principles of Internal Medicine, 14th Edition. Philadelphia: McGraw-Hill, 1998: 2060-80
- (4) Wilson DR, D'Souza L, Sarkar N, Newton M. New-onset diabetes and ketoacidosis with atypical antipsychotics, American College of Neuropsychopharmacology, 1999