


A Critical Curriculum on Psychotropic Medications




A Critical Curriculum on Psychotropic Medications

- **Principal Investigator:** - David Cohen, Ph.D.
- **Research Coordinator:** - Inge Sengelmann, M.S.W.
- **Professional Consultants:**
 - David O. Antonuccio, Ph.D. (psychology)
 - Kia J. Bentley, Ph.D. (social work)
 - R. Elliott Ingersoll, Ph.D. (counseling & psychology)
 - Stefan P. Kruszewski, M.D. (psychiatry)
 - Robert E. Rosen, J.D., Ph.D. (law)
- **Flash production and design:**
 - Sane Development, Inc., and Cooper Design, Inc.
- **Voice narration and Flash editing:**
 - Saul McClintock




CriticalThinkRx was made possible by a grant from the Attorneys General Consumer and Prescriber Grant Program, funded by the multi-state settlement of consumer fraud claims regarding the marketing of the prescription drug Neurontin®




Module 1

Why a Critical Skills Curriculum on Psychotropic Medications?



Part A


Curriculum Rationale, Funding and Contents



Curriculum Rationale

Physicians write prescriptions, but other professionals often influence who gets prescribed and why

Training for these professionals is mostly haphazard and often influenced by the pharmaceutical industry



6

Curriculum Objectives

Help practitioners in mental health and child welfare sharpen *critical thinking skills* to deal with complex and evolving issues about psychotropic medication

7

Critical thinking

- ✓ involves assessing beliefs, arguments and claims to arrive at well-reasoned judgments
- ✓ uses standards such as clarity, accuracy, relevance, and completeness

8

Critical thinking

- ✓ asks “who benefits?”
- ✓ is sensitive to the influence of vested interests on information
- ✓ emphasizes the ethical implications of treatment decisions

9

CriticalThinkRx

A prescription for critical thinking about psychotropic medications

10

Curriculum funding

- Received from the *Attorneys General Consumer & Prescriber Education Grant Program (CPGP)*
- CPGP is overseen by the Attorney General offices of Florida, New York, Ohio, Oregon, Texas and Vermont (plus two rotating states)

11

Funding source of CPGP

2003: Attorneys General of 50 states charged Warner Lambert, a subsidiary of Pfizer, Inc., with conducting an unlawful marketing campaign promoting the off-label uses of the anticonvulsant drug Neurontin

12

Neurontin settlements

2004: The company settled for \$430 million

- \$21 million was earmarked for research and education aimed at health professionals

13

CPGP awards grants

2006: CPGP funded 28 applications in 19 states

- *CriticalThinkRx*, funded at Florida International University, is the only project targeting non-medically trained professionals in child welfare and mental health

14

CPGP aims to improve prescribing practices by educating health professionals about

- ✓the drug development and approval process
- ✓pharmaceutical industry marketing
- ✓knowledge and skills to evaluate drug information critically

15

CPGP requires that

- ✓the curriculum be maintained in the public domain, freely accessible by anyone
- ✓the investigators and their consultants forego funding from the pharmaceutical industry for the duration of their grants

16

Selection of content

Systematic literature searches were conducted in 2006-2007 on databases in medicine, pharmacology, public health, social work, counseling, and psychology

- Materials were selected based on relevance and accuracy

17

Mainstream views

Researchers agree that clinical practice has far outpaced the empirical evidence, yet...

- Mainstream mental health practice subscribes to a “medical” model supporting medication of children with little evidence of safety or efficacy



18

Content bias

CriticalThinkRx offers alternative views based on empirical evidence to stimulate critical thinking and a more balanced evaluation based on ethical codes of practice

19

Content orientation

CriticalThinkRx emphasizes the ethical dictate: **“First, do no harm”**

CriticalThinkRx tries to close gaps between research and practice to maximize opportunities to help clients and avoid harm

20


Curriculum design

Modules designed by experienced researcher/clinician with input from independent consultants in counseling, psychology, psychiatry, social work, and law

21

Principal Investigator

David Cohen, Ph.D., L.C.S.W.



- Professor of Social Work, Florida International University, Miami, and a private practitioner
- Author of numerous publications on psychiatric drugs, medicalization, and law and psychiatry
- His latest books are *Your Drug May Be Your Problem* (2nd rev. ed, 2007) and *Critical New Perspectives on ADHD* (2006)

22

Research Coordinator

Inge Sengelmann, M.S.W.

- M.S.W. with a background in journalism and corporate communication
- Clinician focused on holistic approaches to the treatment of trauma-related mood and behavioral problems



23

Consultant: Counseling

R. Elliott Ingersoll, Ph.D.

- Professor of Counseling, College of Education and Human Services, Cleveland State University
- A licensed psychologist and clinical counselor in Ohio, he has authored books, book chapters, and articles on psychopharmacology, spiritual approaches to counseling, and Integral theory in mental health
- Author, *Psychopharmacology for Helping Professionals: An Integral Exploration* (2006)



24

Consultant: Social Work

Kia J. Bentley, Ph.D., L.C.S.W.



- Professor, Director of the Ph.D. Program, and Associate Dean for Strategic Initiatives in Social Work at Virginia Commonwealth University, where she has taught since 1989
- Author, *The Social Worker & Psychotropic Medication* (3rd ed., 2006) (with Joseph Walsh)
- Editor, *Psychiatric Medication Issues for Social Workers, Counselors and Psychologists* (2003)

25

Consultant: Psychology

David O. Antonuccio, Ph.D.



- Professor, Department of Psychiatry, University of Nevada School of Medicine
- Fellow, American Psychological Association; Diplomate, clinical psychology, American Board of Professional Psychology
- His articles on the comparative effects of psychotherapy and pharmacotherapy have received extensive national coverage and are models of careful scholarship
- Has received many prestigious awards for his outstanding contributions to clinical science and research

26

Consultant: Psychiatry

Stefan P. Kruszewski, M.D.



- Harvard Medical School graduate and board-certified in adolescent psychiatry
- Pennsylvania-based clinician and scientist working with U.S. and international judicial, legislative, and regulatory bodies
- His publications appear in *American Journal of Psychiatry* and *BMJ*

27

Consultant: Law

Robert E. Rosen, J.D., Ph.D.



- Professor of Law, University of Miami, Coral Gables, FL
- Has taught courses in children and the law, professional responsibility, and sociology and the law
- Has served as member of Miami-Dade's Community-Based Care Alliance, and is a reviewer for Foster Care Review
- Holds a J.D. from Harvard Law School, and a Ph.D. in sociology from the University of California at Berkeley
- Former fellow, Harvard's Program in Ethics and the Professions

Use of drug names

Most prescription drugs have a generic and a brand name (e.g., fluoxetine/Prozac)

In this course, charts show both names, but discussions use brand names because they are more familiar to laypersons

29



30

A recent tragic case raises questions about the use of psychiatric medications in young children

31

Case 1: Rebecca Riley

(April 11, 2002 - Dec. 13, 2006)

What went wrong?

Concerns raised before death of 4-year-old girl

Teachers, a social worker, a nurse and a therapist had noticed something wasn't right about Rebecca Riley. A



© Copyright 2007 The Patriot Ledger

32

CBS NEWS Oct. 4, 2007 12:30pm
 CBS EVENING NEWS THE EARLY SHOW 48 HOURS MYSTERY 30 MINUTES
60 MINUTES

What Killed Rebecca Riley?

Katie Couric Reports On The Diagnosis Of Bipolar Disorder In Kids
 Sept. 30, 2007

Girl's death stirs debate over psychiatric meds

Parents of 4-year-old accused of intentionally overmedicating daughter

AP Associated Press
 Updated: 3:11 p.m. ET March 29, 2007

HULL, Mass. - In the final months of Rebecca Riley's life, a school nurse said the little girl was so weak she was like a "floppy doll."

The preschool principal had to help Rebecca off the bus because the 4-year-old was shaking so badly.

And a pharmacist complained that Rebecca's mother kept coming up with excuses for why her daughter needed more and more medication.

33

Some salient facts

In 2002, then again in 2005-2006, Massachusetts' DSS investigated complaints that the three Riley children might be sexually or physically abused and neglected by their parents

DSS ruled complaints unfounded

34

By 2006, all three Riley children were diagnosed with Bipolar I Disorder and prescribed psychotropic drug cocktails by same child psychiatrist from Tufts Medical Center

- Parents were also diagnosed and mother received Paxil
- As discussed in next modules, diagnosing children with Bipolar Disorder I is a questionable and controversial practice

35

Rebecca, the youngest child, was first medicated at age 2

- By age 4, she was taking Seroquel (antipsychotic), Depakote (anticonvulsant), and clonidine (antihypertensive)
- She also took 2 over-the-counter cold medicines

36

Dec. 13, 2006: Rebecca Riley is found dead on her parents' bedroom floor


- Autopsy later indicated cause of death as **“intoxication due to the combined effects”** of clonidine, Depakote, and two cough medications
- **“The amount of clonidine alone in Rebecca’s system was fatal.”**

(Commonwealth of Massachusetts, Feb. 5, 2007) 37

Parents indicted ...

Michael Riley, 34, and Carolyn Riley, 32, indicted in 2007 for the 1st degree murder of their daughter Rebecca (charge later reduced to 2nd degree murder)

- Parents charged with giving her “excessive amounts” of clonidine
- Child’s doctor told mother Rebecca “was already on a high dose of clonidine” and a higher dose could kill the child



© Copyright 2007 The Patriot Ledger

(Commonwealth of Massachusetts, Feb. 5, 2007) 38

Case leads to resignations...

GOODBYE TO DSS CHIEF

Agency has been under fire since parents accused of killing Hull girl

By KEN MAGUIRE Associated Press

BOSTON - The embattled head of the state’s child welfare system is resigning five months after his agency was criticized for its action - or lack of action - in the death of a 4-year-old girl in Hull.



Lewis “Harry” Spence, commissioner of the Department of Social Services since 2001, has been under fire for the agency’s handling of the Hull case in which the parents of the dead girl are charged with killing her with an overdose of prescription drugs.

He also has been criticized for the department’s handling of another high-profile child-abuse case involving a comatose child from Westfield.

Gov. Deval Patrick plans to replace Spence with Angelo McClain, a former DSS worker who now works for ValueOptions, a New Jersey-based health care company, according to a person with direct knowledge of the decision.

Patrick planned to announce Spence’s departure today along with other changes in top positions, a source told The Associated Press.

Spence did not return calls to his cell phone seeking comment.

Lewis “Harry” Spence

39

... puts careers on the line

February, 2007

Psychiatrist to suspend practice; denies wrongdoing The Boston Globe

By Liz Kowalczyk, Globe Staff | February 8, 2007

Dr. Kayoko Kifuji, the psychiatrist who treated Rebecca Riley in the months before the Hull girl died from an overdose of prescription drugs, agreed yesterday to immediately stop treating patients while the state investigates her role in the case.

April, 2008

HOME / NEWS / LOCAL

Doctor is sued in death of girl, 4 The Boston Globe

Her psychiatrist treated her with powerful drugs

By Shelley Murphy
Globe Staff / April 4, 2008

Email | Print | Single Page | Text size - +

40

CBS EVENING NEWS

March 10, 2007

(CBS) Rebecca Riley’s death shocked the Boston community. Did her parents deliberately give her overdoses of psychiatric drugs as prosecutors suggest? Or are her doctors to blame — as defense lawyers argue — for prescribing powerful medications when she was just 2 years

41

Girl’s pill numbers disputed: The prescriptions Carolyn Riley gave 4-year-old were very close to allowed amount, defense says

By JULIE JETTE
The Patriot Ledger



The Patriot Ledger

March 10, 2007

42

Case shines light on therapists' roles...

An LCSW made 12 home visits in summer 2006, working with Rebecca and her 6-year-old sister

- Therapist was "initially concerned" about the medication regimen, since she "did not observe any behavior consistent with the diagnoses"

(Commonwealth of Massachusetts, Feb. 5, 2007) 43

... and on school personnel

In her pre-school, Rebecca was observed to be very lethargic and have "a tremor in her hand"

Mother was observed to be "lethargic" and "fall asleep during interviews"

44

Case stirs heated debate among doctors over bipolar diagnoses

The Boston Globe

Backlash on bipolar diagnoses in children
MGH psychiatrist's work stirs debate

By Scott Allen, Globe Staff | June 17, 2007

45

Leads one doctor to hold another "morally culpable"

LAWRENCE DILLER *The Boston Globe*
Misguided standards of care
By Lawrence Diller | June 19, 2007

"... I felt compelled to name Joseph Biederman, head of the Massachusetts General Hospital's Pediatric Psychopharmacology clinic, as morally culpable in providing the 'science' that allowed Rebecca to die."

-- Lawrence Diller, M.D.

46

FDA "black box" warnings on Depakote ignored?

FDA-approved Depakote black box warning label:

"HEPATOTOXICITY: HEPATIC FAILURE RESULTING IN FATALITIES HAS OCCURRED IN PATIENTS RECEIVING VALPROIC ACID AND ITS DERIVATIVES. EXPERIENCE HAS INDICATED THAT CHILDREN UNDER THE AGE OF TWO YEARS ARE AT A CONSIDERABLY INCREASED RISK OF DEVELOPING FATAL HEPATOTOXICITY...."

"PANCREATITIS: CASES OF LIFE-THREATENING PANCREATITIS HAVE BEEN REPORTED IN BOTH CHILDREN AND ADULTS RECEIVING VALPROATE. SOME OF THE CASES HAVE BEEN DESCRIBED AS HEMORRHAGIC WITH A RAPID PROGRESSION FROM INITIAL SYMPTOMS TO DEATH."

47

Case 2:

"Susan," 10 years old

Parents divorced 5 years ago, custody awarded to mother

Father seeking shared custody—only sees Susan a few times a year

Susan presented behavior problems since the age of 3

48

Loss and instability

Susan's life filled with losses of friends, pets, homes, adopted-away brother

Since age 5, Susan moved 10 times, attended 7 schools, was assessed by 20 physicians and therapists

49

Multiple diagnoses

Diagnosed with ADHD, OCD, bipolar disorder

Lives in a residential treatment center

Her file describes many behavioral outbursts, attributed to "bipolar disorder"

50

Since age 5, Susan has taken:

- ✓5 antipsychotics
- ✓4 anticonvulsants
- ✓3 stimulants
- ✓3 antidepressants
- ✓2 benzodiazepines
- ✓2 other sedatives (incl. antihypertensive)
- ✓lithium

51

Susan now takes:

- ✓2 anticonvulsants
- ✓1 antipsychotic
- ✓1 stimulant, and
- ✓1 antihypertensive



No evaluations of medication...


A psychologist and a social worker conducted separate assessments of Susan's situation for the Court

Neither commented on Susan's drug treatment or suggested any connections between the medications and her behavioral outbursts


53

No one expressed any concern about giving 5 psychiatric drugs (including 4 central nervous system depressants) to a 10-year-old

54



- What are the client's symptoms or observed behaviors of concern, who has observed them?
 - Has the client experienced any recent or chronic life events or stressors that may contribute to the problems?
- 57

- Could any of client's problems be caused by current medication?
- 
- 58

- Does the client's psychiatric diagnosis truly reflect the client's problems? Is the diagnosis useful to plan for interventions with this client?
 - What interventions have been tried to address client's problems? By whom, and with what results?
 - Are alternative interventions available to address client's problems? Why have they not yet been tried?
- 59

- Why is medication being prescribed for this client? What other medication has been prescribed currently or in the past?
 - How long before we see improvements? How will the improvements be measured?
 - How long will the patient be on the medication? How will a decision to stop be made?
- 60

- If client is a minor, is the medication designed to benefit the child, or the child's caregivers?



61



62

- Why is this particular medication prescribed for this client?
- How long has it been on the market? Is it FDA-approved for use in children? Are there any FDA "black box" warnings about this medication?
- What is the recommended dosage? How often will the medication be taken? Who will administer it?

63

- Have any studies been evaluated by professionals working with this child?
- How much scientific support is there for its helpfulness with other children with similar conditions?
- How much scientific evidence exists to support safety and efficacy of this drug in children, alone or in combination with other psychotropic medications?

64

- Has this medication been shown to induce tolerance and/or dependence? What withdrawal effects may be expected when it is discontinued?

65

- Do any laboratory tests need to be done before, during, after use of this medication?
- Are there other medications or foods the child should avoid while on this medication?
- What are all the potential positive and adverse effects of this medication?

66

- How will the effects of the medication be monitored? By whom? Where will they be documented? What should be done if a problem develops?
- How will the use of medication impact other interventions being provided?

67

- How much does this medication cost and who is paying for it?
- Are there cheaper, generic versions of this medication?



68



69

- What is the experience of the physician prescribing the medication?
- Would you consider the physician's prescribing habits as cautious and conservative?
- Does this physician have any financial relationships with pharmaceutical companies? Have these been disclosed to patients?

70

- Have all the risks and benefits of this medication, and those of alternate interventions, been evaluated and discussed by the physician with the client or the client's family?
- Is there an adequate monitoring schedule and follow-up?

71

- Do I or my client/client's family have the opportunity to speak regularly with the physician and other healthcare providers about the medication's effects? Should my feedback be expressed in writing?

72



- Has a comprehensive assessment (e.g., biopsychosocial, holistic, integral) been conducted? Does it offer plausible reasons for the client's problems?
- Are there other explanations for the child's behavior?

74

- If necessary, do I have access to supervision to help me think through the medication issues?
- How knowledgeable is my supervisor about psychotropic medications?

75

- Am I familiar with all the risks and benefits of this medication, as well as those of alternate interventions? Have I discussed them with the client/client's family?
- Do I know how the client/client's family feel about the use of medication?

76

- What is my role and has it been clearly delineated with all other providers?
- Has the client/client's family been provided with all the information necessary to provide informed consent? Do they understand their choices?

77

- Do I feel confident that I can recognize the effects, adverse or otherwise, of this medication on my client? How should I record my observations?
- Will I be able to educate my client about these effects so he/she can raise concerns with the prescribing physician?

78

- What alternative services/interventions does this family need or want?
- Can I provide these or help them obtain access?

79

This course, in the remaining modules, is intended to help you answer the preceding questions

80


A Critical Curriculum
on Psychotropic Medications

Module 1


The End



81




A Critical Curriculum on Psychotropic Medications




A Critical Curriculum on Psychotropic Medications

- **Principal Investigator:**
 - David Cohen, Ph.D.
- **Research Coordinator:**
 - Inge Sengelmann, M.S.W.
- **Professional Consultants:**
 - David O. Antonuccio, Ph.D. (psychology)
 - Kia J. Bentley, Ph.D. (social work)
 - R. Elliott Ingersoll, Ph.D. (counseling & psychology)
 - Stefan P. Kruszewski, M.D. (psychiatry)
 - Robert E. Rosen, J.D., Ph.D. (law)
- **Flash production and design:**
 - Sane Development, Inc., and Cooper Design, Inc.
- **Voice narration and Flash editing:**
 - Saul McClintock





CriticalThinkRx was made possible by a grant from the Attorneys General Consumer and Prescriber Grant Program, funded by the multi-state settlement of consumer fraud claims regarding the marketing of the prescription drug Neurontin®



Module 2


Increasing Use of Psychotropics

Public Health Concerns





Part A

Medicating Youth



Surveys and insurance databases show increasing use



5-8 million children in the U.S. (8-11% of all children) receive prescriptions for psychotropic medications

(Medco, 2006; St. Luke's Health Initiatives, 2006)

6

Prescriptions of psychotropics to youths tripled in the 1990s and are still rising in this decade

In some drug classes, rates in children rival adult rates



(Olson et al. 2002, 2006; Thomas et al. 2006; Zito et al. 2000, 2002, 2003)

7

Drug treatment without any other form of therapy is becoming the norm



(Olson et al. 2002, 2006; Thomas et al. 2006; Zito et al. 2000, 2002, 2003)

8

A worldwide phenomenon...



...but the proportion of children prescribed psychiatric drugs remains 2 to 20 times higher in the U.S., Canada, and Australia than in other developed nations

(Wong et al. 2004)

9

In the U.S., “cultural” differences remain



White children are twice as likely as Black and Latino children to receive prescriptions

- Difference appears unrelated to socio-demographic, access, or clinical factors, and may relate to parental attitudes

(Cooper et al. 2006; Dos Reis et al. 2005; Leslie et al. 2003)

10

Off-Label Uses and Polypharmacy



The New York Times

November 23, 2006

Proof Is Scant on Psychiatric Drug Mix for Young

11

“Off-label” use common

The practice of administering medications for indications or age groups not approved by the FDA, as indicated on the drug’s “label”



(Vitiello, 2001; Zito et al. 2003)

12

75% of all medication use in children is off-label



(Vitiello, 2001; Zito et al. 2003)

13

Concerns about off-label use


“Bearing in mind that some off-label use is perfectly justifiable, it is fair to say that much of it is not justifiable. If there is not evidence presented to the FDA about a given indication, it is certainly a user-beware situation.”



- Jerry Avorn, M.D., Professor of Pharmacology, Harvard Medical School, and author, *Powerful Medicines* (2005)

14

Polypharmacy common




40% or more of all psychiatric drug treatments today involve polypharmacy

(Bhatara et al. 2004; Olsson et al. 2002; Safer et al. 2003)

15


Polypharmacy: concomitant or multiple psychotropic medication use



16

Concomitant = ≥ 2 drugs taken on the same day

Multiple = ≥ 2 drugs taken during a given period



17

Concerns about polypharmacy

Basic empirical support of efficacy in children is lacking for *most individual* medication classes

No studies have established the safety and efficacy of combination treatments in children

(Bhatara et al. 2004; Jensen et al. 1999; Martin et al. 2002; Vitiello, 2001)

18

Increases behavioral toxicity

Behavioral toxicity = drug-induced adverse effects and behavioral changes, including apathy, agitation, aggression, mania, suicidal ideation and psychosis

(Safer, Zito & dosReis, 2003)

19

The “prescribing cascade”

Adverse effects are often confused with symptoms of disorders, leading to co-morbid diagnoses, and even more complex drug regimens



(Safer, Zito & dosReis, 2003)

20

Examples of behavioral toxicity

TABLE 4. Single Case Reports of Adverse Drug Events Associated With Use of Concomitant Psychotropic Medication for Youths

Study	Medications	Diagnosis	Age	Gender	Adverse Drug Events
Salice et al. (89)	Fluoxetine, promethazine, methylphenidate, clonidine	ADHD, conduct disorder, and Tourette's syndrome	9	Male	Death
Preda et al. (90)	Perphenazine, carbamazepine, valproic acid, lorazepam, thioridazine	Bipolar I disorder and adjustment disorder	10	Female	Psychosis
Budman et al. (91)	Pemoline, paroxetine, haloperidol	OCD, ADHD, and Tourette's syndrome	12	Male	Acute dystonia
Levy et al. (92)	Amikrophiline, fluoxetine, clonidine	ADHD and conduct disorder	7	Male	Serotonin syndrome
Fisman et al. (93)	Risperidone, clomipramine, erythromycin	ADHD, OCD, and Tourette's syndrome	15	Male	Deteriorated mental state

(Safer, Zito & dosReis, 2003)

21

Medicating Preschoolers



22

Similar patterns in preschoolers

Use of most classes of psychotropics among 2-4 year-olds continues to increase

- Almost half of those receiving prescriptions received two or more medications



(Coyle, 2000; Rappley, 2006; Zito et al. 2000)

23


Newer drugs top the list

Fastest increases have been in newer drugs without established efficacy or safety profiles



(Pathak et al. 2004; Rappley, 2006; Zito et al. 2000)

24




2006: more than 1,100 Florida Medicaid children under age 6 received atypical antipsychotics

(St. Petersburg Times, 2007)

25

Concerns

Treatment of preschoolers with psychiatric drugs has barely been studied




(Rappley, 2006 ; Vitiello, 2001; Waller et al. 2005; Zito et al. 2000)

26

Insufficient evidence to...

- Provide guidelines for treatment
- Establish efficacy of treatment
- Guarantee safe use
- Evaluate short- and long-term consequences on development



(Rappley, 2006 ; Vitiello, 2001; Waller, Lewellen & Bresson, 2005; Zito et al. 2000)

27

SCIENTIFIC AMERICAN

May 30, 2007

Kids on Meds -- Trouble Ahead

Antidepressants, designed for adults, may be altering the brains of kids who take them

By Paul Raeburn

28

Youths in Foster Care

More likely to be medicated



29

CBS EVENING NEWS

Are Drugs Being Misused On Foster Kids?
Three's Growing Concern: That Anti-Psychotic Drugs Are Being Missed On Children In Foster Care

msnbc

States wrestle with medicating foster kids
 Critics worry psychiatric drugs flow too freely to forgotten children

USA TODAY

Health and Behavior | Inside News

For foster kids, oversight of prescriptions is scarce

Updated 5/2/2006 12:21 AM ET

E-mail | Save | Print | RSS

National foster care

Children in child welfare settings are **2 and 3 times more likely** to be medicated than children in the general community



(Breland-Noble et al. 2004; Raghavan et al. 2005)

31

Group homes

After controlling for demographic *and* clinical factors, youths in group homes still **twice as likely** to be medicated than youths in therapeutic foster care


(Breland-Noble et al. 2004; Raghavan et al. 2005)

32

Concerns in Florida

Reports in 2001 and 2003 highlighted problems with:

- Medication without signed consent
- Medication without medical evaluations and proper follow-up monitoring
- High rates of polypharmacy




(Green, Hawkins & Hawkins, 2005; Florida Statewide Advocacy Council, 2003)

33

Florida concerns led to law

Senate Bill 1090 introduced in 2005 to restrict the state's ability to medicate foster children without the proper consent of their parents or a judge and required improved tracking of these children



34

“No List of Kids on Mood Drugs”

The Miami Herald  (September, 2006)


Child welfare officials acknowledged lacking an accurate list of children in state care receiving psychiatric drugs

- Advocates called use of these drugs in children “chemical restraints” used to control behavior

35

Part B

Public Health Concerns



36



37

Numbers of American children on psychotropics: 2006

Stimulants: 3.6 million
 Antidepressants: 2 million
 Anticonvulsants: 900,000
 Antipsychotics: 540,000

The New York Times
 (Medco Health Solutions, 2006)

38

FDA U.S. Food and Drug Administration U.S. Department of Health and Human Services

2006 FDA warning on cardiovascular effects also alerts doctors to stimulant-induced psychosis and hallucinations

The New York Times
 August 22, 2006
 F.D.A. Strengthens Warnings on Stimulants
 By GARDNER HARRIS

39

FDA U.S. Food and Drug Administration U.S. Department of Health and Human Services

2004: FDA issued a “Public Health Advisory” about all antidepressants, warning of drug-induced:

- Anxiety and panic attacks
- Agitation and insomnia
- Irritability and hostility
- Impulsivity and severe restlessness
- Mania and hypomania

40

FDA U.S. Food and Drug Administration U.S. Department of Health and Human Services

FDA “black box” warns:

“Antidepressants increase the risk of suicidal thinking and behavior (suicidality) in short-term studies in children and adolescents with Major Depressive Disorder and other psychiatric disorders”

41

CBS NEWS June 6, 2007 2:16pm FDA Orders Antidepressant Warning


msnbc Home » Health » Mental Health

Health FDA urges new warnings on antidepressants

2005: FDA extends “black box” warnings to children and adolescents

2007: FDA extends “black box” warnings to young adults 18-24



42





Antipsychotics

Skyrocketing numbers despite safety concerns

43

44


45

Antipsychotics = Fastest rise

Number of **non-institutionalized** 6-18 year-olds on antipsychotics:

1993:	50,000
2002:	532,000

(Olson et al. 2006)




St. Petersburg Times (2007)

More than 18,000 kids on Florida Medicaid prescribed atypical antipsychotics in 2006

Kids on meds: an explosion
In the last seven years, the number of children in the Medicaid fee-for-service plan who received antipsychotics has more than doubled. Prescriptions have more than tripled, and the cost to taxpayers is up more than 500 percent.

Children taking antipsychotics



47

Nationwide, antipsychotics typically prescribed to children for non-psychotic conditions

Most frequent diagnoses:

- disruptive behavior disorders, including ADHD (38%), and mood disorders (32%)


(Olson et al. 2006)

48

In Florida too...

2006: Only 8% of Florida Medicaid children receiving antipsychotics had a diagnosis of psychosis

- Half were diagnosed with attention or conduct disorders



49

Antipsychotics = polypharmacy

77% to 86% of youths taking antipsychotics do so with other drugs



(Medco, 2006; Olsson et al. 2006)

50

Safety and efficacy unknown

"We don't know the first thing about safety and efficacy of these drugs even by themselves in these young ages, let alone when they are mixed together."


Dr. Steven Hyman, former NIMH director, Harvard University provost

The Boston Globe (2006)

51

Adverse effects of "atypicals"

	Clozaril	Risperdal	Zyprexa	Seroquel	Geodon	Abilify
Clinical name	Clozapine	Risperidone	Olanzapine	Quetiapine	Ziprasidone	Aripiprazole
Major symptoms reported						
Diabetes	Severe	Mild	Severe	Moderate	Minimal	Minimal
Weight gain	Severe	Moderate	Severe	Moderate	Mild	Mild
Sedation	Severe	Mild	Moderate	Moderate*	Minimal	Minimal
Tardive dyskinesia	None	Minimal	Minimal	Minimal	Minimal	Minimal



(Correll, 2006; USA Today, 2006)

52



"Doctors need to be judicious when prescribing antipsychotic drugs to children. The use of these drugs can have the pediatric patient trading a behavioral condition for a lifelong metabolic condition that can lead to significant health complications"

—Robert Epstein, M.D., chief medical officer, Medco

53



54

The Washington Times Advertise | RSS | Site

News Elections Opinion Sports Sections Blogs Photos

Nation/Politics


Spending on psychotherapeutic drugs soars
 By Joyce Howard Price
 THE WASHINGTON TIMES
 April 1, 2007

Spending on antidepressants and other prescription drugs to treat mental disorders climbed from \$7.9 billion in 1997 to \$20 billion in 2004, an increase of more than 150 percent, a new federal report says.

55

2004: 17% of total drug spending for children was for psychotropics

- *greater than cost of antibiotics and asthma drugs*



State insurance increases likelihood of medication

Medicaid-enrolled children are more likely to:

- Receive psychotropics
- Be treated with multiple medications
- Receive medications as sole treatment


(Goodwin et al. 2001; Martin et al. 2002, 2003)

57

Use of newer antipsychotics grows faster

1996-2001: increased most dramatically in these Medicaid populations:

- Preschool children (61%)
- Ages 6-12 (93%)
- Ages 13-18 (116%)




(Cooper et al. 2004; Olsson et al. 2006; Patel et al. 2005)

58

Department of Health and Human Services
 OFFICE OF INSPECTOR GENERAL

MEDICAID'S MENTAL HEALTH DRUG EXPENDITURES

Medicaid pays more for psychotropic drugs than other Federal buyers...




INSPECTOR GENERAL
 AUGUST 2003
 OIG-04-03-0099

59

Medicaid programs struggle to contain costs

1997 - 2004: Tripling of Medicaid spending on psychotropics attributed to the expanding use of expensive atypical antipsychotics



(Duggan, 2005; Stagnitti, 2007; OIG, 2003)

60

Antipsychotics top Medicaid spending on psychiatric drugs

10 state Medicaid programs paid \$562 million on 25 psychotropic drugs
- 67% of this total spent on nine antipsychotics

(Duggan, 2005; OIG, 2003; Stagnitti, 2007)



Average prescription price for top 2 antipsychotics, 1993 vs. 2001

1993: Haldol, Mellaril = \$29

2001: Zyprexa, Risperdal = \$286

(Duggan, 2005)

62

Florida Medicaid (fee-for-service) spending on atypical antipsychotic drugs, 2002-2007

\$1.1 billion

(Farley, R., *St. Petersburg Times*, April 12, 2008)



Part D

Conclusions and Recommendations

64

Usage is increasing

Usage of all psychiatric drug classes has skyrocketed during past decade in all age groups, all ethnic/racial groups, all settings



65

Ongoing debate

Debate persists on whether disorders are under- or over-diagnosed, and under- or over-treated, with heated arguments from supporters and critics in professional and public discourse



66

Supporters argue...

- Up to 1/5 of youth have a “DSM-diagnosable disorder”
- Popularly-accepted causes of disorders are neurobiological
- Medications remove “blame”
- Stimulants greatly impact ADHD-like behavior



Critics reply...

- Medication use outpaces research evidence
- Growing use leads to increase in pediatric adverse effects
- Medicating the developing brain may lead to long-term negative changes in functioning
- No pathophysiological variable is associated with any DSM disorder



68

Fastest rise: Antipsychotics

Antipsychotics with serious adverse effects growing faster than any other drug class

- More frequently used in polypharmacy and for non-psychotic disorders, with no research evidence



Racial issues

Black children: fastest-growing group being prescribed antipsychotics

- Increase related to enormous rise in the diagnosis of bipolar disorder in this population



Soaring State Medicaid spending


Largest spending increases on antipsychotics

- Until now, states appear unable to contain such fast-rising drug costs



Young children

Children are particularly vulnerable to harm by psychiatric drugs because their brains are still developing



Research is needed to track subtle changes in children’s developing personality resulting from drug’s impact on brain

72

Children in foster care

Little empirical evidence exists to support the use of drug interventions in traumatized children

- Clinicians need to consider risk/benefit analysis of drugs vs. evidence of effective psychosocial interventions

73

Children in foster care

Experts recommend antipsychotics should not be considered first-line treatment for childhood trauma because of their serious adverse effects



74


A Critical Curriculum on Psychotropic Medications

Module 2

The End




75



A screenshot of the CriticalThinkRx website. The header features the logo and the tagline "A prescription for critical thinking about psychotropic medications". Below the header is a navigation menu with links: Home, About Us, Curriculum, Research, Links and Resources, Ask Your Peers, CE Credits, and Contact Us. A welcome message follows, stating that the project is funded by the State Attorneys General Consumer and Prescriber Grant Program. Below the text is an image of a pill bottle spilling pills.

A Critical Curriculum on Psychotropic Medications



A Critical Curriculum on Psychotropic Medications

Principal Investigator:
- David Cohen, Ph.D.

Research Coordinator:
- Inge Sengelmann, M.S.W.

Professional Consultants:

- David O. Antonuccio, Ph.D. (psychology)
- Kia J. Bentley, Ph.D. (social work)
- R. Elliott Ingersoll, Ph.D. (counseling & psychology)
- Stefan P. Kruszewski, M.D. (psychiatry)
- Robert E. Rosen, J.D., Ph.D. (law)

Flash production and design:

- Sane Development, Inc., and Cooper Design, Inc.

Voice narration and Flash editing:


- Saul McClintock



www.CriticalThinkRx.org

CriticalThinkRx was made possible by a grant from the Attorneys General Consumer and Prescriber Grant Program, funded by the multi-state settlement of consumer fraud claims regarding the marketing of the prescription drug Neurontin®


Module 3 The Drug Approval Process

Part A
The FDA and Drug Regulation

5

All drugs intended for prescription in this country must be **approved** by the U.S. Food & Drug Administration (FDA)



There are huge financial and health stakes in drug approvals



7


The FDA was established by Congress in 1906 to enforce standards on purity of medicinal compounds

Today, the FDA’s Center for Drug Evaluation and Research (CDER) oversees testing and approval of medications

8

The CDER conducts no drug tests of its own—drug firms (sponsors) pay for and conduct all tests

Based on data submitted by sponsors, CDER judges a drug’s “efficacy” and “safety”




(Avorn, 2004)

9

Some FDA mandates

- ☑ grant permission to test drugs on humans
- ☑ review data on safety and efficacy
- ☑ set criteria for drug approval
- ☑ grant or deny approval of new drugs
- ☑ require more studies, disclosure of risks
- ☑ impose fines on drug makers
- ☑ order drugs removed from market



10

1938 Federal Food, Drug and Cosmetic Act:

Basis for FDA regulation of drugs

- Passed after 100 deaths in 1937 from a toxin in a batch of sulfa drugs

(Ballentine, no date)

11

FDA’s drug testing rules tightened after **thalidomide**, prescribed to pregnant women in Europe in 1960, caused birth defects

12

As a result, 1962 amendments to *Food, Drug, & Cosmetic Act* of 1938 required sponsors to:

- ✓ demonstrate efficacy in controlled trials
- ✓ report serious adverse effects to FDA
- ✓ list all known risks (on drug label and in drug ads to doctors)

13

More recent FDA laws have been controversial

Some scientists, advocacy groups, and legislators often accuse the FDA of treating industry, not the public, as its client

(Hawthorne, 2005; Sharav, 2007)

14

Prescription Drug User Fee Act, 1992

To speed up approval times, FDA collects fees from sponsors

User fees now make up over 50% of CDER's budget

(Avorn, 2007)



Impact of user fees

Since 1992 and the birth of user fees, the FDA has slashed its own testing laboratories and network of independent drug safety experts in favor of hiring more people to approve drugs for the pharmaceutical industry

(Harris, 2004)

16

“User fees have undoubtedly constrained the FDA’s independence and influenced its decisions.”

Marcia Angell, former editor, *New England Journal of Medicine*

FDA's User-Fee Habit

washingtonpost.com
By Cindy Skrzycki
Tuesday, April 3, 2007; D01

17

Draft Guidance on Direct-to-Consumer Advertising, 1997

After 15 years of industry pressure, the FDA allowed sponsors to advertise prescription drugs directly to consumers

- DTCA is praised for providing drug information to consumers
- DTCA is criticized for increasing drug costs and promoting least effective drugs

(Gellad & Lyles, 2007; Hollon, 1999)

18

**Pediatric Research Equity Act, 2003
& Pediatric Exclusivity Act, 2004**

FDA can request studies to be conducted on children, giving sponsors an extra 6 months of exclusive marketing for every drug studied

- Acts are praised for stimulating research on drug effects and indications in children
- Acts are criticized for giving drug firms unneeded profits and using kids as guinea pigs for unnecessary drug testing

19

Part B
**FDA's
Drug Approval Process**

20

Few drugs make it to market

5,000 molecules screened in the lab = 1 obtains FDA approval as a medication

From start to finish, sponsor will spend \$100 - \$400 million to obtain FDA approval

(Goozner, 2004; Ng, 2004)

21

FDA requires that drugs intended for prescription undergo pre-clinical and clinical testing



22


**Pre-clinical testing:
2-4 years**

A promising molecule is tested in laboratory and on animals

- to establish its main biological activity and
- to rule out that it causes cancer, mutations, and birth defects



If drug remains promising after pre-clinical testing, sponsor may apply to start clinical trials on humans



Phase I trials: 1-2 years

Drug is given to 20-80 healthy volunteers to establish safe dosage levels, main adverse effects, “abuse potential”

25

Phase II trials: 2-3 years

Drug is given to 300-500 people with the illness for which the drug is supposed to be marketed

- The goal is to show promising therapeutic effects in order to justify the next phase of trials

26

Phase III trials: 2-4 years

In *randomized controlled trials* (RCTs), 1000-3000 diagnosed patients from many sites are *randomly* assigned to receive either the drug or a placebo

- Neither investigators nor patients are supposed to know who is receiving what (“double-blind”)

27

FDA approval requires only 2 positive Phase III trials, even if more trials are negative

Positive trial: on a symptom rating scale, drug-treated group shows *statistically significant* advantage over placebo-treated group

(FDA, 1998)



A drug showing “efficacy”

- ✓ has shown <5% chance of being worse than placebo
- ✓ has *not* shown that it helps patient’s condition to remit, or that it works better than another drug

(Avorn, 2004)

29

With 2 positive Phase III trials, sponsor can make a **New Drug Application (NDA)**, requesting FDA approval to market drug for a specific indication and age group covered in the trials

30

FDA reviews pre-clinical and clinical studies and decides whether the drug's benefits outweigh its risks



31

Drug label

Label summarizes information from pre-clinical and clinical trials
Exact contents are negotiated in private by FDA and sponsor
A shortened form must appear in all drug packaging and advertising, except broadcast
Label is considered the authoritative drug information

32

Phase IV trials: Post-marketing surveillance

As a condition for approval, FDA usually requests sponsor to conduct post-marketing trials
These trials evaluate the drug under ordinary conditions, with ordinary patients
Phase IV trials give more realistic view of drug's harms and benefits

33

Part C

Limitations of Clinical Trials

34

To discover new drugs for physical diseases

Researchers start with a *target* of drug action identified by understanding how a disease affects the body at the cellular/molecular levels

35

Not the same process for mental disorders...

Cellular/molecular biology of mental disorders is *unknown*—drugs tested for these problems don't target known biological anomalies
These drugs are selected based on their *effects on animal behavior* and *expected effects on people's complaints and behavior*

(Moncrieff & Cohen, 2005)

36

No biological markers exist

To repeat - mental and emotional problems *are not* equivalent to physical diseases


No cause has been shown to be exclusively biological

There is *no biological marker* for any DSM “primary mental disorder,” including schizophrenia

(Charney et al., 2002) 37

Flaws in clinical trials

Analysts and critics have revealed many problems with the design and conduct of clinical trials of psychotropic drugs



Overall conclusion:
Clinical trials do not provide definite basis to determine benefits or risks of drugs

(Cohen, 2002; Safer, 2002) 38

If this or any other citation is to a book, PsychRights will probably purchase it. Just let me know.

Trials at all phases neglect most psychoactive effects

Practice: Trials focus on measuring narrowly selected complaints and behavior

Problem: Main psychological alterations produced by drugs remain unknown

(Jacobs & Cohen, 1999; Cohen & Jacobs, 2007) 39

Phase II & III trials are very short

Practice: Most last only 3-8 weeks, and up to 70% of subjects drop out before trial’s end

Problem: Only some acute effects are detected—not those emerging over a longer time

(Cohen & Jacobs, 2007) 40

Subjects are wrongly assumed to have the “same” disorder

Practice: In a depression drug trial, a subject meeting DSM criteria for depression is eligible

Problem: 200 distinct symptom combinations = DSM diagnosis of depression

Also, subjects usually meet DSM criteria for several diagnoses

The “sameness” of subjects’ problems—needed for a valid comparison of treatments—is not established

(Beutler & Malik, 2002; Cohen & Jacobs, 2007; Emslie et al. 2002) 41

Inert pills are used as comparisons

Practice: Drugs with psychoactive effects are compared to inert sugar pills

Problem: Placebos can be active (causing physical sensations) or inert (no sensations)

Because they are more powerful, active placebos are almost never used

Also, sponsors routinely screen and exclude placebo responders from clinical trials

(Abboud, 2004; Fisher & Greenberg, 2003) 42

The “blind” is often broken

Practice: It’s assumed that patients and investigators are “blind” to treatment status

Problem: Obvious side effects in drug-treated subjects cue everyone about which treatment they’re getting. This breaks the “blind”—making objective studies impossible

(Fisher & Greenberg, 1993) 43

pretty sure already have.

High doses of comparison drugs are used

Practice: When comparing a new drug to an older drug, very high doses of the older drug are used

Problem: The older drug produces more side effects, making the newer drug appear safer

(Geddes et al., 2000) 44

Outcomes are researcher-rated rather than patient-rated

Practice: Main outcome measures are rated by *researchers*

Problem: In all Phase III pediatric trials of antidepressants, *not one of 10* parent- or child-rated scales showed advantage for the drug

(Jureidini et al., 2004) 45

Adverse effects are carelessly investigated

Practice: Most trials elicit side effects by asking subjects general questions once a week, or waiting for subjects to report them *spontaneously*

Problem: This *underestimates* rates of side effects, especially psychological and behavioral ones, giving false impression of drug’s safety

(Greenhill et al., 2003) 46

Adverse effects are mis-coded

Practice: Sponsor decides which effects qualify as “adverse drug events” and how to name them

Problem: Many adverse events are coded as something else, giving false impression of drug’s safety

(Breggin, 2002) 47

Strattera pediatric trial: Mis-coding why patients dropped out

What the researcher wrote	How the sponsor coded it	How it was re-coded after FDA reanalysis
“Parents felt ‘too many side effects’; stopped drug early; Abdominal pain, nausea, anxiety”	Protocol Violation	Adverse Event
“Increasing behavior problems, worsening oppositional behavior; depression”	Physician Decision	Adverse Event

(Lillytrials.com, 2007) 48

Post-treatment ratings unreported

Practice: Sponsor gathers data for weeks *after* subjects stop treatment, but does not submit them to FDA

Problem: How subjects rate their treatment *once they're off drugs* may contradict their ratings while *on drugs*. This discrepancy is rarely discussed or explored

(Healy & Farqhar, 1998) 49

Post-marketing trials rarely conducted

As of late 2006, more than 70% of promised Phase IV trials had not yet started...

Status	Percentage
Not yet started	71%
Terminated or completed, reports submitted	11%
Terminated before completion, no report submitted	<1%
Behind schedule	3%
In progress, on schedule	13%

(Avorn, 2007) 50

The preceding limitations of clinical trials give clinicians and policymakers false ideas about how medications can help and how they can harm people

- FDA approval by itself does not guarantee that a drug is either *safe* or *efficacious* for its intended uses

(Strom, 2006) 51

The increasing involvement of industry in clinical trials has further muddled this worrisome situation

52

Part D Blurring Science and Marketing

53

Huge payoffs can follow an FDA drug approval


Zyprexa sales since 1996: \$20 billion

These create enormous incentives to turn clinical trials into marketing tools

(Smith, 2005)

For the FDA, a clinical trial is a limited test of the efficacy of a product

For the sponsor, it's a ticket to get its product past the FDA hurdle—and possibly to blockbuster status



(Smith, 2003) 55

How sponsors turn trials into marketing tools

- ☑ design studies solely to get positive results
- ☑ suppress and twist negative results
- ☑ publish positive results multiple times

(Quick, 2001) 56

The NEW ENGLAND JOURNAL of MEDICINE

Selective Publication of Antidepressant Trials and Its Influence on Apparent Efficacy

Erick H. Turner, M.D., Annette M. Matthews, M.D., Efiha Linardatos, B.S., Robert A. Tell, L.C.S.W., and Robert Rosenthal, Ph.D.

“According to the published literature, it appeared that 94% of the trials conducted were positive. By contrast, the FDA analysis showed that 51% were positive.”

57

Contract Research Organizations (CROs)

To get drugs approved by the FDA, sponsors outsource clinical trials to CROs, a \$15 billion/year business

These private firms make it easier to:

- Enroll thousands of subjects
- Conduct more multi-site trials
- Shield trials from public scrutiny

(Hunley, 2007) 58

Conflicts in research

“It’s a house of cards built on a fundamental conflict of interest. The problem is that drug companies have inordinate influence over the evaluation of their own products. That, on the face of it, doesn’t make sense.”


- Marcia Angell, former editor, *New England Journal of Medicine*, author, *The Truth About the Drug Companies*



59

Funder’s drugs come out ahead

In 90% of studies pitting one newer antipsychotic against another, the best drug was the study sponsor’s drug




(Heres et al., 2006) 60

I have this book (if it can be found)

Independent studies don't favor newer drugs

NIMH's (CATIE) study compared 5 antipsychotics in largest schizophrenia trial. Older, cheaper drug worked as well (or as poorly)

- Regardless of drug, 3/4 of patients stopped treatment because they did not improve or had intolerable side effects

 The NEW ENGLAND JOURNAL of MEDICINE
(Lieberman et al., 2005)

61

The New York Times

November 22, 2002

Madison Ave. Has Growing Role In the Business of Drug Research

By MELODY PETERSEN

"You cannot separate advertising and marketing from the science anymore."

- Arnold S. Relman, MD, Professor Emeritus, Harvard Medical School, and former editor, *New England Journal of Medicine*

THE WEEK

The Best of the U.S. and International Media

The Corruption of Medicine

Several top medical journals recently admitted that studies they published on new medications have been tainted by undisclosed financial ties between researchers and drug companies. Does Big Pharma have too much influence over drug research?
9/22/2006

63

Part E

Problems in Drug Safety After Marketing

64

Because of the limitations of clinical trials, detecting adverse effects from drugs falls to **post-marketing surveillance**, when drugs are commonly prescribed, and used for longer periods, in more natural conditions, by more varied patients



(Strom, 2006)

This is when most adverse effects, and a more accurate portrait of the drug's risk-benefit ratio, emerge

Yet such post-marketing monitoring also appears spotty

(Lasser et al., 2002)

66

Newer drugs more likely to have hidden risks

50% of warnings occur within 7 years of a drug's introduction

Half of the withdrawals occur within 2 years

(Lasser et al., 2002)

67

Black Box Warnings

If the adverse drug reaction is serious enough to require extraordinary monitoring or special screening, the FDA will ask the drug sponsor to insert a "black box warning" in all marketing and product information to alert clinicians and consumers of the nature of the risk

68

Safety questions are "answered" post-marketing

51% of drugs get label changes
20% of drugs get new black box warnings
3-4% of drugs are withdrawn

(Strom, 2006)

69

Former and current FDA officials, outside scientists, and advocates for patients say the FDA's efforts to monitor the ill effects of drugs on the market are insufficient

70



Report: FDA so underfunded, consumers are put at risk

(December 3, 2007; http://www.usatoday.com/news/washington/2007-12-02-fda_N.htm)



FDA Is Broken, Endangers American Lives
Report Blames Congress for Cutting FDA's Budget

December 6, 2004

The New York Times

At F.D.A., Strong Drug Ties and Less Monitoring

71

Example: Prozac, 2004

Prozac was on the market for 17 years before FDA warned of increased suicidality



Sponsors of several SSRIs have been accused of not disclosing all the data from clinical trials

72

Example: Vioxx, 2004

Vioxx was taken by 20 million Americans before Merck withdrew it after links to heart attacks and strokes

Merck accused of not disclosing all the data from clinical trials



FDA Public Health Advisory: Safety of Vioxx

73

Serious Adverse Events (SAEs)

- Fatal or life-threatening, cause disability or require hospital stay

Only 1% to 10% of all drug-related SAEs are actually reported to the FDA through MedWatch



(Moore, Cohen & Furberg, 2007)

74

Thousands die annually

Reports to Medwatch of fatal drug reactions tripled between 1998-2005

- Over **80,000** deaths suspected from medications were reported by health professionals and others during that 7-year period

(Moore, Cohen & Furberg, 2007)

75

26,000 deaths suspected to be linked to 15 drugs, including:

- 3 antipsychotics and
- 1 antidepressant

Clozaril, Risperdal, Zyprexa, Paxil

(Moore, Cohen & Furberg, 2007)

76

Table 4. Most Frequent Suspect Drugs in Death and Serious Nonfatal Outcomes, 1998-2005

Drug Name	Rank/Deaths	Drug Class
Death outcome		
Oxycodone	1/5548	Opioid analgesic
Fentanyl	2/3545	Opioid analgesic
Clozapine	3/3277	Antipsychotic
Morphine	4/1616	Opioid analgesic
Acetaminophen	5/1393	Analgesic
Methadone	6/1258	Opioid analgesic
Infliximab	7/1228	DMARD
Interferon beta	8/1178	Immunomodulator
Risperidone	9/1093	Antipsychotic
Etanercept	10/1034	DMARD
Paclitaxel	11/1033	Antineoplastic
Acetaminophen-hydrocodone	12/1032	Combination analgesic
Olanzapine	13/1005	Antipsychotic
Rofecoxib	14/932	NSAID
Paroxetine	15/850	Antidepressant

(Moore, Cohen & Furberg, 2007)

77

Part F

Conclusions and Recommendations

78

FDA's independence in question

As a result of inordinately close ties to drugmakers, the FDA appears to have compromised its independence and its mandate to protect the public from dangerous products

79

Clinical trials provide skewed portrait of drug risks and benefits

Predictable limitations of trials suggest that their positive findings cannot generalize to real-life clinical conditions

Trials are especially poor at detecting adverse effects

80

Most psychological alterations produced by drugs are unstudied

Drugs' main psychological and behavioral effects can remain unknown even years after their approval by FDA and use by millions of people



Clinical trials ≠ objective evaluations of drug effects

Excessive involvement of sponsors in testing drugs may have tainted the research process, turning many clinical trials into "infomercials"



82

Conflicts of interest = suppression of negative trial findings

"Selective reporting of clinical trial results may have adverse consequences for researchers, study participants, health care professionals, and patients."

(Turner et al. 2008)

83

Need for skepticism and vigilance

Professionals should view announcements of clinical trial findings with skepticism and review them critically



84

Use new drugs cautiously

The first users of a newly marketed FDA-approved drug are the true research subjects

Public Citizen recommends waiting 7 years after marketing to use new drugs

“The public misunderstands drug safety, believing that a drug is safe at the time of marketing.”

(Strom, 2006)

85

Your role in post-marketing surveillance?

Non-medical professionals and consumers can play an important role in *observing* and *reporting* adverse drug reactions to FDA, thus helping to create a more accurate portrait of medications and their impact on people’s lives



86

A Critical Curriculum on Psychotropic Medications


Module 3

The End



www.CriticalThinkRx.org

87




A Critical Curriculum on Psychotropic Medications

A Critical Curriculum on Psychotropic Medications

- **Principal Investigator:**
 - David Cohen, Ph.D.
- **Research Coordinator:**
 - Inge Sengelmann, M.S.W.
- **Professional Consultants:**
 - David O. Antonuccio, Ph.D. (psychology)
 - Kia J. Bentley, Ph.D. (social work)
 - R. Elliott Ingersoll, Ph.D. (counseling & psychology)
 - Stefan P. Kruszewski, M.D. (psychiatry)
 - Robert E. Rosen, J.D., Ph.D. (law)
- **Flash production and design:**
 - Sane Development, Inc., and Cooper Design, Inc.
- **Voice narration and Flash editing:**
 - Saul McClintock




CriticalThinkRx was made possible by a grant from the Attorneys General Consumer and Prescriber Grant Program, funded by the multi-state settlement of consumer fraud claims regarding the marketing of the prescription drug Neurontin®



Module 4

Pharmaceutical Industry Influences on Prescribing



Part A



Expanding Drug Markets

5

Pharmaceutical drugs = Big business




World sales:
\$643 billion in 2006
\$685 billion projected for 2008

(IMS Health, 2006, 2007; Pharmaceutical Executive, 2007; Los Angeles Times, 2007) 6

Brand-name drugs

Manufacturer holds an exclusive patent to market them for about 15 years

- 40% of prescription volume
- 90% of revenues




(IMS Health, 2007; Pharmaceutical Executive, 2007)

7

Generic drugs

Once patent on marketing a brand-name drug expires, drug becomes a “generic,” and sells for much less, as other manufacturers may apply to market it



(IMS Health, 2007; Pharmaceutical Executive, 2007)

8

“Blockbuster” drugs

Generate more than \$1 billion of revenue each year

Are heavily marketed, so their manufacturer can make profits during the marketing exclusivity period


7 of the top 10 companies have 1 psychotropic drug among their top 3 blockbusters



(Pharmaceutical Marketing, 2006)

9

Antidepressants, antipsychotics, anticonvulsants: among top 6 drug classes sold in U.S.



(Pharmaceutical Executive, 2007; IMS Health, 2006)

Growing consensus:

Psychotropics are not popular because they are particularly effective

...“medicalization” and “disease mongering” also stimulate drug use

11

“Medicalization”

- Defining or treating a problem as a *medical* disease, requiring *medical* treatments


(Conrad & Leiter, 2004; Mintzes, 2002)

12

"Disease mongering"

- Turning ordinary ailments into diseases
- Framing conditions as being severe and widespread
- Seeing mild symptoms as serious
- Seeing risks as diseases

(Moynihan, Health, & Henry, 2002; Moynihan, 2002)



13

Disorders Made to Order

Pharmaceutical companies have come up with a new strategy to market their drugs: First go out and find a new mental illness, then push the pills to cure it.

Brendan I. Koerner
July/August 2002 Issue

MotherJones

Disease awareness campaigns turn healthy people into patients


Owen Dyer London



DOI: 10.1371/journal.pmed.0030189.g001


Pills are often marketed as a solution to human anxieties and dissatisfactions

14



Prescription Psychiatry

How big a dose?




Psychiatry by Prescription

Do psychotropic drugs blur the boundaries between illness and health?

PLOS MEDICINE
a peer-reviewed open-access journal published by the Public Library of Science

The Latest Mania: Selling Bipolar Disorder

David Healy



This is one of a series of articles on disease mongering in the April 2006 issue




Figure 1. Articles Elicited by Meshline Using the Mesh Term "Mood Stabilizer"

DOI: 10.1371/journal.pmed.0030189.g001

bipolar disorder. Help your doctor to help you.

treated with antidepressants and the manias with antipsychotics or lithium.

16

Part B




Marketing Expands Drug Markets

17

Cost of marketing and promoting drugs in U.S.

Industry estimates: \$29.9B



Independent estimates: \$57.5B

The NEW ENGLAND JOURNAL of MEDICINE
(Donohue, Cevalco & Rosenthal, 2007)

PLOS MEDICINE
a peer-reviewed open-access journal published by the Public Library of Science
(Gagnon & Lexchin, 2008)

18

Drug company marketing targets all players in the health care system




19

It influences physicians to prescribe through:

Gifts:

- free lunches
- drug samples
- continuing medical education
- payments for lecturing, consulting and research



20

It influences physicians to prescribe by:

- ✓ funding countless activities of professional organizations
- ✓ drug advertising in professional journals
- ✓ paying doctors to serve on “expert committees” that create and promote guidelines for drug treatments used by other doctors

21

It influences consumers to seek drugs through:

- ✓ direct-to-consumer-advertising (DTCA)
- ✓ “disease awareness” campaigns
- ✓ funding “patient advocacy” groups
- ✓ online medical information and promotions

22

It influences legislators and government agencies to approve drugs and create favorable conditions for drugmakers through:

- ✓ lobbying at all levels of government
- ✓ large donations to political parties
- ✓ payment of “user fees” to the FDA


23

It influences experts to evaluate drugs positively by:

- ✓ paying researchers to run clinical trials and develop treatment guidelines
- ✓ signing “secrecy agreements” with researchers to conceal negative drug information
- ✓ paying academics and researchers to lend their names to articles they have not written (“ghostwriting”)

24

Drug Reps



25

100,000 drug reps in the United States



~ 1 for every 6 doctors

(Oldani, 2004; Greene, 2004; Fugh-Berman & Ahari, 2007)

26

Doctors who meet frequently with reps:

- ✓ increase prescribing of newer, costlier drugs
- ✓ reduce prescribing of generics
- ✓ increase nonrational prescribing
- ✓ use rep as main information source

(Dana & Loewenstein, 2003; Reist & VandeCreek, 2004, Schwartz et al. 2001; Wazana, 2000)

27

Reps know just which doctors to target and how

Health Information Organizations combine purchased pharmacy data, AMA physician data, and patient data to determine which drugs individual physicians prefer for which diagnoses and which patient groups

This *prescription tracking* is used to tailor marketing to physicians and evaluate effects of promotions on their prescribing behavior

(Fugh-Berman, 2008)

28



Gift-giving

Very effective, even when
doctors don't think so



29

The Boston Globe

Does a drug firm's free lunch influence doctors?


By Scott Lassman | May 18, 2007

Physicians and the Pharmaceutical Industry Is a Gift Ever Just a Gift?

JAMA[®]
The Journal of the American Medical Association

30

Are doctors “on the take”?



31

The NEW ENGLAND JOURNAL of MEDICINE

A National Survey of Physician–Industry Relationships

Among a sample of 3,200 physicians:

- 83% received food at work
- 78% received drug samples
- 35% were reimbursed for CME
- 28% were paid to give lectures or recruit patients in trials

(Campbell et al., 2007)

32

The New York Times

Psychiatrists Top List in Drug Maker Gifts

By GARDINER HARRIS
Published: June 27, 2007

1997- 2005: drug companies paid Minnesota doctors \$57 million

- psychiatrists received \$6.7 million

(Ross et al., 2007; The New York Times, 2007)

33

1 in 3 Minnesota psychiatrists received money from drugmakers

“One in three Minnesota psychiatrists has received funding from drug manufacturers in the past five years, including seven past presidents of the Minnesota Psychiatric Society, two state drug policy advisers and 17 faculty psychiatrists at the University of Minnesota.”

(Olson, 2007)

34

The New York Times

May 10, 2007

Psychiatrists, Children and Drug Industry’s Role

By GARDINER HARRIS, BENEDICT CAREY and JANET ROBERTS

Psychiatrists receiving money from drug companies more likely to prescribe “off-label” antipsychotics to children

Prescription for Influence Beyond the Label

Average number of prescriptions for atypical antipsychotics for children written by Minnesota psychiatrists who received the following amounts of money from the drug makers from 2000 to 2005:

PAYMENTS	PRESCRIPTIONS*
\$5,000 or more	223
Under \$5,000	67

* For children enrolled in Minnesota’s fee-for-service Medicaid program
Sources: Minnesota Board of Pharmacy; Minnesota Medicaid

The New York Times

“Free” samples...

- ✓introduce drug into doctor’s office
- ✓generate sales, influence brand choice
- ✓Mostly go to wealthy/insured patients
- ✓63% of total promotional spending

Return-on-investment:

\$10 in sales for every \$1 spent

(Adair & Holmgren, 2005; Backer et al. 2000; Chew et al. 2000; Cutrona et al. 2008; ugh-Berman & Ahari, 2007)

36

Small gifts are powerful

Studies suggest that *the most powerful form of influence might be small gifts*


The more gifts a doctor received, the more he/she believed that they had no influence on prescribing

(Reist & VandeCreek, 2004; Dana & Loewenstein, 2003; Oldani, 2004) 37



The “gift cycle”

A three-way exchange of gifts between doctors, drug reps, and patients



(Reist & VandeCreek, 2004; Dana & Loewenstein, 2003; Oldani, 2004) 38

“Ask your doctor...”



39

1997: FDA allows full-scale, direct-to-consumer advertising (DTCA) of prescription drugs
- DTCA only allowed in the U. S. and New Zealand

(Gellad et al. 2007)

40

DTCA increases drug use by

- ✓encouraging people to visit doctor
- ✓encouraging patients to request advertised drugs
- ✓influencing doctor’s behavior through patient requests

(Gellad et al., 2007; Donohue & Bernd, 2004; Wolfe, 2002; Consumer Reports, 2007)

41

DTCA increases spending by

stimulating sales of newer, costlier drugs above older generics



(Gellad et al., 2007; Donohue & Bernd, 2004)

42

Accuracy of DTC ads questioned

1995 to 2004: FDA sent 1,359 warning letters to drug companies for false or misleading advertising

Only 4 FDA staffers review thousands of ads

(Donohue et al., 2007; Zalesky, 2006)

43

Example: 2007 Geodon ad “false and misleading”

2007 FDA letter: maker exaggerated claims of efficacy and did not mention risks of neuroleptic malignant syndrome, tardive dyskinesia, hyperglycemia and diabetes



GEODON “exaggerated claims, downplayed risks”

44

Industry funds “patient advocacy” groups

2005-2006: \$29 million to 6 groups
- 7%-91% of the groups’ budgets
Groups rarely disclose funding
Funds decline when drugmakers don’t benefit

(Philadelphia Inquirer, 2006; Los Angeles Times, 2007)

45

- o **NAMI** National Alliance on Mental Illness received \$11.7 million from 18 drugs firms in three years
- o **CHADD** Children and Adults with Attention Deficit/Hyperactivity Disorder is funded by Shire PLC, the #1 ADHD drugmaker
- o **DBSA** Depression and Bipolar Support Alliance receives more than half its funding from drug firms

(Philadelphia Inquirer, 2006; Los Angeles Times, 2007)

46

NAMI, CHADD, and **DPSA**, among “patient advocacy” groups receiving most industry funding, promote view of distress as *chronic brain disease*, requiring latest drugs and neurobiological research

47

Continuing Medical Education

“Educating” to expand markets?

48

Medical Education Communication Companies (MECCs) earned over \$1 billion in 2004 to deliver industry-sponsored continuing medical education (CME)

(Retman, 2001; Elliott, 2004; Wazana, 2000)


Industry-sponsored CME highlights sponsor’s drugs and is associated with increased prescriptions of those drugs

(Retman, 2001; Elliott, 2004; Wazana, 2000)

Concerns in U.S. Senate

APRIL 2007

Concern over drug firms’ influence on CME, and its impact on off-label drug use



(Report to Committee on Finance, US Senate, April 2007)

“Ghost” Marketing


Industry marketers and scientific journals

“Ghostwriting”

Pharmaceutical firms hire MECCs to write academic papers favorable to their products

MECCs then hire academics to publish the articles under their name without disclosure about the true source

(Moffat & Elliott, 2007)



Even without ghost-writing...

A drug firm may pay a journal \$1 million for reprints, creating enormous incentive for the journal to publish a favorable article

A former editor of *British Medical Journal* called journals “extensions of marketing arms” of drug firms and urged journals to *stop publishing all clinical trials*, and only evaluate them critically

(Moffat & Elliot, 2007; Smith, 2004; *The New York Times*, 2002)

55



Government and the politics of influence

56

Pharmaceutical Researchers and Manufacturers of America (PhRMA) represents pharmaceutical and biotechnology companies in the U.S.



57

PhRMA hired hundreds of lobbyists to help pass the Medicare Part D bill in 2004

Originally estimated to cost taxpayers \$534 billion, Medicare Part D forbids the government from negotiating drug prices



PhRMA head is Billy Tauzin, former Republican congressman from Louisiana

Drug industry lobbyists outnumber Congressmen 2:1

2006: Drug interests employed about **1,100** lobbyists, including **40** former members of Congress



Under The Influence
NEW YORK, April 1, 2007

(Center for Public Integrity, 2007; *CBS News/60 Minutes*, 2007)

59

Large investments in lobbying

2005 - 2006: \$182 million spent on federal lobbying

2005 - 2006: \$100 million spent on campaign contributions

Sales of top 20 lobbying spenders = 77% of the US drug market

(*CBS News/60 Minutes*, 2007; Center for Public Integrity, 2007)

60

Defending industry interests

Main goal in 2007:

- Oppose laws that would strengthen FDA's ability to monitor drug safety
- Fight bills that would allow Medicare to negotiate drug prices, which could reduce government drug spending by 60%

(CBS News/60 Minutes, 2007; Center for Public Integrity, 2007)

61

Part C

Conclusions and Recommendations

62

Conclusions

Industry promotion of expensive drugs permeates all phases of the life-cycle of drugs

Deceptive drug marketing is "pervasive, dangerous and primarily aimed at doctors"

63

Skepticism of industry grows

Previously "hidden" practices are increasingly exposed and scrutinized

Government hearings and legislative efforts highlight concerns over public health and public spending

64

Some doctors call for limits

Asking for stringent regulation to eliminate conflicts of interest:

- no gifts, no speaking at industry-sponsored CME, no ghostwriting, disclose research and consulting contracts, replace free samples with vouchers for patients



(Troyn et al., 2006; Washington Post, 2006)

65

Medical students take action

More Med Schools Show Pharma The Door

July 2nd, 2007 8:56 am By Ed Silverman

Last month, the American Medical Student Association ranked med schools based on their freebie policies, using a PharmFree scorecard. Since then, several schools reacted with embarrassment over their rankings.

- **Only 5 of 116 medical schools got an "A": for having a policy restricting drug industry access to students and faculty**

66

But medical schools lag behind

- The International Committee of Medical Journal Editors (ICMJE) requires full disclosure of drug companies' role in research
- But even major journals still can't ensure transparency
- A study of 108 medical schools' agreements to conduct research for drug firms found that ICMJE guidelines were rarely followed
- Researchers have little access to data or power over publishing

(Rivera & Cummings, 2002)

(Schulman et al., 2002)

67

States attempt legislation and sue drug firms

Most states have introduced bills or resolutions aimed at marketing

Several states are suing drugmakers for off-label promotion of antipsychotics and for hiding drug risks (see Module 5)

(Reist & VandeCreek, 2004; Zalesky, 2006)

68

9 in 10 Americans favor reforms

Consumer Reports survey finds strong backing for drug reforms

As Congress prepares to vote on the most significant prescription drug safety legislation in 45 years, a new *Consumer Reports* poll finds that the American public strongly backs a number of reforms. Safety issues rose to the top, with 9 of every 10 Americans supporting reforms that would require warning labels and follow-up studies on drugs with safety problems, and public disclosure of all clinical drug trials.

ConsumerReports (2007)

69

Recommended reforms to research

Create a public registry of all clinical trials

Fund clinical trials publicly, and cease drugmakers' ties to clinical research

Make raw clinical trial data accessible for independent analyses

(Antonuccio & Healy, 2008; NJPIRG Law & Policy Center, 2006)

70

Researchers' commitment?

Because research participants expose themselves to risk, information derived from them should not be misused, suppressed, or distorted

Researchers should promise to make *all raw research data available publicly*, or forego approval from Institutional Review Boards

(Antonuccio & Healy, 2008)

71

Teach prescribers, academics and consumers to:

- ✓critically evaluate drug marketing
- ✓rely on independent sources of information
- ✓implement best practices to minimize industry influence in schools, professional organizations, and mental health providers

(NJPIRG Law & Policy Center, 2006)

72


A Critical Curriculum
on Psychotropic Medications

Module 4


The End



73




A Critical Curriculum on Psychotropic Medications




A Critical Curriculum on Psychotropic Medications

- **Principal Investigator:**
 - David Cohen, Ph.D.
- **Research Coordinator:**
 - Inge Sengelmann, M.S.W.
- **Professional Consultants:**
 - David O. Antonuccio, Ph.D. (psychology)
 - Kia J. Bentley, Ph.D. (social work)
 - R. Elliott Ingersoll, Ph.D. (counseling & psychology)
 - Stefan P. Kruszewski, M.D. (psychiatry)
 - Robert E. Rosen, J.D., Ph.D. (law)
- **Flash production and design:**
 - Sane Development, Inc., and Cooper Design, Inc.
- **Voice narration and Flash editing:**
 - Saul McClintock




CriticalThinkRx was made possible by a grant from the Attorneys General Consumer and Prescriber Grant Program, funded by the multi-state settlement of consumer fraud claims regarding the marketing of the prescription drug Neurontin®

Module 5



Specific Drug Classes: Use, Efficacy, Safety



4

Part A


Overview: Psychotropic Drugs Used with Children and Adolescents



5

“Psychotropic” or “psychoactive” drugs

affect the central nervous system and alter feeling, thinking, and behaving



6

“Approved use” means...

FDA has reviewed limited data on safety and efficacy for one indication, usually in one population

A “label” for the drug is established to guide dosage and describe observed side effects



7

Fewer than 10% of psychotropic drugs are FDA-approved for any psychiatric use in children



8

Focus: Stimulants



9

Stimulants approved by FDA for pediatric use

Brand Name	Generic Name	Psychiatric Indication	Age group
Adderall, Adderall XR, Dexedrine, Dextrostat	amphetamine, dextroamphetamine	ADHD, narcolepsy	3 +
Concerta, Ritalin, Daytrana, Metadate Focalin, Focalin XR	methylphenidate, dexamethylphenidate	ADHD	6 +
Vyvanse	lisdextroamphetamine		
Strattera (not considered a stimulant)	atomoxetine		

10

Stimulants act quickly

Stimulants change behavior *within one hour* in 60-70% of children who take them

11

Long-term evidence of benefits doubtful

APA Report noted lack of data supporting long-term efficacy or safety

- Stimulants show minimal efficacy in general life domains of the child, including social and academic success



(APA Working Group on Psychoactive Medications for Children and Adolescents, 2006; MTA Cooperative Group, 2004)

12

Short-term desirable effects of stimulants at usual doses

- ✓ Increase alertness and wakefulness
- ✓ Induce sense of well-being (euphoria)
- ✓ Improve accuracy on brief physical and mental tasks



(Bezchlibnyk-Butler & Jeffries, 2005)

Effects misconstrued as therapeutic in children

- ✓ Increased repetitive, persistent behavior
- ✓ Decreased exploration and social behavior
- ✓ Increased compliance

(Breggin, 1998)

14

In Module 5 references, but couldn't find in big one

Undesirable *behavioral* effects of stimulants

- Nervousness, restlessness
- Insomnia
- Agitation
- Depression, "zombie" look
- Irritability, Aggression
- Psychological dependence
- Mania, Psychosis

(Bezchlibnyk-Butler & Jeffries, 2005)

15

Undesirable *physical* effects of stimulants

- Increased blood pressure
- Dizziness, headaches
- Palpitations
- Stomach cramps, nausea
- Appetite/weight loss
- Stunted growth
- Cardiac arrest

(Bezchlibnyk-Butler & Jeffries, 2005)

16

Stunted growth

Decreases in growth averaging $\frac{3}{4}$ " and 6 lbs. without evidence of rebound 3 years after stopping treatment



(Swanson et al., 2007)

17

Emergency room visits

2,500 children visited ERs in 2004 after taking stimulants for ADHD, most due to accidental overdoses

- 1 in 4 children had heart or blood pressure symptoms including palpitations, chest pain or fainting



(Waters, 2007)

18

2006: FDA warning on stimulants

- ✓ increased risk of sudden death in patients with heart problems
- ✓ increased aggression, mania and/or psychotic symptoms (including hallucinations)

The New York Times

August 22, 2006
F.D.A. Strengthens Warnings on Stimulants

19


Definite risk of tolerance and dependence

Stimulants prescribed to children are Drug Enforcement Administration (DEA) "Schedule II Drugs," indicating a high risk of tolerance and dependence

RITALIN LA[®] is a federally controlled substance (CII) because it can be abused or lead to dependence. Keep RITALIN LA[®] in a safe place to prevent misuse and abuse. Selling or giving away RITALIN LA[®] may harm others, and is against the law.

20

Focus: Antidepressants



21

FDA-approved antidepressants for pediatric use

Brand Name	Generic Name	Psychiatric Indication	Age group
Sinequan	doxepin	OCD	12+
Anafranil	clomipramine		10+
Luvox	fluvoxamine		8+
Zoloft	sertraline		6+
Tofranil	imipramine		
Prozac	fluoxetine	Depression, OCD	7+

22

CDC: Antidepressants most prescribed drugs in U.S.



Antidepressants such as Paxil, Prozac and Lexapro are among America's most-prescribed drugs.

CNN.com/health 2007

23

But are they effective?

Meta-analyses of drug vs. placebo studies show 75-82% of the response was duplicated by placebo

- 57% of studies submitted to FDA failed to show a difference between drug and placebo

(Moncrieff et al., 2004; Kirsch et al., 2002; Kirsch & Sapirstein, 1998)

24

Unimpressive evidence from FDA's complete adult database

"[I]n 189 trials of 53,048 adult subjects with psychiatric disorders ... Approximately 50% of subjects who received active drug and 40% of subjects who received placebo were designated as responders."

(Stone & Jones, 2006)

The entire scientific case for antidepressants rests on this 10% difference—which may result from biases in the conduct of clinical trials

25

FDA analysis of pediatric trials concurs

Only 3 of 15 published and unpublished randomized controlled trials show SSRIs as more effective than placebo in depressed children

None of the studies found drugs better on client- or parent- rated measures

(Laughren, 2004)

26


No evidence that older antidepressants (tricyclics or MAO inhibitors) have any efficacy with depressed youths

(Somers-Flanagan & Somers-Flanagan, 1996)

27

Short-term desirable effects at usual doses

- ✓ Increased physical activity
- ✓ Elevated mood
- ✓ Decreased expressions of distress such as crying, hopelessness
- ✓ Improved sleep and appetite



(Bezchlibnyk-Butler & Jeffries, 2005)

28

Undesirable *behavioral* effects of antidepressants

- Anxiety, nervousness
- Agitation, irritability
- Mood swings, mania
- Aggressiveness
- Thoughts of suicide
- Attempted or actual suicide

(Antonuccio et al., 1999; Preda et al., 2001; Healy, 2003)

29

Undesirable *physical* effects of antidepressants

- Gastrointestinal distress (nausea, vomiting, stomach pain, constipation, diarrhea)
- Sexual problems (loss of libido, anorgasmia, erectile dysfunction)
- Sleep disruption (insomnia, hypersomnia)
 - Urinary retention
 - Blurred vision
 - Weight gain
- Headaches, dizziness

(Antonuccio et al., 1999; Preda et al., 2001; Healy, 2003)

30

Six clusters of withdrawal effects likely upon abrupt discontinuation of SSRI antidepressants

1. Neurosensory (vertigo, tingling & burning)
2. Neuromotor (tremor, spasms, visual changes)
3. Gastrointestinal (nausea, vomiting, diarrhea, weight loss)
4. Neuropsychiatric (anxiety, depression, crying spells, irritability, suicidal thinking)
5. Vasomotor (heavy sweating, flushing)
6. Other (insomnia, vivid dreaming, fatigue)

(Schatzberg et al., 2006) 31

Antidepressants double risk of suicidality

FDA U.S. Food and Drug Administration U.S. Department of Health and Human Services

2005: FDA issues “black box” warning of “Suicidality in Children and Adolescents”:
“Antidepressants increase the risk of suicidal thinking and behavior (suicidality)”

(22 RCTs testing 9 antidepressants: 2.3% rate of serious suicidal events among drug-treated children, vs. 1.2% among placebo treated—no completed suicides)

32

“Activation” syndrome: A more common risk

FDA also warns of *increased agitation, irritability, aggression, worsening anxiety, severe restlessness, and other unusual behaviors* in youth treated with antidepressants

(Breggin, 2006) 33

Concern over “prescription cascade”

Continued exposure to the drug can lead to effects misinterpreted as psychiatric symptoms (such as mania), leading to increases in dosage or additional drugs—when reducing or stopping the drug would relieve the patient’s discomfort

(Breggin, 2006) 34

Focus: Anticonvulsant Drugs



35

Anticonvulsants on U.S. market (antiepileptics, antiseizure drugs)

Brand Name	Generic Name	Yr of intro
Tegretol, Equetro	carbamazepine	1968, 2004
Neurontin	gabapentin	1993
Lamictal	lamotrigine	1994
Depakene, Depakote	valproate	1995
Topamax	topiramate	1997
Trileptal	oxcarbazepine	2000

36

Anticonvulsants FDA-approved for pediatric seizure disorders



Brand Name	Generic Name	Approved Indications	Age
Tegretol, Equetro	carbamazepine	NO PSYCHIATRIC INDICATIONS	Any
Gabitril	gabapentine		12 +
Depakote, Depakene	divalproex sodium, valproate		10 +
Topamax	topiramate		3 +
Neurontin	gabapentin		2 +
Lamictal	lamotrigine		
Trileptal	oxcarbazepine		

Anticonvulsants widely promoted as “mood stabilizers”

Use started in 1980s-1990s due to dissatisfaction with lithium and antipsychotics in treatment of Bipolar Disorder

Use spread rapidly with the promotion of “mood stabilizer” expression and of Bipolar Disorder diagnosis in children

(Healy, 2006)



Polypharmacy without psychotherapy

More than 90% of children diagnosed with Bipolar Disorder received more than 1 psychoactive drug

Less than 40% received psychotherapy

(Moreno et al., 2007)

Scant empirical support

No studies confirm the efficacy and safety of anticonvulsants to treat Bipolar Disorder in children and adolescents

“Despite the frequent use of antiepileptic drugs in the treatment of juvenile bipolar disorder, migraine, and neuropathic pain, the data are insufficient to make recommendations regarding the efficacy of antiepileptics in these conditions in children and adolescents.” (Golden et al., 2006)

(Kowatch et al., 2000, 2005; National Institute of Mental Health, 2000; Ryan, Bhatara & Peret, 1999)

Most trials are open, small, and show limited response in youth

Half of all participants in an open trial of lithium, divalproex, or carbamazepine did not respond to treatment

- 58% received *at least one* mood stabilizer *plus* a stimulant, an atypical antipsychotic, or an antidepressant

(Lopez-Larson & Frazier, 2006)

43

Desired behavioral effects of anticonvulsants

- ✓ Reduce aggression and impulsivity
- ✓ Calm restlessness and excitability

(Bezchlibnyk-Butler & Jeffries, 2005)

44



2008: FDA warns anticonvulsants double risk of suicidal behavior or ideation

Risk is highest in treatment of epilepsy—which rules out psychiatric status as confounding variable

45

Undesired *behavioral* effects of anticonvulsants

- Depression, sedation
- Hostility and irritability
- Anxiety, nervousness
 - Hyperactivity
 - Abnormal thinking
- Confusion and amnesia
 - Slurred speech
- Sedation, sleepiness

(Bezchlibnyk-Butler & Jeffries, 2005)

46

Undesired *physical* effects of anticonvulsants

- Nausea and dizziness
- Vomiting and abdominal pain
- Headaches and tremors
 - Fatal skin rashes
 - Hypothyroid
 - Blood disorders
- Pancreatitis, liver disease
- Birth defects and menstrual irregularities
 - Withdrawal seizures


(Bezchlibnyk-Butler & Jeffries, 2005; Gonzalez-Heydrich et al., 2003)

47

Birth defects of concern given new patient profiles

Anticonvulsants cross placenta and increase the risk of fetal malformations and cognitive impairments in children exposed in utero

- *Highest rates for valproate and carbamazepine*



(Adab et al., 2006)

FDA black-box warnings

Depakote	Liver toxicity (particularly for under 2 yrs of age); birth defects; pancreatitis
Tegretol	Aplastic anemia and agranulocytosis (severe reduction in white blood cells)
Lamictal	Serious rash requiring hospitalization; Stevens-Johnson Syndrome for under 16 yrs of age (fatal sores on mucuous membranes of mouth, nose, eyes and genitals)
All anticonvulsants	Suicidal ideation and behavior

49



“Atypical” (newer, 2nd generation) antipsychotics on U.S. market

Brand Name	Generic Name	Yr of intro
Clozaril	clozapine	1989
Risperdal	risperidone	1994
Zyprexa	olanzapine	1996
Seroquel	quetiapine	1997
Geodon	ziprasidone	2001
Abilify	aripiprazole	2002
Invega	paliperidone	2007

51

FDA-approved psychiatric indications of atypicals

Brand Name	Indication	Age
Risperdal	Autism, bipolar mania, schizophrenia	5 +
Abilify	Schizophrenia	10+
Clozaril	Treatment resistant schizophrenia	Adults only
Zyprexa	Bipolar mania, schizophrenia	
Seroquel		
Geodon		
Symbyax		
Invega		

52

FDA-approved psychiatric indications of typicals for children

Brand Name	Generic Name	Psychiatric Indication	Age
Orap	pimozide	Tourette's Disorder (for Haldol non-responders)	12 +
Haldol	haloperidol	Schizophrenia, Tourette's Disorder	3 +
Mellaril	thioridazine	Schizophrenia	2 +

Typicals make up less than 5% of FL Medicaid prescriptions of antipsychotics

53

“Typical” & “Atypical” antipsychotics


Since 1950s, antipsychotics were used to treat psychoses, despite high toxicity and limited effectiveness

Newer, expensive “atypical” antipsychotics were heavily promoted in the 1990s as safer and more effective

54

Put this in section about no stats from State

Yet, newer no better than older...




2005: largest-ever schizophrenia treatment study finds atypicals neither more effective nor better tolerated than older drug

- 75% of patients quit either drugs within 18 months due to inefficacy or intolerable side effects

(Lieberman et al., 2005) 55

Non-psychotic diagnoses in children treated with atypicals

Diagnosis	% of Florida Medicaid children on antipsychotics (2006)
ADHD / Conduct Disorder	48
Nonpsychiatric, Anxiety, Other Psychiatric	27
Bipolar / Depression	13
Schizophrenia / Psychosis	8
Austism / Mental Retardation	4

 (2007) 56

Assuming this is the CATIE study, I have it.

“Aggression” said to account for most of the antipsychotic prescribing in children and adolescents

(Patel et al., 2005) 57

But do antipsychotics effectively control aggression?

The latest randomized-controlled trial found *placebo more effective* than either a typical (haloperidol) or atypical (risperidone) antipsychotic to reduce aggression in patients with intellectual disability

Trial had no drug company sponsorship

(Tyrer et al., 2008) 58

“Antipsychotic drugs should no longer be regarded as acceptable routine treatment for aggressive behavior in people with intellectual disability.”

(Tyrer et al., 2008) 59

Few pediatric clinical trials of atypicals for any indication

As of 2006, only a few studies of direct AAP comparisons with placebo

Most studies are short-term (3-6 weeks) and results favor the funder’s drugs

(McDonagh et al., 2006) 60

“There are no studies that have shown (atypicals) are safe, or for that matter, that they are effective for children...The bottom line is that the use of psychiatric medications far exceeds the evidence of safety and effectiveness.”

Ronald Brown, Chair,

2006 American Psychological Association
Task Force on Psychotropic Drug Use in Children

**St. Petersburg
Times** (2007)

61

Dopamine-blocking action of all antipsychotics explains

- ✓ indifference, sedation, drowsiness, apathy
 - ✓ reduced spontaneity and affect
- ✓ reduced ability to monitor one's state
 - ✓ increased abnormal movements
- ✓ cognitive and motor impairments
- ✓ confusion and memory problems
- ✓ depression, mood swings, agitation

(Bezchlibnyk-Butler & Jeffries, 2005)

62

Desirable effects of antipsychotics at usual doses

- ✓ suppress psychotic symptoms (delusions, hallucinations, agitation)
- ✓ suppress manic symptoms (euphoria, expansiveness, irritability)

(Bezchlibnyk-Butler & Jeffries, 2005)

63

Effects misconstrued as therapeutic

- ✓ increased indifference
- ✓ reduced spontaneity and affect
- ✓ reduced ability to monitor one's state
- ✓ increased compliance with social norms

(Bezchlibnyk-Butler & Jeffries, 2005)

64

Undesirable **behavioral** effects of antipsychotics

- Cognitive and motor impairments
- Sedation, drowsiness
- Confusion and memory problems
- Anxiety
- Depression, mood swings
- Abnormal thinking
- Hostility, aggression

(Bezchlibnyk-Butler & Jeffries, 2005)

65

Undesirable **physical** effects of antipsychotics

- Weight gain, high blood sugar
- Abnormal movements (all body parts)
- Diabetes
- Cardiac problems
- Liver problems, jaundice
- Neuroleptic malignant syndrome
- Death

(Bezchlibnyk-Butler & Jeffries, 2005; Lindenmayer et al., 2003; Meyer, 2001)

Hormonal dysfunctions

Elevated prolactin levels cause:

- ✓sexual and menstrual disturbances
- ✓infertility
- ✓decreased bone density

(Bezchlibnyk-Butler & Jeffries, 2005; Correll & Carlson, 2006; Patel et al., 2005)

Extrapyramidal symptoms (abnormal movements)

Akathisia: inner distress, rocking, pacing, agitation

Dystonia: sudden, bizarre muscle spasms

Dyskinesia: rhythmic movements of face, mouth and tongue, sometimes of hands and feet

Parkinsonism: rigid muscles, loss of facial expression, unsteady gait, drooling

(Campbell, Rapaport & Simpson, 1999)

68

Tardive dyskinesia risk highest for typical antipsychotics

Long-lasting abnormal movements affect 12% to 35% of children who receive typical antipsychotics for more than 3 months

(Campbell, Rapaport & Simpson, 1999)

69

Weight gain and diabetes

50% of patients on antipsychotics gain 20% of their weight (primarily as fat)

Weight gain linked to "metabolic syndrome"

3 Schizophrenia Drugs May Raise Diabetes Risk, Study Says

By ERICA GOODE
Published: August 25, 2003

The New York Times

(Bezchlibnyk-Butler & Jeffries, 2005; Correll & Carlson, 2006; Patel et al., 2005)

Neuroleptic malignant syndrome

Can occur with any antipsychotic agent, at any dose, at any time

Symptoms: extreme muscular rigidity, high fever, & altered consciousness

1-2% rate per year

Fatal if untreated

(Bezchlibnyk-Butler & Jeffries, 2005; Silva et al., 1999)

71

3 atypicals suspected in nearly 4,500 deaths reported to FDA, 1998-2005

Clozaril: 3,277 deaths

Risperdal: 1,093 deaths

Zyprexa: 1,005 deaths

(Moore, Cohen & Furberg, 2007)

72

FDA “black-box” warnings

All atypicals	Increased mortality in frail elderly
Clozaril	Serious risk of agranulocytosis (severe drop in white blood cells), seizures, myocarditis, and other cardiovascular and respiratory effects
Seroquel	Risk of suicidality in children and adolescents

73

“For many adults, and a small number of children, these agents can be an important component of treatment. However, it’s so rare to find an example where evidence-based alternatives were exhausted prior to starting an atypical antipsychotic in a child that I have not found one yet in three years of searching.”

Mark E. Helm, MD, MBA
 Medical Director, Evidence-Based Prescription Drug Program
 University of Arkansas Medical Sciences
 College of Pharmacy, 2007

74

Part B

Lawsuits against drug makers shed light on illegal promotion and serious risks

The New York Times
 December 18, 2006

Drug Files Show Maker Promoted Unapproved Use

By ALEX BERENSON

75


States sue drug makers for illegal marketing of unapproved uses

to recover money states paid to purchase atypical antipsychotics and the costs of medical care for the people injured by these drugs

(Pringle, 2007; Kesselheim & Avorn, 2007)



Patients sue, charging that drug makers did not adequately warn about severe weight gain, pancreatitis, diabetes, and other risks



(Pringle, 2007; Kesselheim & Avorn, 2007)

The New York Times

January 5, 2007

Lilly Settles With 18,000 Over Zyprexa

By ALEX BERENSON

78

Zyprexa lawsuits

2007: Several states sue Eli Lilly for downplaying or hiding data linking use of the drug to weight gain and hyperglycemia

- *Most of those states' Medicaid spending on antipsychotics is for Zyprexa*

79

2007: Zyprexa settlements top \$1.2 billion, so far

Eli Lilly has paid more than **\$1.2 billion** to settle 30,000+ Zyprexa lawsuits

- The settlements required data on rates of adverse effects be kept *secret*

(Berenson, 2008)

80

2008: Feds, Eli Lilly negotiate \$1 billion Zyprexa fine

If a deal is reached, it would be the largest fine ever paid by a drug company for breaking the federal laws governing how drugmakers can promote their medicines

The New York Times
Thursday, February 7, 2008

Lilly Considers \$1 Billion Fine To Settle Case

81



Department of Justice

FOR IMMEDIATE RELEASE
FRIDAY, SEPTEMBER 28, 2007
WWW.USDOJ.GOV

2007: Bristol-Myers Squibb pays \$515 million over illegal marketing and pricing of Abilify, Serzone, other drugs

82

Litigation has

- exposed shady practices of pharmaceutical manufacturers
- uncovered previously hidden data about adverse events
- helped doctors reassess risks and benefits of some drugs and think critically about the available "evidence"

(Kesselheim & Avorn, 2007)

83

Part C Conclusions and Recommendations

84

Evidence “poor” for the use of psychotropics in children

- Little or no evidence of efficacy and safety of long-term use of these drugs in children
- Clear evidence of harm and risk of serious adverse events, including death
- Risk-benefit ratio especially poor for antidepressants, anticonvulsants, and antipsychotics

85

Need to rethink risk-benefit ratio

Risks for adverse events, including death, increase with the number of concomitant drugs administered
 Risks for adverse events are higher in children, who are receiving adjusted adult dosages of drugs rarely studied in children

(Brown & Sammons, 2002; Riddle, Kastelic & Frosch, 2001; Vitiello, 2001) 86

Side effects leading to multiple medications?

After initial medication, side effects may be viewed as mental disorders and drugged, in a “prescribing cascade” of polypharmacy that keeps children at risk with no sign of behavioral improvement

87

Available evidence does not justify use of psychotropic drugs as first-line treatments for children and adolescents



Reassess all cases?

Given known risks and dearth of valid studies showing benefits, cases of children receiving psychiatric medications should be reassessed

Children are involuntary patients. To support continuing psychotropic drug treatment, *rock-solid* rationale should be provided in every single case

89

A Critical Curriculum on Psychotropic Medications


Module 5


The End



www.CriticalThinkRx.org

90





A prescription for critical thinking about psychotropic medications

[Home](#) [About Us](#) [Curriculum](#) [Research](#) [Links and Resources](#) [Ask Your Peers](#) [CE Credits](#) [Contact Us](#)


Welcome to **CriticalThinkRx**, a project funded by the State Attorneys General Consumer and Prescriber Grant Program. The project is one of 28 in 19 states funded by a multi-state settlement of consumer fraud claims against a pharmaceutical company regarding the marketing of a psychotropic drug.

A Critical Curriculum on Psychotropic Medications




A Critical Curriculum on Psychotropic Medications

- **Principal Investigator:**
 - David Cohen, Ph.D.
- **Research Coordinator:**
 - Inge Sengelmann, M.S.W.
- **Professional Consultants:**
 - David O. Antonuccio, Ph.D. (psychology)
 - Kia J. Bentley, Ph.D. (social work)
 - R. Elliott Ingersoll, Ph.D. (counseling & psychology)
 - Stefan P. Kruszewski, M.D. (psychiatry)
 - Robert E. Rosen, J.D., Ph.D. (law)
- **Flash production and design:**
 - Sane Development, Inc., and Cooper Design, Inc.
- **Voice narration and Flash editing:**
 - Saul McClintock



www.CriticalThinkRx.org

CriticalThinkRx was made possible by a grant from the Attorneys General Consumer and Prescriber Grant Program, funded by the multi-state settlement of consumer fraud claims regarding the marketing of the prescription drug Neurontin®



Module 6

Non-Medical Professionals and Psychotropic Medications: Legal, Ethical and Training Issues




Part A

Legal limits to prescribing psychotropic medications

5

Who can prescribe?

Most states grant full or partial prescriptive authority to licensed physicians, dentists, advanced nurse practitioners, pharmacists, podiatrists, and optometrists



(NASW, 2005; Norfleet, 2002; Wiggins & Wedding, 2004)

6

How about social workers?

Survey of a national sample of 176 practitioners in late 1990s

- 52% opposed to obtaining prescription privileges
- 19% in favor
- the rest said “maybe” or did not respond

(Piotrowski & Doelker, 2001)



Professional associations' stances

American Psychological Association supports psychologists' efforts to gain prescriptive authority

National Association of Social Workers views prescription as beyond the scope of the profession

American Psychiatric Association actively opposes all such initiatives from non MDs



Part B

**Ethical and Legal Issues:
Competence and Training
Informed Consent
Confidentiality**



15

Professional *competence* is a core principle in the codes of ethics and standards for practice of various helping professions

(ACA, AMHCA, APA, NASW)

16

To maintain competence, professional codes recommend

Education and training
Consultation
Supervision
Continuing education

(ACA, AMHCA, APA, NASW)

17

Competence requires

- ✓ knowledge of valid information relevant to practice
- ✓ regular critical review of literature and emerging information
- ✓ participation in relevant and unbiased CE

(ACA, AMHCA, APA, NASW)

18

No specific standards address working with clients and others around medication-related issues

In the absence of standards, Codes advise exercising careful judgment and taking responsible steps to ensure competence and protect clients from harm

(ACA, AMHCA, APA, NASW)

19

Knowledge = Competence Training = Knowledge

Without knowledge about drugs, counselors, psychologists and social workers are ill-prepared to meet their clients' needs

Psychopharmacology should be part of training for non-medical practitioners

(Barnett & Neel, 2000; Bauer, Ingersoll & Burns, 2004; Bentley, 2005; Carlson, Thaler & Hirsch, 2005; Dziegielewski & Leon, 1998; Farmer, Walsh & Dziegielewski, 1998; Ingersoll, 2000)

20

Knowledge increases confidence and empowers non-medical professionals to participate fully in multidisciplinary environments

(Farmer, Walsh & Bentley, 2006; Dziegielewski, 1998; Littrell, 2003)

Education vs. indoctrination

Students & practitioners must be *educated* rather than *indoctrinated*, and should be exposed to controversies, uncertainties in knowledge, and well-argued alternatives to popular views

(Dziegielewski, 1998; Gomory & Lacasse, 2001; Littrell, 2003)

22

Special guidelines needed

Use of polypharmacy
Integrating psychosocial and biological therapies
Specific groups, such as children, older persons, pregnant women
Ethical and critical thinking skills in the age of "Big Pharma"

(Buelow & Chafetz, 1996; Chafetz & Buelow, 1994; Dunivin & Southwell, 2000; Freimuth, 1996; Levant & Shapiro, 2002; Smyer et al., 1993)

23



24

Why obtain informed consent?

Informed consent is the bedrock of professional practice in a free society

- It promotes the **right to self determination**, prevents harm and provides for the **client's best interest**

(Cohen & Jacobs, 2000; Strom-Gottfried, 1998; Littrell & Ashford, 1995; Littrell, 2003)

What is informed consent?


A systematic *process* intended to guarantee the client's right to choose, to privacy and to safety



(Dell et al 2008; Littrell & Ashford, 1995; Littrell, 2003; Strom-Gottfried, 1998)

What is *not* informed consent?


Having a client sign-off on services without a clear understanding of the information, including uncertainties about the treatment



(Cohen & Jacobs, 2000; Littrell & Ashford, 1995; Reamer, 2003) 27

Validity of consent forms

Blanket consent forms lack specificity and have been challenged in court
Signing a blank consent form to be completed later *is not* valid consent



(Littrell & Ashford, 1995; Reamer, 2003; Strom-Gottfried, 1998) 28

Standards for valid consent

1. Avoid coercion and undue influence
2. Assess client competence to consent
3. Specify procedures or actions in the form
4. Inform clients of the right to refuse or withdraw consent
5. Provide adequate information on risks, benefits **and** alternatives to treatment

(Reamer, 2003)

29

Coercion or undue influence

Practitioners who *want* clients to agree to treatments or procedures may be exercising undue influence and will jeopardize validity of their consent

(Dell et al 2008; Littrell & Ashford, 1995; Reamer, 2003; Strom-Gottfried, 1998)

“Adequate” information

- ✓ Critical findings on usefulness, ineffectiveness and reported information on harm
- ✓ Description of the hoped-for benefits and how success will be evaluated
- ✓ Alternatives to treatment being proposed
- ✓ Costs of treatment

(Littrell, 2003; Littrell & Ashford, 1995; Strom-Gottfried, 1998) 31

Knowledge of alternatives

Lack of knowledge about the alternatives to proposed treatment invalidates informed consent

Competence by providers in a variety of treatment methods is essential to informed consent

(Littrell, 2003; Littrell & Ashford, 1995; Strom-Gottfried, 1998) 32

Encourage questions

Informed consent should serve to empower clients to make intelligent decisions about their care, not protect practitioners from liability

Practitioners must ensure the persons receiving the information understand it, and should encourage questions

(Littrell, 2003; Cohen & Jacobs, 2000; Strom-Gottfried, 1998; Tan et al., 2007)

Competence to consent

“The capacity to act on one’s own behalf, to understand and weigh potential outcomes, to anticipate future consequences of a decision.”

(Tan et al., 2007)

34

Assessing competence to consent

In youths, assessment considers intelligence and cognitive functioning, maturity, impact of any distress, seriousness and urgency of situation, and impact of youth’s relationships

Refusing to consent does not mean incompetence

(Dell et al 2008; Tan et al 2007)

Cognitive capacity of children

By about age 9, children reach the same conclusions as adults, but by different strategies

By about age 14, minors show the same risk-benefit reasoning as adults and can participate in the consent process

(Dell et al 2008; Spetie & Arnold, 2007)



Respect for autonomy

Older children and adolescents should participate in the consent process in order to protect them from being subjected to treatment procedures against their will, and to respect their developing autonomy and personhood



(Dell et al 2008; Spetie & Arnold, 2007)

37

Third-party representation



Those who cannot give consent require a third party to act “in their best interests”

There are many views on just what this means...

(Spetie & Arnold, 2007)

38

What about preschoolers?

Are parents fully able to carry out their advocacy role?

Their capacity to act in their young child’s best interest warrants careful evaluation



(Dell et al 2008; Spetie & Arnold, 2007)

39

“The clinician must be watchful for caregivers who may have ulterior motives and want a child to be medicated for their own convenience, or because pharmacotherapy may simply be ‘easier’ than behavioral therapy, or as is more often the case, caregivers who have unrealistic expectations about what benefits a treatment may potentially hold for the child.”

(Dell et al., 2008, p. 105)

Constitutional right to refuse or withdraw consent

Clients have the right to refuse or withdraw consent at any time and must be informed of this right

State and federal courts have consistently ruled that it is unfair to allow forced medication without “adequate” procedural guidelines

(Bentley, 1993)

41

Forced treatment remains a most controversial issue

Although a fixture of mental health interventions, involuntary treatment must be *literally* “option of last resort”

Opponents of forced treatment assert that it violates one’s fundamental human rights, creates distrust of helpers, and undermines the foundation for recovery

(Bassman, 2005)

42

Taking psychotropic medications, having a psychiatric diagnosis, or experiencing major distress, does not by itself provide grounds for being denied the right to refuse or withdraw consent



(Bentley, 1993)

43

Confidentiality



44

Confidentiality vs. privacy

U.S. Constitution guarantees privacy rights, not confidentiality, to the individual

Confidentiality is essential to develop trust between client and professional



(Corcoran, Gorin & Moniz, 2005; Hanson & Sheridan, 1997; Millstein, 2000)⁴⁵

“Duty to protect”

However, the state can breach confidentiality if it has a rationale for seeing the information, such as the “duty to protect” client or others from harm



(Corcoran, Gorin & Moniz, 2005; Millstein, 2000)

46

Relinquishing confidentiality

Managed care organizations and publicly-funded payers require information from providers about clients’

- psychiatric diagnoses
- treatment procedures
- progress and outcomes



(Bilynsky & Vernaglia, 1998; Corcoran, Gorin & Moniz, 2005; Millstein, 2000)

Ethical mandates

Clients must be informed of, *and authorize*, all disclosures made to insurers and advised of the potential risks of such disclosure *before* disclosure is made

(Reamer, 2001; Millstein, 2000)

48

Part C

Emerging Legislative Issues


Concerns over medicating children lead to new laws



49

States respond to concerns

In 2004, several states passed laws limiting recommendations from school personnel about medications, and requiring their training where administration of drugs was allowed



2005: U.S. House of Representatives passes *Child Medication Safety Act* (H.R. 1790)

- Bill seeks to protect children from being forced to take psychotropic drugs as a pre-condition for attending public school, and intends to restore parental authority over decisions about their children's health

51

Florida limits school's roles F.S. 1006.0625

Public schools cannot require students to receive psychotropic medication as a condition for attending school

"Any medical decision made to address a student's need is a matter between the student, the student's parent, and a competent health care professional chosen by the parent."

52

F.S. 39.407 places limits on medicating children in state custody

Children under state care can be medicated only after obtaining *express and informed consent* from the parent, or, if parental rights have been terminated, receiving authorization from a judge

Florida and other states now require state agencies to keep list of foster care children on meds—but no register in U.S. tracks health effects of prescriptions on kids

53

Mental health screening debate

Joining the list of issues hotly debated is a 2003 Presidential task force recommendation to screen all school-aged children for mental health problems



(President's New Freedom Commission Report on Mental Health, 2003)

Early detection or pharmaceutical ploy?

Pros: early detection and treatment of disorders



Cons: invalid diagnoses and screening instruments; drug companies attempt to increase market share for psychiatric drugs

55

Part D

Psychotropic Medications and Children: “First Do No Harm”



56

“Children and adolescents are deemed vulnerable populations, at risk of being harmed by unethical or suboptimal practice and research and are in need of protection.”



(Spetie & Arnold, 2007, p. 15)

57

Medications have socio-cultural implications and impact children's identities



(Dell et al, 2008; Floersch, 2003)

58

How do children interpret their taking drugs?

To make sense of everyday medication treatment, children develop “illness narratives”

They may learn to see themselves as “defective” and unable to control their actions



(Dell et al 2008; Floersch, 2003)

59

Medication “messages”

“Better living through chemistry”:

Children learn to use drugs to deal with behavioral, emotional, academic and social difficulties



(Dell et al 2008; Floersch, 2003; Jacobs, 2006)

60

Competent practice involves listening and responding to how youths make sense of their medication experience

This requires therapeutic and personal interpretation

(Dell et al 2008; Floersch, 2003; Rappaport & Chubinsky, 2000) 61

In child and adolescent psychiatry, medication decisions are infrequently guided by scientific knowledge, as data on safety and efficacy for most psychotropics in youths remains limited

(Jensen et al., 1999; Matsui et al. 2003; Spetie & Arnold, 2007; Vitiello, 2003)



“The bottom line is that the use of psychiatric medications far exceeds the evidence of safety and effectiveness”

Ronald Brown, Chair,
2006 American Psychological Association (APA)
Working Group on Psychoactive Medications for
Children and Adolescents

St. Petersburg
Times
(2007)

63

“Whether one subscribes to the Hippocratic dictum ‘first, do no harm’ or takes a risk-benefit approach to treatment, it is impossible to discount possible unwanted treatment effects.”



(APA Working Group on Psychoactive Medications for Children and Adolescents, 2006, p. 27)

Part E

Conclusions and Recommendations

65

Non-medical professionals may neither prescribe, dispense, or administer drugs, but they may discuss any medication-related issue with their clients, including how their clients can attain their goals with the use or non-use of medications

66

Legal implications

Even professionals who do not prescribe are being called to testify in court about matters that directly concern treatment of clients with psychotropic medications

67

Training for competence

To remain competent in this emerging field requires basic education and training, including critical perspectives on drug use and marketing

68



Professionals working with children receiving psychotropic drugs must take responsibility for their education, and be accountable to clients and society for *their own decisions* about medication-related issues

69

Ethical standards

A practitioner's involvement in referring children for medication, encouraging medication compliance, and monitoring effects, must rest on the *highest* ethical standards

70



Can anyone ethically reassure clients about the safety of psychiatric drugs for children when information is not yet available?

(Littrell, 2003)

71

Balancing risks and benefits

When considering treatments, practitioners have an *ethical responsibility* to balance potential benefits with potential risks *and* to discuss both with parents as well as older children to *obtain informed consent from both*

72

“The potential for benefit from these medications must be balanced against the risks of not only the physical side effects, but also the social stigma, cost, inconvenience, and even family disapproval that can accompany even the most seemingly clear-cut, evidence-based treatment recommendation.”

(Dell et al., 2008, p. 99)

Given all the known risks associated with psychotropic drugs, attempting psychosocial therapies to treat problems in children *prior to considering medication* is an *ethical priority*

74

“First do no harm”

Use of psychotropic medications that have been *reported to have serious adverse effects in children—including death—should be halted* until research demonstrates that both short- and long-term benefits outweigh the already known risks



Avoid psychotropic drug use in young children until

- ✓ evidence-based psychosocial interventions have been exhausted
- ✓ rationally-anticipated benefits outweigh the likelihood of risks
- ✓ parents/guardians are fully informed
- ✓ close monitoring is in place


(Vitiello, 2001)

76


A Critical Curriculum
on Psychotropic Medications

Module 6

The End




A Critical Curriculum on Psychotropic Medications



A Critical Curriculum on Psychotropic Medications


- **Principal Investigator:**
 - David Cohen, Ph.D.
- **Research Coordinator:**
 - Inge Sengelmann, M.S.W.
- **Professional Consultants:**
 - David O. Antonuccio, Ph.D. (psychology)
 - Kia J. Bentley, Ph.D. (social work)
 - R. Elliott Ingersoll, Ph.D. (counseling & psychology)
 - Stefan P. Kruszewski, M.D. (psychiatry)
 - Robert E. Rosen, J.D., Ph.D. (law)
- **Flash production and design:**
 - Sane Development, Inc., and Cooper Design, Inc.
- **Voice narration and Flash editing:**
 - Saul McClintock



CriticalThinkRx was made possible by a grant from the Attorneys General Consumer and Prescriber Grant Program, funded by the multi-state settlement of consumer fraud claims regarding the marketing of the prescription drug Neurontin®


Module 7

Medication Management: Professional Roles and Best Practices



Part A

Non-medical roles and medication management




Historical roles of non-medical helpers

To serve as resources for physicians and allied professionals:

- First, giving clients information about their medications;
- Then, identifying obstacles to compliance;
- Later, advocating for clients

(Bentley, Walsh, & Farmer, 2005) 6


A 2001 national survey of clinical and mental health social workers identified 31 possible tasks and activities related to medication



(Bentley, Walsh, & Farmer, 2005)

Survey found some tasks “frequently” performed with clients


- ✓ Discussing clients’ feelings about taking medications
- ✓ Making referrals to physicians
- ✓ Discussing how medications may work with other interventions



(Bentley, Walsh, & Farmer, 2005)

Tasks “often” performed with clients


- ✓ Helping weigh pros and cons of taking medication
- ✓ Monitoring clients’ compliance with medication
- ✓ Discussing medication problems



(Bentley, Walsh, & Farmer, 2005)

Tasks “rarely” performed

- ✓ Assessing and documenting adverse effects
- ✓ Educating about medications
- ✓ Suggesting changes in medications to physicians




(Bentley, Walsh, & Farmer, 2005)

Assuming roles is complicated by:


- ✓ priority of some professional values and ethics, such as client’s right to self-determination
- ✓ questions about validity of medical model for explaining human distress
- ✓ gaps and uncertainties in evidence about medications
- ✓ influence of pharmaceutical companies on the entire mental health system

(Walsh, Farmer, Taylor & Bentley, 2003)

Increasing demands to regulate medicated clients clash with professional values, creating a “professional dissonance”




(Taylor & Bentley, 2005)



Public and professional attitudes

Overall, the public does not embrace psychiatric medications as a solution to children's problems

- 70% of adult Americans refuse to use medication for children labeled "oppositional" or "hyperactive"
- Only 10% see medication as the most effective component of treatment, and 66% believe it is used as a substitute for other interventions




(McLeod, et al. 2004)

Practitioners divided

Some find drug treatment of youth helpful or essential

Others find drugs used as a form of social control, misused as a remedy for frustrated parents or overtaxed system, or ineffective




(Moses & Kirk, 2006)

15

Helping parents find solutions

When faced with a distressed child, parents may perceive few options in a world where insurers, medical providers and schools pressure them to medicate their children



(McLeod et al., 2004)

16

Unbiased sources of information

Non-medical professionals should serve as "unbiased sources of information" to help parents find the right solutions for their children and to promote alternatives based on critically-evaluated evidence

(Bradley, 2003; Buccino, 2006; McLeod et al., 2004)

17

"Vigilant and critically minded"

Non-medical professionals are urged to maintain an "*informed but critical*" stance by developing adequate knowledge about the benefits and adverse effects of psychotropic drugs, and remain "*vigilant, and critically minded*"

(Moses & Kirk, 2006, pp. 220-221)

18

Yet be familiar with basic psychopharmacology

including uses, side effects, dosages, and drug interactions in order to be effective in this complex environment

(Bradley, 2003; Buccino, 2006)

19

Part B

Evolving roles in medication management



In today's *collaborative, multi-disciplinary environment*, non-medical practitioners are called upon to play many roles on behalf of clients taking medication



21

Physician's Assistant

Traditionally the most common role for professionals legally limited in their scope of work with medications, they

- Help clients follow doctor's recommendations
- Not expected to give advice about decisions involving the prescription



(Bentley & Walsh, 2006)

Consultant

- Evaluates client to assess for referral to physicians
- Prepares clients to talk with the prescribing physician
- Monitors client's subjective experience of medication
- Assesses client's ability to pay for expensive drugs

(Bentley & Walsh, 2006)

23

Counselor

- Coaches and teaches by providing information and advice about medications
- Teaches problem solving, helps identify alternatives, assists in making decisions



(Bentley & Walsh, 2006)

24

Monitor

Helps client observe positive and negative effects of medication
Evaluates client's medication responses, in psychological, interpersonal, and social realms, and effects on self-image and identity
Discusses the monitoring process with clients, families and physicians

(Bentley & Walsh, 2006)

25

Advocate

Presents client's expressed wishes to those in the medical or mental health system
Ideally, has a peer relationship with the physician and participates in all phases of medication decision-making
Possesses knowledge of psychopathology, medications, and related laws and regulations



(Bentley & Walsh, 2006)

26

Teacher

Provides educational materials and other information to clients about:

- The purposes, actions and effects of medications
- Problem-solving regarding medication issues and adverse effects
- Practical suggestions to help clients take medication appropriately

(Bentley & Walsh, 2006)

27

Researcher

Conducts and publishes research in medical and non-medical literature about the full range of psychotropic medication issues



(Bentley & Walsh, 2006)

28

An emerging clinical role: *easing clients off meds*

Helping clients withdraw from psychiatric drugs or helping simplify medication regimen
Contingent on practitioner competence and a "rational, person-centered" approach
Guidelines exist for non-medical practitioners to recognize and address discontinuation effects

(Cohen, 2007; Meyers, 2007; Rivas-Vasquez et al., 1999)

29

Effective collaboration with clients, physicians and other providers of care



Traditional

Reflects dominance of medical profession
 Characterized by limited, unclear or subservient roles of non-medical professionals



(Bentley & Walsh, 2006; Bronstein, 2003)

Interdisciplinary

Improves services to the client and work satisfaction for professionals
 - May not translate in all environments and training in effective models is needed

(Bentley & Walsh, 2006; Bronstein, 2003)

32


Transformational

Enhances the contributions of all members of a team
 Assumes non-hierarchical relationships where physicians integrate psychosocial aspects of care and involve non-medical professionals in decision-making

(Bentley & Walsh, 2006; Bronstein, 2003)

33

Components of an Interdisciplinary Model



(Bronstein, 2003)

34

Elusive qualities of successful collaboration?

- A favorable political and economic climate
- Shared vision, attainable goals
- Open and frequent communication
- Trust, adaptability, respect
- Clear roles but flexibility in assuming them
- Competent, well-trained practitioners
- A leader with strong interpersonal skills

Unfortunately, these qualities may be absent in interdisciplinary settings

(Bentley & Walsh, 2006; Bronstein, 2003)

35

Collaboration to enhance client's self-determination

Collaboration between clients, families and professionals as *partners* in the helping process is key to respecting the client's right to self-determination

When partnership with other professionals is difficult, focus should be on empowering clients with information so that they make choices in collaboration with prescribers

(Bentley & Walsh, 2006; Cohen, 2007; Slavin, 2004; Weene, 2002)

36

Needed—but difficult to accomplish: A balance between...

- ✓ the rights of individuals, families and society
- ✓ the costs and benefits of using psychotropic medication
- ✓ the non-medical practitioner’s role in medication management and the legitimacy and uniqueness of other helping professions

(Bentley & Walsh, 2006) 37



Integrating drugs and psychosocial treatment introduces complex dynamics that require attention and management

Managing parallel treatment requires navigating

- ✓ the relationships among client, prescriber and therapist
- ✓ competing ideologies held by providers

(Bentley & Walsh, 2006; Bradley, 2003) 39

Dimensions of partnership in medication management

Dimension	Traditional model	Partnership model
Goals of medication	Reduce symptoms	Improve quality of life; emphasis on client priorities
Who selects medication	Physician provider	Client collaboration to help define options
Education focus	Increasing compliance	Improving client’s ability to manage recovery
Monitoring and evaluating	Physician evaluates clinical status and compliance	Client and providers evaluate range of outcomes and options
Self-care by client	Largely ignored in mental health	Integrated into consultations with client and family
Control and status	Providers control processes and hold status positions	Emphasis on client control, and client’s experiences valued
Refusal and reluctance	Seen as related to denial and paranoia	Seen as a right to be respected in all but emergency situations

(Bentley & Walsh, 2006, p. 223) 40

Part C

Tools for Competence

Assessments, Referrals, Court Affidavits and Medication Monitoring

Comprehensive assessments

Understanding the person in the context of their experiences

Working Definition

An ongoing, systematic data collection about a client's functioning

A process of problem selection and specification guided by a person-in-environment, systems orientation

(Jordan & Franklin, 2003)

43

An individualized process

views the whole person in context, including all factors contributing to their distress and strengths, and changes required to improve coping and mastery

- the person's own perspective is key to understand their situation

(Austrian, 2005; Jordan & Franklin, 2003)

44

Elements of assessment

1. **Exploration** of client's unique story and facts
2. **Inferential thinking** to evaluate meaning of the facts of their story
3. **Evaluation** to assess client functioning, strengths and weaknesses in context
4. **Problem definition** based on the first three steps and in collaboration with client
5. **Intervention planning** based on preceding four steps and in context of environment

(Austrian, 2005; Jordan & Franklin, 2003)

45

Mental status examination

- Appearance, speech, attitude, motor behavior
- Mood, range and appropriateness of affect
- Hallucinations, depersonalization, derealization
- Remote, recent, and immediate memory
- Level of consciousness, orientation
- Impulse control
- Judgment and insight

(Austrian, 2005; Jordan & Franklin, 2003)

46

"Integral" assessment approach requires knowledge of

- the client's experience (the individual viewed subjectively/from within)
- the client's behavior (the client viewed objectively/from without)
- the client's culture (the client's system viewed subjectively/from within)
- the client's social system (the client's system viewed objectively/from without)

(Marquis, 2008; Ingersoll, 2002)

47

Referrals

Best practices in referring clients for psychiatric evaluation

Few empirical evaluations

Few researchers have investigated effective referral practices, despite frequency of this activity

Tentative guidelines are offered



(Bentley, Walsh & Farmer, 2005)

49


Quality referrals

1. Establish and maintain collaborative relationships with prescribers
2. Share *up-to-date* information about medications with clients and families
3. Help clients and families articulate and manage the meaning of medication
4. Prepare clients and families for the medication evaluation
5. Follow up on the referral
6. Manage legal and ethical concerns

(Bentley, Walsh & Farmer, 2005)

50

- Prescription
- Reason for the prescription
- Expectations of benefit
- Probability of benefits
- Alternative treatments available
- Risks of the medication
- Expenses involved (direct/indirect)
- Decision



(Chewning & Sleath, 1996, in Bentley & Walsh, 2006)

51

A medication evaluation should be requested only if the child's symptoms do not improve or worsen significantly after good psychosocial interventions have been attempted



52

If drugs are considered, all practitioners should evaluate if there is clear evidence of favorable benefit-to-risk ratio

Drugs unapproved for that age group cannot be recommended without special consideration



53

Affidavits to judges regarding medication suggestions for children in state care
A recommended checklist



Psychosocial situation and stressors

1. Describe the observed behaviors of concern & who has observed them, when and where
2. Describe past, recent, or chronic stressors in the child's life that may be contributing to any of the observed behaviors

55

Psychosocial assessment

3. Summarize the results of your own assessment of this child's situation: what, in your judgment, could explain how this child is now acting?
4. If the child has been on medication, could the symptoms be adverse effects of the medication? List sources to justify your conclusion

56

Assessment of interventions

5. Describe any previous interventions to address the problems identified in your assessment
6. Describe how these interventions have been evaluated, and their results
7. What other interventions might address this child's problems? To what extent are they available for this child? Why or why not?

57

Medication history

8. List medications (names, dosages, times per day) the child takes now and over the past 2 years
9. Have you participated in evaluating the child's progress on medication? What specific goals have been expected, how has their attainment been evaluated?

58

Medication monitoring, evaluation

10. Have you evaluated for adverse effects, behavioral or other? Have you used any rating scales? How well, in your own careful, overall judgment, is this child tolerating his or her medication?

59

Informed consent

11. Do you have any information on this child's attitude to the medication?
12. How have the risks and benefits of the medication, as well as those of alternate interventions, been assessed and discussed with parents or caregivers?

60

Future monitoring

- 13. If the child is placed on medication, describe your specific role in monitoring its effects.
- 14. What reasons do you have to expect that the proposed medication will be beneficial to this child?

61

Medication monitoring



Attending to anticipated and unanticipated effects

Monitoring helps clients and families

- Keep track of medication effects
- Cope with bothersome effects
- Solve medication-related issues
- Make decisions about treatment using critically-evaluated information
- Prevent medication errors

(Shojania, 2006)

63

Clients may not know

Clients typically fail to link behavioral drug effects to their drug, and may incorrectly believe they are suffering from additional unrelated physiological or psychological symptoms

Do not dismiss unusual effects, watch out for amplified usual effects, and educate clients about risk of “prescribing cascade”

(Otis & King, 2006)

64

Formal monitoring essential

Without formal monitoring, only a fraction of drug problems are recognized

Structured medication reviews have been shown to be more valid and improve client’s quality of life

(Otis & King, 2006; Greenhill et al., 2004; Jordan et al., 2004; Kalachnik, 1999)

65

Tools for monitoring

Drug effect checklists — existing or individualized for client’s situation (*see checklist handout in website*)

- Use before starting the medication
- Use after starting the medication



(Jordan et al., 2004)

66

Adapted from: Kafetsios JE. Measuring side effects of psychopharmacologic medications. *Mental Rehabilitation & Recovery, Disability Research Review* 1995; 38:203-31; Rasmussen et al. 2002. Discontinuation syndrome. *Biological Psychiatry* 1995; 44:74-87. (3) Beethoven/Gulcher & Jeffrey, 2005. *Clinical handbook of psychotropic drugs* (10th ed). Seattle: Hogrefe.

CRITICAL THINK Rx

MEDICATION MONITORING—ADVERSE EVENT CHECKLIST


Client's name: _____ Date of assessment: _____
 Assessor: _____

Drug(s) and dosage: _____

Instructions: Fill out once a month or more, before, during, and for 3 months after medication use. Inquire about the presence of each event over the past 7 days. If present, score as 1 (mild), 2 (moderate), or 3 (severe). If not present, leave blank. For items listing different or opposite events (e.g., "increased" or "decreased" appetite, circle the appropriate one.)

Psychological	1, 2, 3	Gastrointestinal	1, 2, 3
1. Agitation (restless, nervous, hyperactive)		43. Increased or Decreased appetite	
2. Confusion, cognitive difficulties		44. Weight Gain or Loss	
3. Memory problems, forgetfulness		45. Abdominal pain or cramps, Stomach bloating	
4. Irritability (easily upset, angry)		46. Increased thirst	
5. Irritability		47. Nausea, vomiting	
6. Trouble concentrating or paying attention		48. Diarrhea	
7. Insomnia, trouble falling or staying asleep		49. Constipation	
8. Hypersomnia, trouble waking up		50. High blood sugar	
9. Crying spells, sadness		51. Other: _____	
10. Anxiety, tension, Panic (racing heart, breathless)		Musculoskeletal/Neurological	
11. Lethargy, apathy, drowsiness, agitation		52. Disequilibrium, unsteady gait, poor coordination	
12. Nightmares, intense dreaming		53. Spinning, swaying, lightheaded	
13. Feeling detached or unreal		54. Weakness, fatigue	
		55. Numbness, burning or tingling sensations	
		56. Slowed movements, muscle rigidity	
		57. Muscle cramps, stiffness, twitches, jerks	
		58. Restlessness, pacing, rocking, can't sit still	
		59. Tremor (slight shaking/ trembling of limbs or muscles)	
		60. Any abnormal involuntary movements	
		61. Other: _____	
		Skin	
		62. Increased or Reduced sweating	
		63. Increased sensitivity to sun	
		64. Chills or feelings of warmth	
		65. Rash, hives / Dry skin, crusty	
		66. Acne	
		67. Easy bruising	
		68. Pale, yellowing skin	
		69. Hair loss or Abnormal hair growth	
		70. Other: _____	
		Genito-Urinary	
		71. Menstrual disturbances (absent or irregular periods)	
		72. Difficulty urinating / Increased urination	
		73. Erection, night bedwetting	
		74. Difficulties with orgasm	
		75. Erectile dysfunction	
		76. High or low sexual desire / activity	
		77. Other: _____	
		Cardiovascular	
		78. High blood pressure	
		79. Arrhythmia (irregular heartbeat)	
		80. Tachycardia (abnormally fast heartbeat)	
		81. Dizziness	
		82. Fainting	
		83. Rapid heart rate	
		84. Slow heart rate	
		85. Swelling	
		86. Tiredness	
		87. Weakness	
		88. Weight gain	
		89. Weight loss	
		90. Other: _____	

Systematic monitoring must be carried out to evaluate the wide-ranging effects of medications on behavior, mood, as well as physical and emotional development



68

Children should be evaluated for

- Emotional development (to examine whether the drug induces or worsens certain problems)
- Cognitive development
- Physical growth (i.e., weight and height)
- Pubertal development (to examine drug effects on course of puberty)

(Greenhill et al., 2003)

69

Medication guidelines for child welfare

Medication should only be used as part of a comprehensive treatment plan integrating behavioral interventions

- not used in lieu of other treatments or supports
- based on adequate information, including full biopsychosocial and medical assessment
- resting on informed consent

(Bellonci & Henwood, 2006)

70

With children (after rock-solid justification for medication has been provided)

- ✓ adjust doses to a minimum to minimize side effects
- ✓ periodically attempt to take child off medication
- ✓ avoid polypharmacy
- ✓ continually reassess risk-to-benefit ratio

(Bellonci & Henwood, 2006)

71


Medical monitoring schedule

Children on psychotropic medications should be seen no less than every three months *at a bare minimum*

FDA guidelines for antidepressants require more frequent monitoring due to risks

(Bellonci & Henwood, 2006)

72




**Red flags:
Additional
monitoring concerns**

- Children under five years of age
- Children on 2 or more medications
- Children in state custody


(Bellonci & Henwood, 2006) 73

“Psychotropic medications for young children should be used only when anticipated benefits outweigh risks. Parents should be fully informed and decisions made only after carefully weighing these factors. Children and adolescents must be carefully monitored and frequently evaluated as the side effects common to some medications are particularly difficult for children.”

National Alliance for Mental Illness (NAMI)
Policy Research Institute, 2004



Part D



**Conclusions and
Recommendations**

Beyond biology...

...medications affect the psychological and social concerns of clients, leading non-medical providers to be increasingly involved in medication issues

76

What is needed?

Education and training about psychiatric medications for non-medical professionals

Guidelines regarding responsibilities with respect to medication, including dealing with ethical and legal issues such as obligations to report adverse effects

Improved collaboration with *clients as partners* and with medical providers as part of interdisciplinary teams—though key concern remains empowering clients to make their own decisions

77

Training on

- the impact of meanings of medication-taking
- monitoring clients for adverse effects
- skills in educating clients about risks and benefits of psychotropic medications
- finding and critically evaluating research on specific medications
- understanding the strong ideological, economic and political influences on prescription writing in the U.S.

78

Research on

- ☑ how medications and psychosocial interventions interact
- ☑ how medications affect child's self-control, self-image, and personal responsibility (autonomy)
- ☑ how medications affect therapeutic relationships

79


A Critical Curriculum on Psychotropic Medications

Module 7

The End



80



A Critical Curriculum on Psychotropic Medications

A Critical Curriculum on Psychotropic Medications

Principal Investigator: **Research Coordinator:**

- David Cohen, Ph.D.
- Inge Sengelmann, M.S.W.

Professional Consultants: **Flash production and design:**

- David O. Antonuccio, Ph.D. (psychology)
- Sane Development, Inc., and Cooper Design, Inc.
- Kia J. Bentley, Ph.D. (social work)
- Voice narration and Flash editing:**
- R. Elliott Ingersoll, Ph.D. (counseling & psychology)
- Saul McClintock
- Stefan P. Kruszewski, M.D (psychiatry)


– Robert E. Rosen, J.D., Ph.D. (law)



CriticalThinkRx was made possible by a grant from the Attorneys General Consumer and Prescriber Grant Program, funded by the multi-state settlement of consumer fraud claims regarding the marketing of the prescription drug Neurontin®



Module 8

Alternatives to Medication: Evidence-Based Psychosocial Interventions



Part A

What is Evidence-Based Practice?

A philosophy *and* a process designed to unite research and practice in order to

maximize chances to help clients
minimize harm to clients (in the name of helping)



(Gambrill, 2006)

Deeply participatory

EBP is “anti-authoritarian”—it urges all involved to question claims about what is known *and unknown* about treatments

(Gambrill, 2006)

8

EBP difficulties

- ☑ Threats to business-as-usual
- ☑ Limited training and supervision
- ☑ Concerns about cultural sensitivity
- ☑ Worries that “cook book” methods mask real-world complexity

(Barratt, 2003; Chorpita et al. 2007; Duncan & Miller, 2006)

9

An intervention should have *at least some* unbiased observations or tests supporting its usefulness with particular problems and clients

10

Some criteria for judging an intervention

- ☑ Sound theoretical basis
- ☑ Low risk for harm
- ☑ *Unbiased* research exists
- ☑ Therapist and client concur

11

Available “evidence” no guarantee of usefulness

Published evidence is influenced by funding sources, researcher biases, and conventional wisdom

Statistically significant differences between treatment groups means simply that more clients in one group had some type of response (partial to complete)

(Hoagwood et al. 2001; Ingersoll & Rak, 2006)

12

However, on average, *all major therapies produce equivalent results.*

Clients' improvement may result from *factors common to every therapy*

(Etkins, 2007; Hubble, Duncan, & Miller, 1999) 13

Most improvement has little to do with therapy or technique

Factor	% improvement explained
Client + outside therapy factors	87
Client-therapist alliance	8
Therapist allegiance to model	4
Therapist technique	1

(Hubble, Duncan, & Miller, 1999; Wampold, 2001) 14

Healthy skepticism

“We would do well ... to remain optimistically humble on the matter of evidence-based practices in mental health” by accepting that all assumptions are “provisional and reversible”

(Norcross, Beutler & Levant, 2006, p. 11) 15


A clinician's “rubric” for EBP

“Adhere when possible, adapt when necessary, assess along the way”

(Amaya-Jackson & DeRosa, 2007, p. 388) 16


Choosing proper interventions rests on

- ☑ a clear understanding of the problem from a person-in-situation perspective
- ☑ addressing the complexity of the problem
- ☑ a policy of “First, do no harm”



Part B

Deconstructing the Diagnosis:



What is this child's problem in behavioral terms?

Bio-psycho-social or bio-bio-bio?

- ✓ Complex problems in living reduced to “brain disorders”
- ✓ Complex life events reduced to “triggers”
- ✓ Medicalization of distress and disability leading to false hopes of “quick fix” via pills

(Read, 2005)

19

We often ignore environmental influences on behavior

- ☑ Poor parenting, neglect, abuse
- ☑ Schools’ failure to motivate children
- ☑ Poverty, lack of access to resources
- ☑ Violence in media, society, neighborhood
- ☑ Culture’s emphasis on instant gratification
- ☑ Drug culture (“take,” not “talk”)
- ☑ Lack of tolerance for differences

(Bentley & Collins, 2006)

20

Children’s distress: “Disorders” or complex adaptations to distressing life experiences?



By seeing children as real persons with their own view of their situation, one ascribes a different meaning to their behavior

(Donovan & McIntyre, 1990)

21

“Understanding” rather than “diagnosing”

A developmental-contextual approach views actions as “communicative”: attempts by individuals to cope, adapt, struggle with their life experiences



(Donovan & McIntyre, 1990)

22

Here’s a list of feelings and behaviors from DSM-IV-TR criteria of “disorders” commonly diagnosed in children

Note the similarities...

“Attention-Deficit/Hyperactivity Disorder (ADHD)”

Feels:

- Angry, irritable, frustrated



Acts:

- Fidgets, squirms
- Easily distracted, forgetful (difficulty thinking, concentrating)
- Interrupts others (acts impulsively)
- Acts aggressively

24

“Major Depressive Disorder”

Feels:

- Sad, empty
- Afraid, anxious
- Angry, irritable, frustrated



Acts:

- Eats, sleeps too little (or too much)
- Moves, speaks slowly
- Acts impulsively
- Acts aggressively
- Easily distracted (difficulty thinking, concentrating)

25

“Anxiety Disorder”

Feels:

- Afraid, anxious
- Angry, irritable, frustrated



Acts:

- Cries, throws tantrums
- Freezes, clings
- Fidgets (psychomotor agitation)

26

“Conduct Disorder”

Feels:

- Angry, irritable, frustrated, hostile



Acts:

- Bullies and threatens
- Fights
- Steals, lies
- Runs away
- Destroys property

27

“Oppositional Defiant Disorder”

Feels:

- Angry, irritable, frustrated, hostile



Acts:

- Disobedient
- Loses temper
- Argues with adults
- Annoys people
- Refuses to follow rules

28

“Bipolar Disorder”

Feels:

- Alternating sad and euphoric
- Alternating fearful and reckless
- Angry, irritable, frustrated

Acts:

- Easily distracted (difficulty thinking, concentrating)
- Moves, speaks fast (agitation)
- Acts impulsively
- Acts aggressively
- Does not sleep well

29

“Psychotic Disorder”

Feels:

- Sad, empty
- Blunted feelings, expressionless
- Angry, irritable, frustrated
- Afraid, anxious

Acts:

- Apathetic
- Refuses to speak
- Dresses inappropriately
- Cries frequently
- Sees or hears things

30

“Post-Traumatic Stress Disorder”

Feels:

- Sad
- Afraid, anxious
- Angry, irritable, frustrated
- Helpless, guilty, shameful

Acts:

- Agitated, impulsive, re-enacts trauma
- Hypervigilant: distrustful, withdraws
- Dissociated: forgets and can't focus



31

“Reactive Attachment Disorder”

Feels:

- Afraid, anxious
- Angry, irritable, frustrated

Acts:

- Watchful, frozen
- Avoids attachments
- Seeks approval or can't be comforted
- Disregards danger cues



32

The common elements

Experiencing negative emotions
(sadness, fear, anger, irritability)

Difficulty controlling oneself
(impulsivity, aggression, inattention)

Seeing self and world negatively
(hopelessness, helplessness, shame, withdrawal)

33

What are we medicating?

Negative emotions leading to disruptive actions—especially under stressful conditions that tax the child's adaptive capacities

(Schore, 1994, 2003)

34

Most commonly medicated

Impulsive aggression
“a key therapeutic target across multiple disorders”



(Jensen et al. 2007, p. 309)

35

DSM's scientific value seriously challenged in all disciplines

- ✓ internal inconsistency in the manual (rejects categorical approach in intro but then lists 300+ categories)
- ✓ overlap between categories leads to “comorbidity”—with no increase in understanding
- ✓ persistent problems of unreliability, especially with children's diagnoses
- ✓ lack of fit between categories and empirically observed symptom clusters

(Caplan, 1995; Duncan et al. 2007; Maj, 2005; Kirk & Kutchins, 1992, 1994; Jacobs & Cohen, 2004; Mirowsky & Ross, 1990)

36

More recent DSM critiques...

- ✓ more behaviors now seen as “mental disorders” (from 106 in 1952 to 365 in 1994)
- ✓ political lobbying determines inclusion or exclusion of diagnoses
- ✓ all DSM task force members on mood and psychotic disorders tied to drug industry
- ✓ practitioners focus on diagnosis rather than client, losing client’s actual story
- ✓ still no “gold standard” validity—no specific bio-marker linked to *any* disorder

(Andreasen, 2006; Tucker 1998; Charney et al. 2005; Kutchins & Kirk, 1998)

37

Critical list of DSM “accomplishments”

- ☑ increases people’s interest to classify psychosocial problems as medical disorders
- ☑ Helps justify more studies to see how many people can fit how many DSM categories (which often change)
- ☑ led to modest increase in diagnostic reliability since 1980
- ☑ now used by most practitioners in main schools of thought—mostly to obtain third-party reimbursement?
- ☑ brings financial revenues to the American Psychiatric Association from sales of DSMs and training materials
- ☑ strengthened psychiatry’s leadership in mental health system (as official definer of mental distress)

38

Part C

Empirically-supported psychosocial interventions for children and adolescents



**Focus:
Trauma, Resilience and Child Welfare**



Trauma and early loss

For thousands of children every year, loss and trauma due to disrupted attachments to biological parents result in foster care placements

(Jones Harden, 2004; Racussin et al. 2005)

41


Additional, placement-related traumas

- ✓ Emotional disruption of out-of-home placement
- ✓ Adjusting to a foster care setting
- ✓ Relative instability of foster care
- ✓ High turnover of workers

(Jones Harden, 2004; Racussin et al. 2005)

42

Neurobiology of attachment



Brains develop in a *socially dependent manner*, through secure attachments and *consistent, competent* adults attuned to the needs of the child

(Schore, 1994, 2001, 2003; van der Kolk, 2003)

43

<p><u>Child's "job"</u>: to form close, trusting attachments with caregivers</p>		<p><u>Adolescent's "job"</u>: to expand attachments using secure base with caregivers</p>
--	---	---

(Gunnar et al. 2006; Mash & Barkeley, 2006; Moran, 2007; Wolfe & Mash, 2006)

44

Trauma, abuse, and neglect

- ☑ disrupt a child's ability to form secure attachments
- ☑ impair brain development and regulation
- ☑ make self-control difficult
- ☑ alter identity and sense of self

(Bowlby, 1988; Cook et al. 2005; Courtois, 2004; Creeden, 2004; Jones Harden, 2004; van der Kolk, 1994)

45



Resilience

The ability to function well despite living or having lived in adversity rests mainly on normal cognitive development and involvement from a caring, competent adult

(Agaibi & Wilson, 2005; Masten et al. 1990; Schofield & Beek, 2005)

46

- ✓ Risk and protective factors in the foster child, foster-families, agencies, and birth family interact to produce upward or downward spirals
- ✓ Understanding resilience helps create interventions that produce positive turning points in children's lives

(Schofield & Beek, 2005)

47

Three key elements

1. Secure base: is child strengthening sense of security and able to use foster-parents as a secure base?
2. Sense of permanence: is placement stable and foster-parents offering family membership?
3. Social functioning: is child functioning well in school, with peers?

(Schofield & Beek, 2005)

48

Treatment goals


- ✓ Enhance sense of personal control and self-efficacy
- ✓ Maintain adequate level of functioning
- ✓ Increase ability to master, rather than avoid, experiences that trigger intrusive re-experiencing, numbing, and hyper-arousal

(Ford et al. 2005; Kinniburgh et al. 2005) 49

What could help?

Activating child's internal reparative mechanisms through *dyadic interventions* and creating secure attachments

- dyadic therapy mobilizes the completion of interrupted biological and emotional developmental processes



(Amaya-Jackson & DeRosa, 2007; Courtois, 2004; Ford et al. 2005; Pearlman & Courtois, 2005) 50

A sensorimotor approach

Children's internal stimuli, can trigger dysregulated arousal, causing emotions to escalate

- Integration of cognitive, emotional and sensorimotor levels is crucial for recovery

(Ogden, 2006) 51


Why would this help?

Child develops the ability to take in, sort out, process, and interrelate information from the environment – leading to self-organization of internal states and self-control of behavior

(DeGangi, 2000; Kinniburgh et al. 2005; Schore, 2003; van der Kolk, 2006) 52

How would this help?

By enhancing children's:



- ✓ social skills
- ✓ ability to understand and express feelings
- ✓ ability to cope with anger and distress
- ✓ ability to problem-solve and think helpful thoughts
- ✓ skills to self-direct and create goals

(Bloomquist, 1996; Kinniburgh et al. 2005) 53

Alternatives to medication


- ☑ Consistent, structured, supportive adult supervision
- ☑ Opportunities for self-expression and physical activity, to give children a sense of mastery over their minds and bodies



(DeGangi, 2000; Faust & Katchen, 2004) 54

Helpful activities


- ☑ Teaching problem-solving and pro-social skills
- ☑ Modeling appropriate behaviors
- ☑ Teaching self-management
- ☑ Helping children learn to comply and follow rules



(DeGangi, 2000; Faust & Katchen, 2004) 55

Helpful interactions


- ☑ Desensitizing hyper-reactivity
- ☑ Promoting self-calming and modulation of arousal states
- ☑ Organizing sustained attention
- ☑ Facilitating organized, purposeful activity



(DeGangi, 2000)

Expected outcomes

Children learn to develop appropriate responses, self-organization and control, which in turns leads to




MASTERY AND SELF-ESTEEM

(Kinniburgh et al. 2005) 57

Many treatment alternatives


Symptom-focused: Behavioral, cognitive-behavioral, and interpersonal therapies, attachment-based therapies, trauma-focused therapies

System-focused: Treatment foster care (TFC), Multi-dimensional treatment foster care (MTFC)



(Farmer et al. 2004; Racussin et al. 2005)

Focus: Dysregulated “moods”




“Depression” and “Anxiety”



60

The New York Times
Talk Therapy Pivotal for Depressed Youth



February 6, 2007
In Rigorous Test, Talk Therapy Works for Panic Disorder
By [BENEDICT CAREY](#)

61

Link to child maltreatment

Abuse leads to “hypervigilance” to threat, resulting in anxiety and hopelessness

Neglect results in dysregulated “moods”

(Greenwald, 2000; Lee & Hoaken, 2007)

62

“Traumatized children tend to communicate what has happened to them ... by responding to the world as a dangerous place by activating neurobiologic systems geared for survival, even when objectively they are safe”

(van der Kolk, 2003, p. 309)

Therapy or no therapy?

Some 30-40% recover without intervention

Approximately 50% of treated patients improve within 8 weeks

A friendly sympathetic attitude and encouragement are key

(Roth & Fonagy, 1996)

64

Consensus strongly favors cognitive-behavioral therapy (CBT) as **first-line treatment above medications**

(APA Working Group, 2006; March, 1995; Roth & Fonagy, 1996; Velting et al. 2004)

65

Other effective interventions

1. Interpersonal psychotherapy
2. Psychodynamic psychotherapy
3. Exposure-based contingency management
4. Problem-solving and coping-skills training

(APA Working Group, 2006; Roth & Fonagy, 1996)

66

Patient preference

When given a choice, patients express a preference for psychosocial interventions over medications



(APA Working Group, 2006)

67

“Bipolar Disorder” and “Schizophrenia”

68

Very rare in children (~1%)

Diagnosis controversial:

- no laboratory “test”
- “symptoms” may be manifestations of ordinary developmental differences

(Birmaher, 2003; Birmaher & Axelson, 2006; Cepeda, 2007; Correll et al. 2005; Danielson et al. 2004; Irwin, 2004; Findling, Boorady & Sporn, 2007; Roth & Fonagy, 1996)

69

High risk of over-diagnosis

NIMH Review: 95% of 1500 children referred for high clinical suspicion of childhood-onset schizophrenia did not meet DSM criteria after careful inpatient observation off all medications

No evidence that they would have developed psychosis if left untreated

(Shaw & Rapoport, 2006)

70

Link to child maltreatment

Child abuse and neglect considered a causal factor for psychosis and “schizophrenia”

- Content and severity of psychotic symptoms related to severity of past abuse

(Cepeda, 2007; Morrison et al. 2005; Read & Ross, 2003; Read et al. 2004, 2005)

71

Many children improve when treated with family-based psychosocial interventions, *even without medications*

- High rates of “relapse” observed on medication

(Birmaher, 2003; Birmaher & Axelson, 2006; Cepeda, 2007; Correll et al. 2005; Danielson et al. 2004; Findling et al. 2007; Irwin, 2004; Roth & Fonagy, 1996)

72

Effective psychosocial treatments

Child- and Family-Focused CBT combined with interpersonal and “social rhythm” therapy to stabilize mood, activities and sleep

Community support and social acceptance through day programs and sports/cultural activities

(Findling et al., 2007) 73

Who recovers and why?

Psychiatric literature is mostly silent about the characteristics of people who fully recover from psychosis and how and why they do so

(Siebert, 2000) 74

Focus:
Disruptive behaviors



Disruptive behaviors:
the most frequent reason for referral of children to mental health services

(Brestan & Eyberg, 1998; Butler & Eyberg, 2006) 76

For disruptive behaviors and conduct “disorders”

Family-based behavioral interventions



(APA Working Group, 2006; Brestan & Eyberg, 1998; Diamond & Josephson, 2005; Kazdin, 2005, 2000, 2000b; Kazdin & Weisz, 2003; Thomas, 2006)

The New York Times (2006, December 22)

TROUBLED CHILDREN
Parenting as Therapy for Child's Mental Disorders



TJ Van de Walle's attention deficit problems have improved in response to parenting techniques, his mother Dawn, right, said.

78

Effective parenting: the most powerful way to reduce child and adolescent problem behaviors



(Caspe & Lopez, 2006; Johnson et al. 2005; Kumpfer et al. 2003) 79

Strongest evidence base

1. Parent management training (PMT)
2. Problem-solving skills training (PSST)
3. Brief strategic family therapy (BSFT)
4. Functional family therapy (FFT)

(Brestan & Eyberg, 1998; Butler & Eyberg, 2006; Farley et al. 2005; Kazdin, 2003; Kazdin & Whitley, 2003; Springer 2006; Thomas, 2006)

80

Goals of parent training

- ☑ Promote parent competencies & strengthen parent-child bonds
- ☑ Increase consistency, predictability & fairness of parents
- ☑ Produce behavior change in children



(Kazdin, 2003; McCart et al. 2006; Webster-Stratton & Reid, 2003) 81

“Problem” children or “problem” adults?

Coercive parenting was the only factor linked to children’s failure to improve their conduct after family treatment

(Webster-Stratton, Reid & Hammond, 2001)

82

Maltreatment consistently linked to aggressive behaviors

- ☑ History of trauma virtually *universal* in youth with conduct “disorders”

(Greenwald, 2000; Lee & Hoaken, 2007)

83

Children in foster care

- ✓ have socio-emotional problems **3 to 10 times more often** than other kids
- ✓ Coercive interactions only result in escalation of aggressive behaviors



(Nilsen, 2007)

84

Parent-training in child welfare

Promising programs exist to train biological and foster parents

Goal is to break the cycle of coercive parenting and child oppositional behavior

(Barth et al. 2005; Nilsen, 2007)

85

“ADHD”

Large evidence base exists for behavioral interventions, incl. parent training, social skills training, and school-based services

- Results equivalent to stimulant medications without the health risks



(APA Working Group; Chronis et al. 2004, 2006)

86

Focus: Mentoring



Children’s development depends upon reciprocal activity with others with whom they have a strong and lasting bond



(Jones Harden, 2004; Rhodes et al. 2006)

Mentorship

A relatively long-term, non-expert relationship between a child and non-parental adult, based on acceptance and support, aiming to foster the child’s potential, where change is a desired but not predetermined goal

(Dallos & Comley-Ross, 2005; Rhodes et al. 2006)

Significant effects

Meta-analysis of 55 studies found significant effects of mentoring programs

- Community-based programs more effective than school-based programs

(DuBois & Silverthorn, 2005)

90

Mentoring in foster care

Survey of 29 programs found mentoring provides a bridge to employment and higher education, helps with transitional problem-solving

(Mech, Pryde & Rycraft, 1995)

91

Common factors for success

- ☑ Frequent contacts
- ☑ Emotional closeness (attunement)
- ☑ Longer duration
- ☑ Structured activities
- ☑ Ongoing training for mentors

(DuBois & Silverthorn, 2005; Gilligan, 1999; Rhodes et al. 2006)

92

Mentors enhance resilience

Sensitive mentoring increased self-esteem and well-being, reduced aggression and opened new relationships beyond care system

- *prevents negative outcomes as youth leave foster care*

(DuBois & Silverthorn, 2005; Gilligan, 1999; Lemon et al. 2006; Legault et al. 2005; Rhodes et al. 1999, 2006; Schofield & Beek, 2005)

93

Reduces violence

“Having someone to count on when needed” softened the impact of trauma and reduced likelihood of youth engaging in violent offenses

(Maschi, 2006)

94

Part D

Conclusions and Recommendations



Medicalized approach to distress and disability pathologizes children's behaviors and ignores the context of their experiences

- “Understanding” rather than “diagnosing” changes the meaning of those behaviors and can lead to more helpful interventions

96

Abuse, neglect and trauma disrupt secure attachment and impair the child's ability to self-regulate

- "Repair" occurs through the formation of secure attachments, rather than by medication



Irritability, impulsivity and aggression appear in criteria for most DSM diagnostic labels used on children

- We are medicating children's negative emotions and immature self-control

98

Growing consensus

Just Say 'No' to Drugs as a First Treatment for Child Problems

(Duncan, Sparks, Murphy, & Miller, 2007)

99

Attempt psychosocial interventions *before* initiating medication

Ample evidence supports their use as effective first-line options for children's behavioral problems, *with no apparent risk of medical harm*

100

Fundamental issues of efficacy and safety of psychotropic medications in children remain unresolved

Therefore, medicating children should be avoided



101

A Critical Curriculum on Psychotropic Medications

Module 8

The End



www.CriticalThinkRx.org

102



COMPLETE CURRICULUM REFERENCES

June 2008

- Abel, D. (2007, February 6). Hull parents arrested in girl's poisoning death. *Boston Globe*. Retrieved from http://www.boston.com/news/local/articles/2007/02/06/hull_parents_arrested_in_girls_poisoning_death?mode=PF
- Abboud, L. (2004, June 18). Drug makers seek to bar "placebo responder" from trials. *Wall Street Journal*, pp. B1, B5.
- Adair, R.F. & Holmgren, L.R. (2005). Do drug samples influence resident prescribing behavior? A randomized trial. *The American Journal of Medicine*, 118, 881-884
- Agaibi, C.E., & Wilson, J.P. (2005). Trauma, PTSD, and resilience: A review of the literature. *Trauma, Violence, & Abuse*, 6(3), 195-216.
- Agnell, M. (2000). Is academic medicine for sale? *New England Journal of Medicine*, 342, 1516-1518.
- Allen, S. (2007, June 17). Backlash on bipolar diagnoses in children: MGH psychiatrist's work stirs debate. *Boston Globe*. Retrieved from http://www.boston.com/yourlife/health/children/articles/2007/06/17/backlash_on_bipolar_diagnoses_in_children?mode=PF
- Aman, M., De Smedt, G., Derivan, A., Lyons, B., Findling, R., et al. (2002). Double-blind, placebo-controlled study of risperidone for the treatment of disruptive behaviors in children with subaverage intelligence. *American Journal of Psychiatry*, 159, 1337-1346.
- Amaya-Jackson, L., & DeRosa, R.R. (2007). Treatment considerations for clinicians in applying evidence-based practice to complex presentations in child trauma. *Journal of Traumatic Stress*, 20(4), 379-390.
- America the Pharmaceutical (2005). *Stanford Medicine Magazine*, Summer 2005. Retrieved from <http://mednews.stanford.edu/stanmed/2005summer/drugs-main.html>
- American Counseling Association. (2005). *Code of ethics for the American Counseling Association*. Alexandria, VA: Author. Retrieved from <http://www.counseling.org/Resources/CodeOfEthics/TP/Home/CT2.aspx>
- American Mental Health Counselors Association. (2000). *Code of ethics of the American Mental Health Counselors Association, 2000 Revision*. Alexandria, VA: Author. Retrieved from <http://www.amhca.org/code/>
- American Psychological Association. (2002). *Ethical principles of psychologists and code of conduct*. Washington, DC: Author Retrieved from <http://www.apa.org/ethics/code2002.pdf>

- Andreasen, N.C. (2007). DSM and the death of phenomenology in America: An example of unintended consequences. *Schizophrenia Bulletin*, 33(1), 108-112.
- Antonuccio, D. O., & Healy, D. (2008). The researcher's credo. *British Medical Journal*, 336 (7643), 532-534.
- APA reports cites critical gaps in evidence for current treatment of children's behavioral and mental health problems. (2006). Retrieved March 28, 2007 from, http://www.apa.org/releases/children_meds.html
- APA Working Group on Psychoactive Medications for Children and Adolescents. (2006). *Report of the Working Group on Psychoactive Medications for Children and Adolescents. Psychopharmacological, psychosocial, and combined interventions for childhood disorders: Evidence-base, contextual factors, and future directions*. Washington, DC: American Psychological Association. Retrieved on October 23, 2007 from <http://www.apa.org/pi/cyf/childmeds.pdf>
- Appelbaum, K. (2006). Pharmaceutical marketing and the invention of the medical consumer. *PloS Med* 3(4): e189
- Austrian, S.G. (2005). *Mental disorders, medications and clinical social work* (3rd ed.). New York: Columbia University Press.
- Avorn, J. (2007). Paying for drug approvals—Who's using whom? *New England Journal of Medicine*, 356 (17), 1697-1700.
- Avorn, J. (2004). *Powerful medicines: The benefits, risks, and costs of prescription drugs*. New York: Knopf.
- Backer, E. L., Lebsack, J. A., Van Tonder, R. J., & Crabtree, B. F. (2000). The value of pharmaceutical representative visits and medication samples in community-based family practices. *The Journal of Family Practice*, 49(9), 811-816
- Ballentine, C. (no date). *Taste of raspberries, taste of death: The 1937 Elixir Sulfanilamide incident*. Retrieved May 20, 2008 from: www.fda.gov/oc/history/elixir.html
- Barnett, J. E., & Neel, M. L. (2000). Must all psychologists study psychopharmacology? *Professional Psychology: Research and Practice*, 31(6), 619-627.
- Barratt, M. (2003). Organizational support for evidence-based practice within child and family social work: A collaborative study. *Child and Family Social Work*, 8, 143-150.
- Barreda, A. (Reporter) (2007). 4-year-old's death raises questions. *WCVB-TV, Channel 5, Boston*. Retrieved from <http://www.thebostonchannel.com/video/10946927/index.html>
- Barth, R.P., Landsverk, J., Chamberlain, P., Reid, J.B., Rolls, J.A., Hurlburt, M.S., Farmer, E.M. Z., James, S., McCabe, K.M., & Kohl, P.L. (2005). Parent-training programs in child welfare services: Planning for a more evidence-based approach to serving biological parents. *Research on Social Work Practice*, 15(5), 353-371.

- Bassman, R. (2005). Mental illness and the freedom to refuse treatment: Privilege or right. *Professional Psychology: Research and Practice*, 36(5), 488-497.
- Bauer, A. L., Ingersoll, E., & Burns, L. (2004). School counselors and psychotropic medication: Assessing training, experience, and school policy issues. *Professional School Counseling*, 7(3), 202-211.
- Bellonci, C., & Henwood, T. (2006) *Use of psychotropic medications in child welfare: The needs and challenges of informed consent, ordering, and tracking of medication for children in state custody*. Presentation to the Tennessee Department of Children's Services. Retrieved on November 28, 2007 from the National Resource Center for Family-Centered Practice and Permanency Planning website:
<http://www.hunter.cuny.edu/socwork/nrcfcp/downloads/ppt/Psychotropic-Medications.ppt>
- Bentley, K. J., Walsh, J., & Farmer, R. (2005). Referring clients for psychiatric medication: Best practices for social workers. *Best Practices in Mental Health*, 1(1), 59-71.
- Bentley, K. J., Walsh, J., & Farmer, R. L. (2005). Social work roles and activities regarding psychiatric medication: Results of a national survey. *Social Work*, 50(4), 295-303.
- Bentley, K.J. (1993). The right of psychiatric patients to refuse medication: Where should social workers stand? *Social Work*, 38(1), 101-106.
- Bentley, K.J., & Collins, K.S. (2006). Psychopharmacological treatment for child and adolescent mental disorders. In C. Franklin, M.B. Harris, & P. Allen-Meares (eds.), *The school services sourcebook: A guide for school-based professionals* (pp.15-30). New York: Oxford University Press.
- Bentley, K.J., & Walsh, J. (2006). *The social worker and psychotropic medication: Toward effective collaboration with mental health clients, families, and providers* (3rd ed.). Belmont, CA: Thomson Brooks/Cole.
- Berenson, Alex. (2006, January 1). Antidepressants seem to cut suicide risk. *The New York Times*. Retrieved from
<http://www.nytimes.com/2006/01/01/health/01depress.html?ex=1187755200&en=9659d0e0e167bcf9&ei=5070>
- Beutler, L.E., & Malik, M. (2002). *Rethinking the DSM: A psychological perspective*. Washington, DC: American Psychological Association.
- Bezchlibnyk-Butler, K. AZ., & Jeffries, J. J. (2005). *Clinical handbook of psychotropic drugs* (15th rev. ed.). Seattle: Hogrefe.
- Bhatara, V., Feil, M., Hoagwood, K., Vitiello, B., & Zima, B. (2004). National trends in concomitant psychotropic medication with stimulants in pediatric visits: Practice versus knowledge. *Journal of Attention Disorders*, 7(4), 217-226.
- Bilynsky, N. S., & Vernaglia, E. R. (1998). The ethical practice of psychology in a managed-care framework. *Psychotherapy: Theory, Research, Practice, Training*, 35(1), 54-68.

- Birmaher, B., & Axelson, D. (2006). Course and outcome of bipolar spectrum disorder in children and adolescents: A review of the existing literature. *Development and Psychopathology, Special Issue: Developmental approaches to bipolar disorder, 18*(4), 1023-1035.
- Bloomquist, M.L. (1996). *Skills training for children with behavior problems: A parent and therapist guidebook*. New York: The Guilford Press.
- Boggs, S.R., Eyberg, S.M., Edwards, D.L., Rayfield, A., Jacobs, J., Bagner, D., & Hood, K.K. (2004). Outcomes of parent-child interaction therapy: A comparison of treatment completers and study dropouts one to three years later. *Child & Family Behavior Therapy, 26*(4), 2004, 1-22.
- Boston Globe. (2007, February 8). DSS case files. A chronology of involvement by the state Department of Social Services with the family of 4-year-old Rebecca Riley. *Boston Globe*. Retrieved from http://www.boston.com/news/local/massachusetts/articles/2007/02/08/dss_case_file/
- Bowlby, J. (1988). *A secure base: Parent-child attachment and healthy human development*. New York: Basic Books.
- Boyd, E.A., & Bero, L.A. (2000). Assessing faculty financial relationships with industry. *JAMA, 284*, 2209-2214.
- Bradley, S.S. (2003). The psychology of the psychopharmacology triangle: The client, the clinicians, and the medication. *Social Work in Mental Health, 4*(1), 29-50.
- BrandWeek. (2007). *Strategy: J&J updates ploy that worked for cookies, encyclopedias. Company moves door-to-door approach online to sell OTC drugs*. June 18, 2007, Retrieved from http://www.brandweek.com/bw/search/article_display.jsp?vnu_content_id=1003599809
- BrandWeek. (2007). *The tracker: This is your show on drugs: Rx brands injected into action*. <http://www.brandweek.com>
- BrandWeek. (2007). *US Weekly, eat your heart out: These celebrities are all on drugs*. <http://www.brandweek.com>
- Breggin, P. R. (2002). Fluvoxamine as a cause of stimulation, mania, and aggression, with a critical analysis of the FDA approved label. *International Journal of Risk and Safety in Medicine, 14*, 71-86.
- Breland-Noble, A.M., Elbogen, E.B., Farmer, E.M.Z., Dubs, M.S., Wagner, H.R., & Burns, B.J. (2004). Use of psychotropic medications by youths in therapeutic foster care and group homes. *Psychiatric Services, 55*(6), 706-708.
- Brestan, E.V., & Eyberg, S.M. (1998). Effective psychosocial treatments of conduct-disordered children and adolescents: 29 years, 82 studies, and 5,272 kids. *Journal of Clinical Child Psychology, Special Issue: Empirically supported psychosocial interventions for children, 27*(2), 180-189

- Briere, J., & Spinazzola, J. (2005). Phenomenology and psychological assessment of complex posttraumatic states. *Journal of Traumatic Stress, 18*(5), 401-412.
- Bronstein, L. R. (2003). A model for interdisciplinary collaboration. *Social Work, 48*(3), 297-306.
- Buccino, D.L. (2006). Social work's role in psychiatric medication. *Social Work, 6*(1), 188-189.
- Buckley, J. (2004). Pharmaceutical marketing—time for change. *Electronic Journal of Business Ethics and Organization Studies, 9*(2), 4-11. Retrieved from http://ejbo.jyu.fi/pdf/ejbo_vol9_no2_pages_4-11.pdf
- Buelow, G. D., & Chafetz, M. D. (1996). Proposed ethical practice guidelines for clinical pharmacopsychology: Sharpening a new focus in psychology. *Professional Psychology: Research and Practice, 27*(1), 53-58.
- Butler, A.M., & Eyberg, S.M. (2006). Parent-child interaction therapy and ethnic minority children. *Vulnerable Children and Youth Studies, 1*(3), Dec 2006, 246-255.
- Caldwell, M. F., McCormick, D. J., Umstead, D., & Van Rybroek, G. J. (2007). Evidence of treatment progress and therapeutic outcomes among adolescents with psychopathic features. *Criminal Justice and Behavior, 34*(5), 573-587.
- Campbell, E.G., Gruen, R.L., Mountford, J., Miller, L.G., Cleary, P.D., & Blumenthal, D. (2007). A national survey of physician-industry relationships. *New England Journal of Medicine, 356*(17), 1742-1750.
- Campbell, M., Rapaport, J., & Simpson, G. (1999). Antipsychotics in children and adolescents, *Journal of the American Academy of Child and Adolescent Psychiatry, 38*(5), 537-545.
- Caplan, P. (1995). *They say you're crazy: How the world's most powerful psychiatrists decide who's normal*. Boston: Perseus Books.
- Carey, B. (2006, April 20). Study finds a link of drug makers to psychiatrists. *New York Times*. Retrieved from:
- Carey, B. (2006, June 6). Use of antipsychotics by the young rose fivefold. *The New York Times*. Retrieved from <http://travel.nytimes.com/2006/06/06/health/06psych.html?fta=y>
- Carey, B. (2006, March 7). Study details link of drugs and thoughts of suicide. *The New York Times*. Retrieved from <http://www.nytimes.com/2006/03/07/health/07depress.html?ex=1187755200&en=e30e21ac48130a13&ei=5070>
- Carey, B. (2007, February 15). Debate over children and psychiatric drugs. *New York Times*. Retrieved from <http://www.nytimes.com/2007/02/15/us/15bipolar.html?ei=5070&en=a56bb57ee6609130&ex=1183780800&pagewanted=print>
- Carey, B. (2007, September 3). More children treated for bipolar disorder. *The New York Times*. Retrieved from

http://www.nytimes.com/2007/09/04/health/04psych.html?_r=2&adxnml=1&oref=slogin&adxnmlx=1188923032-vieIaZfSAYw2C//wX2LPCA&oref=slogin

- Carey, B. (2007, September 4). Bipolar illness soars as diagnosis for the young. *The New York Times*. Retrieved from http://www.nytimes.com/2007/09/04/health/04psych.html?_r=1&adxnml=1&oref=slogin&adxnmlx=1188923032-vieIaZfSAYw2C//wX2LPCA
- Carey, B., & Harris, G. (2006, May 12). Antidepressant may carry suicide risk. *The New York Times*. Retrieved from <http://www.nytimes.com/2006/05/12/health/12depress.html?ex=1187755200&en=e1853b257ebc9120&ei=5070>
- Carlat Psychiatry Report (2007). The latest, greatest treatments for PTSD [online]. Retrieved from <http://www.thecarlatreport.com/index.asp?page=wp530200711859>
- Carlson, J. S., Thaler, C. L., & Hirsch, A. J. (2006). Psychotropic medication consultation in schools: An ethical and legal dilemma for school psychologists. *Journal of Applied School Psychology, 22*(1), 29-41.
- Caspe, M., & Lopez, M.A. (2006). Lessons from family-strengthening interventions: Learning from evidence-based practice. *Harvard Family Research Project* [Report]. Retrieved from <http://www.hfrp.org>
- CBS Evening News (2006, 18 October). *Are Drugs Being Misused On Foster Kids?* Retrieved from <http://www.cbsnews.com/stories/2006/10/18/eveningnews/main2104249.shtml>
- CBS News (2004, October 15). FDA orders antidepressant warnings. Retrieved from <http://www.cbsnews.com/stories/2004/08/04/health/main634089.shtml>
- Center of Drug Evaluation and Research (n.d.). Retrieved May 24, 2007, from http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.Search_Drug_Name
- Cepeda, C. (2007). *Psychotic symptoms in children and adolescents: Assessment, differential diagnosis, and treatment*. New York, NY, US: Routledge.
- Charney, D. S., Barlow, D. H., Botteron, K., Cohen, J. D., Goldman, D., Gur, R. E., Lin, K-M., Lopez, J. F., Meador-Woodruff, J. H., Moldin, S. O., Nestler, E. J., Watson, S. J., & Zalcman, S. J. (2002). Neuroscience research agenda to guide development of a pathophysiologically based classification system. In D. J. Kupfer, M. B. First, & D. A. Regier (eds.), *A research agenda for DSM-V* (pp. 31-84). Washington, DC: American Psychiatric Association.
- Chafetz, M. D., & Buelow, G. (1994). A training model for psychologists with prescription privileges: Clinical pharmacopsychologists. *Professional Psychology: Research and Practice, 25*(2), 149-153.
- Chang, K., Saxena, K., & Howe, M. (2006). An open-label study of lamotrigine adjunct or monotherapy for the treatment of adolescents with bipolar

- depression. *Journal of the American Academy of Child and Adolescent Psychiatry*, 45(3), 298-304.
- Chen, H., Deshpande, A., Jiang, R., & Martin, B. (2005). An epidemiological investigation of off-label anticonvulsant drug use in the Georgia Medicaid population. *Pharmacoepidemiology and Drug Safety*, 14, 629-638.
- Chew, L. D., O'Young, T. S., Hazlet, T. K., Bradley, K. A., Maynard, C., & Lessler, D. S. (2000). A physician survey of the effect of drug sample availability on physicians' behavior. *Journal of General Internal Medicine*, 15(7), 478-483.
- Chong, S., Remington, G., & Bezchlibnyk-Butler, K. (2000). Effect of clozapine on polypharmacy. *Psychiatric Services*, 51(2), 250-252.
- Chorpita, B.F., Becker, K.D., & Daleiden, E.L. (2007). Understanding the common elements of evidence-based practice: misconceptions and clinical examples. *Journal of the American Academy of Child and Adolescent Psychiatry*, 46(5), 647-652.
- Choudry, N.K., Stelfox, H.T., & Allan, D. (2002). Relationship between authors of clinical practice guidelines and the pharmaceutical industry. *JAMA*, 287, 612-617.
- Chronis, A.M., Chacko, A., Fabiano, G.A., Wymbs, B.T., & Pelham, W.E. (2004). Enhancements to the behavioral parent training paradigm for families of children with ADHD: Review and future directions. *Clinical Child and Family Psychology Review*, 7(1), 1-27.
- Chronis, A.M., Jones, H.A., & Raggi, V.L. (2006). Evidence-based psychosocial treatments for children and adolescents with attention-deficit/hyperactivity disorder. *Clinical Psychology Review*, 26(4), 486-502.
- Cohen, D. (2002). Research on the drug treatment of schizophrenia: A critical appraisal and implications for social work education. *Journal of Social Work Education*, 38(2), 1-24.
- Cohen, D. (2007). Helping individuals withdraw from psychiatric drugs. *Journal of College Student Psychotherapy*, 21(3/4), 199-224.
- Cohen, D., & Jacobs, D. (2000). A model consent form for psychiatric drug treatment. *Journal of Humanistic Psychology*, 40(1), 59-64.
- Cohen, D., & Jacobs, D. (2007). Randomized controlled trials of antidepressants: Clinically and scientifically irrelevant. *Debates in Neuroscience*, 1, 44-54.
- Cohen, J.A., Berliner, L., & March, J.S. (2000). Guidelines for treatment of PTSD: Treatment of children and adolescents. *Journal of Traumatic Stress*, 13, 566-568.
- Cohen, J.A., Mannarino, A.P., & Deblinger, E. (2006). *Treating trauma and traumatic grief in children and adolescents*. New York: The Guilford Press.
- Cohen, J.A., Mannarino, A.P., & Rogal, S.S. (2001). Treatment practices for childhood PTSD. *Child Abuse and Neglect*, 25, 123-125.
- Coleman, D. L., Kazdin, A. E., Miller, L. A., Morrow, J. S., & Udelsman, R. (2006). Guidelines for interactions between clinical faculty and the

- pharmaceutical industry: One medical school's approach. *Academic Medicine*, 81(2), 154-160.
- Conrad, P. (1992). Medicalization and social control. *Annual Review of Sociology*, 18, 209-232.
- Conrad, P., & Leiter, V. (2004). Medicalization, markets and consumers. *Journal of Health and Social Behavior*, 45 Suppl, 158-176.
- Cook, A., Spinazzola, J., Ford, J., Lanktree, C., Blaustein, M., Cloitre, M., DeRosa, R., et al. (2005). Complex trauma in children and adolescents. *Psychiatric Annals*, 33(5), 390-398.
- Cooper, W.O., Arbogast, P.G., Ding, H., Hickson, G.B., Fuchs, D.C., and Ray, W. (2006). Trends in prescribing of antipsychotic medications for U.S. children. *Ambulatory Pediatrics*, 6, 79-83.
- Cooper, W.O., Hickson, G.B., Fuchs, C., Arbogast, P.G., & Ray, W.A. (2004). New users of antipsychotic medications among children enrolled in TennCare. *Archives of Pediatrics & Adolescent Medicine*, 158: 753-759.
- Corcoran, K., Gorin, S., & Moniz, C. (2005). Managed care and mental health. In S.A. Kirk (Ed.). *Mental disorders in the social environment: Critical perspectives* (pp. 430-442). New York: Columbia University Press.
- Correll, C. U., Lencz, T., Smith, C. W., Auther, A. M., Nakayama, E. Y., & Hovey, L. et al. (2005). Prospective study of adolescents with subsyndromal psychosis: Characteristics and outcome. *Journal of Child and Adolescent Psychopharmacology*, 15(3), 418-433.
- Correll, C., & Carlson, H. (2006). Endocrine and metabolic adverse effects of psychotropic medications in children and adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 45(7), 771-791.
- Costin, J., & Chambers, S.M. (2007). Parent management training as a treatment for children with oppositional defiant disorder referred to a mental health clinic. *Clinical Child Psychology and Psychiatry*, 12(4), 511-524.
- Courtois, C. A. (2004). Complex trauma, complex reactions: Assessment and treatment. *Psychotherapy: Theory, Research, Practice, Training. Special Issue: The Psychological Impact of Trauma: Theory, Research, Assessment, and Intervention*, 41(4), 412-425.
- Coyle, J. T. (2000). Psychotropic drug use in very young children. *JAMA*, 283(8), 1059-1060.
- Cramer, M. (2007, February 8). DSS dropped inquiry before girl, 4, was found dead; got assurances on medications. *Boston Globe*. Retrieved from http://www.boston.com/news/local/articles/2007/02/08/dss_dropped_inquiry_before_girl_4_was_found_dead?mode=PF
- Curtis, N.M., Ronan, K.R., & Borduin, C.M. (2004). Multisystemic treatment: A meta-analysis of outcome studies. *Journal of Family Psychology*, 18(3), 411-419.

- Dallos, R. & Comley-Ross, P. (2005). Young people's experience of mentoring: Building trust and attachments. *Clinical Child Psychology and Psychiatry*, 10(3), 369-383.
- Daly, R. (2006). Psychiatrists proactive in scope-of-practice battles. *Psychiatric News*, 41(5), 17. Retrieved from <http://pn.psychiatryonline.org/cgi/content/full/41/5/17>
- Dana, J., & Loewenstein, G. (2003). Doctors and drug companies: A social science perspective on gifts to physicians from industry. *JAMA*, 290(2), pp. 252-255.
- Danielson, C. K., Feeny, N. C., Findling, R. L., & Youngstrom, E. A. (2004). Psychosocial treatment of bipolar disorders in adolescents: A proposed cognitive-behavioral intervention. *Cognitive and Behavioral Practice*, 11(3), 283-297.
- Davidson, R.A. (1986). Source of funding and outcome of clinical trials. *Journal of General Medicine*, 1, 155-158.
- DeAngelis, T. (2007). A new diagnosis for childhood trauma? Some push for a new DSM category for children who undergo multiple, complex traumas. *Monitor on Psychology*, 38(3). Retrieved from <http://www.apa.org/monitor/mar07/diagnosis.html>
- DeAngelis, T. (2008). PTSD treatments grow in evidence, effectiveness. *Monitor on Psychology*, 39(1), 40-43.
- DeGangi, G. (2000). *Pediatric disorders of regulation in affect and behavior: A therapist's guide to assessment and treatment*. San Diego, CA: Academic Press.
- Delbello, M. P., Hanseman, D., Adler, C. M., Fleck, D. E., & Strakowski, S. M. (2007). Twelve-month outcome of adolescents with bipolar disorder following first hospitalization for a manic or mixed episode. *American Journal of Psychiatry*, 164(4), 582-590.
- Demb, H., & Nguyen, K. (1999). Movement disorders in children with developmental disabilities taking risperidone. *Journal of the American Academy of Child and Adolescent Psychiatry*, 38(1), 5-6.
- Department of Health and Human Services. (2004). *Warning letter to Janssen Pharmaceutica, Inc.*
- Diamond, G., & Josephson, A. (2005). Family-based treatment research: A 10-year update. *Journal of the American Academy of Child & Adolescent Psychiatry*, 44(9), 872-887.
- Diamond, G., Siqueland, L., & Diamond, G. M. (2003). Attachment-based family therapy for depressed adolescents: Programmatic treatment development. *Clinical Child and Family Psychology Review*, 6(2), 107-127.
- Diller, L. (2007, June 19). Misguided standards of care. *Boston Globe*, Editorial Opinion. Retrieved from http://www.boston.com/news/globe/editorial_opinion/oped/articles/2007/06/19/misguided_standards_of_care/

- Donohue, J.M., & Bernd, E.R. (2004). Effects of direct to consumer advertising on medication choice: The case of antidepressants. *Journal of Public Policy & Marketing*, 23(2), 115-127
- Donovan, D.M., & McIntyre, D. (1990). *Healing the hurt child: A developmental-contextual approach*. New York: W.W. Norton & Company.
- dosReis, S., Zito, J. M., Safer, D. J., & Soeken, K. L. (2001). Mental health services for youths in foster care and disabled youths. *American Journal of Public Health*, 91(7), 1094-1099.
- dosReis, S., Zito, J. M., Safer, D. J., Gardner, J. F., Puccia, K. B., & Owens, P. L. (2005). Multiple psychotropic medication use for youths: A two-state comparison. *Journal of Child and Adolescent Psychopharmacology, Special Issue on Psychopharmacoepidemiology*, 15(1), 68-77.
- Dozier, M., Peloso, E., Lindhiem, O., Gordon, M.K., Manni, M., Sepulveda, S., & Ackerman, J. (2006). Developing evidence-based interventions for foster children: An example of a randomized clinical trial with infants and toddlers. *Journal of Social Issues*, 62(4), 767-785.
- Drugs by the numbers (2005, Summer). *Stanford Medicine Magazine*. Retrieved from <http://mednews.stanford.edu/stanmed/2005summer/drugs-numbers.html>
- DuBois, D. L., & Silverthorn, N. (2005). Characteristics of natural mentoring relationships and adolescent adjustment: Evidence from a national study. *Journal of Primary Prevention, Special Issue: Mentoring with Children and Youth*, 26(2), 69-92.
- DuBois, D.L., Holloway, B.E., Valentine, J.C., & Cooper, H. (2002). Effectiveness of mentoring programs for youth: A meta-analytic review. *American Journal of Community Psychology*, 30(2), 157-196.
- Duggan (2005). Do new prescription drugs pay for themselves? The case of second-generation antipsychotics. *Journal of Health Economics*, 24, 1-31.
- Duncan, B.L., Sparks, J.A., Murphy, J.J., Miller, S.D. (2007). Just say 'no' to drugs as a first treatment for child problems. *Psychotherapy in Australia*, 13(4), 32-40.
- Dunivin, D. L., & Southwell, G. D. (2000). Psychopharmacology training in psychology internships: A brief curriculum. *Professional Psychology: Research and Practice*, 31(6), 610-614.
- Dziegielewski, S. F. (1998). Psychopharmacology and social work practice: Introduction. *Research on Social Work Practice. Special Issue: Psychopharmacology and social work practice*, 8(4), 371-383.
- Dziegielewski, S. F., & Leon, A. M. (1998). Psychopharmacology knowledge and use with social work professionals: A continuing education evaluation. *Professional Development: The International Journal of Continuing Social Work Education*, 1(3), 31-40.
- Elias, M. (2006, May 2). A rush to overprescribe? *USA Today*, retrieved from http://www.usatoday.com/news/health/2006-05-01-kids-overprescribe_x.htm

- Elias, M. (2006, May 2). Adult antipsychotics worsen troubles. *USA Today*, retrieved from http://www.usatoday.com/news/health/2006-05-01-adult-antipsychotics-kids_x.htm
- Elias, M. (2006, May 2). New antipsychotic drugs carry risks for children. *USA Today*, retrieved from http://www.usatoday.com/news/health/2006-05-01-atypical-drugs_x.htm?loc=interstitialskip
- Elkins, D.N. (2007). Empirically supported treatments: The deconstruction of a myth. *Journal of Humanistic Psychology*, 47(4), 474-500.
- Elliott, C. (2004). Pharma goes to the laundry: Public relations and the business of medical education. *Hastings Center Report*, 34(5), 18-23.
- Elliott, C. (2006). The drug pushers. *Atlantic Monthly (April)*, 2-13.
- El-Mallakh, R. S. (1994). Teaching psychopharmacology to psychology interns. *Psychological Reports*, 74(2), 674.
- Emslie, G. J., Heiligenstein, J. H., Wagner, K. D. et al. (2002). Fluoxetine for acute treatment of depression in children and adolescents: A placebo controlled, randomized clinical trial. *Journal of the American Academy of Child and Adolescent Psychiatry*, 41, 1205-1215.
- Families USA. (2005, January). *Big Pharma behaving badly: A survey of selected class action lawsuits against drug companies* (4th ed.). Prepared for Families USA by Miller Faucher and Cafferty, LLP. Retrieved from http://www.familiesusa.org/assets/pdfs/Rx_Lawsuits_Survey_4th_edition_pmdabc2.pdf
- Farley, R. (2007, July 29). The atypical dilemma: Skyrocketing numbers of children are prescribed powerful antipsychotic drugs. Is it safe? Nobody knows. *St. Petersburg Times*. Retrieved from http://www.sptimes.com/2007/07/29/Worldandnation/The_atypical_dilemm.shtml
- Farley, R. (2008, April 12). Drug research: To test or to tout? *St. Petersburg Times*. Retrieved from <http://www.tampabay.com/news/health/article454391.ece>
- Farley, S.E., Adams, J.S., Lutton, M.E., & Scoville, C. (2005). What are effective treatments for oppositional and defiant behaviors in preadolescents? *The Journal of Family Practice*, 54(2), 162-165.
- Farmer, E.M.Z., Dorsey, S., & Mustillo, S.A. (2004). Intensive home and community interventions. *Child and Adolescent Psychiatric Clinics of North America. Special Issue: Evidence-Based Practice, Part I: Research Update*, 13(4), 857-884.
- Farmer, R. L., Walsh, J., & Bentley, K. J. (2006). Advancing social work curriculum in psychopharmacology and medication management. *Journal of Social Work Education*, 42(2), 211-229.
- Faust, J., & Katchen, L. B. (2004). Treatment of children with complicated posttraumatic stress reactions. *Psychotherapy: Theory, Research, Practice, Training*

(Special Issue: *The Psychological Impact of Trauma: Theory, Research, Assessment, and Intervention*) 41(4), 426-437.

- Findling, R., Gracious, B., & Seman, C. (2004). Elevated thyrotropin in bipolar youths prescribed both lithium and divalproex sodium. *Journal of the American Academy of Child and Adolescent Psychiatry*, 43(2), 215-220.
- Findling, R., McNamara, N., Branicky, L., Schluchter, M., Lemon, E., & Blumer, J. (2000). A double-blind pilot study of risperidone in the treatment of conduct disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 39(4), 509-516.
- Findling, R., McNamara, N., Youngstrom, E., Stansbrey, R., Gracious, B., et al. (2005). Double-blind 18-month trial of lithium versus divalproex maintenance treatment in pediatric bipolar disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 44(5), 409-417.
- Findling, R.L, Boorady, R.J., & Sporn, A.L. (2007). *The treatment of bipolar disorder and schizophrenia in children and adolescents*. Medscape CME. Retrieved from <http://www.medscape.com/viewarticle/563314>
- Fisher, S., & Greenberg, R. P. (1993). How sound is the double-blind design for evaluating psychotropic drugs? *Journal of Nervous and Mental Disease*, 181, 345-350.
- Floersch, J. (2003). The subjective experience of youth psychotropic treatment. *Social Work in Mental Health*, 1(4), 51-69.
- Florida Center for Medicaid and the Uninsured (2005). A dilemma for state Medicaid programs: A look at the methods of addressing the rising pharmaceutical expenditures for treatments of mental health disorders. *Health Coverage Updates, Issue Brief*, January, 2005. Retrieved from http://fcmu.phhp.ufl.edu/pdf/1_issue_briefs_and_fact_sheets/fcmu_issue_briefs/IssueBrief03-StateMedicaidProgramsDilemma.pdf
- Florida Statewide Advocacy Council (2003) *Red Item Report: Psychotropic Drug Use in Foster Care*. Retrieved from http://www.floridasac.org/sacweb/documents/red_psychotropic.doc
- Florida Statutes 381.026 Florida's Patient Bill of Rights and Responsibilities. Retrieved from <http://www.doh.state.fl.us/MQA/Profiling/billofrights.htm>
- Food and Drug Administration, Center for Drug Evaluation and Research. (1998). *Guidance for industry: Providing clinical evidence of effectiveness for human drug and biological products*. 6, 9. Available at: <http://www.fda.gov/cder/guidance/1397fnl.pdf>
- Ford, J. D., Courtois, C. A., Steele, K., van der Hart, O., & Nijenhuis, E. R. S. (2005). Treatment of complex posttraumatic self-dysregulation. *Journal of Traumatic Stress*, 18(5), 437-447.
- Fostering Positive Outcomes. (2006). Retrieved from http://www.mentoring.org/program_staff/research_corner/fostering_positive_outcomes.php

- Fox, R. E., Schwelitz, F. D., & Barclay, A. G. (1992). A proposed curriculum for psychopharmacology training for professional psychologists. *Professional Psychology: Research and Practice*, 23(3), 216-219.
- Frazier, J., Meyer, M., Bierderman, J., Wozniak, J., Wilens, T., et al. (1999). Risperidone treatment for juvenile bipolar disorder: A retrospective chart review, *Journal of the American Academy of Child and Adolescent Psychiatry*, 38(8), 960-965.
- Freimuth, M. (1996). Combining psychotherapy and psychopharmacology: With or without prescription privileges. *Psychotherapy: Theory, Research, Practice, Training*, 33(3), 474-478.
- Fristad, M. A. (2006). Psychoeducational treatment for school-aged children with bipolar disorder. *Development and Psychopathology, Special Issue: Developmental Approaches to Bipolar Disorder*, 18(4), 1289-1306.
- Fristad, M. A., Goldberg-Arnold, J. S., & Gavazzi, S. M. (2003). Multi-family psychoeducation groups in the treatment of children with mood disorders. *Journal of Marital & Family Therapy*, 29(4), 491-504.
- Frost, N., Robinson, M., & Anning, A. (2005). Social workers in multidisciplinary teams: Issues and dilemmas for professional practice. *Child & Family Social Work*, 10(3), 187-196. Retrieved July 12, 2007, from PsycINFO database.
- Fugh-Berman, A. (2008). Prescription tracking and public health. *Journal of General Internal Medicine*, DOI: 10.1007/s11606-008-0630-0
- Fugh-Berman, A., & Ahari, S. (2007). Following the script: How drug reps make friends and influence doctors. *PloS Medicine*, 4(4): e150.
- Fugh-Berman, A., Alladin, K., & Chow, J. (2006). Advertising in medical journals: Should current practices change? *PLoS Medicine*, 3(6), e130.
- Gambrill, E. (2006). Evidence-based practice and policy: Choices ahead. *Research on Social Work Practice*, 16(3), 338-357.
- Gambrill, E. D. (2002). Encouraging transparency. *Journal of Social Work Education*, 38(2), 211-215.
- Gardner, F., Shaw, D.S., Dishion, T.J., Burton, J., & Suplee, L. (2007). Randomized prevention trial for early conduct problems: Effects on proactive parenting and links to toddler disruptive behavior. *Journal of Family Psychology*, 21(3), 398-406.
- Geddes, J., Freemantle, N., Harrison, P., & Bebbington, P. (2000). Atypical antipsychotics in the treatment of schizophrenia: Systematic overview and meta-regression analysis. *British Medical Journal*, 32, 1371-1376.
- Gellad, Z.F., & Lyles, K.W. (2007). Direct-to-consumer advertising of pharmaceuticals. *The American Journal of Medicine*, 120(6), 475-480.
- Geller, B., Tillman, R., Craney, J. L., & Bolhofner, K. (2004). Four-year prospective outcome and natural history of mania in children with a prepubertal and early

- adolescent bipolar disorder phenotype. *Archives of General Psychiatry*, 61(5), 459-467.
- Gilligan, R. (1999). Enhancing the resilience of children and young people in public care by mentoring their talents and interests. *Child & Family Social Work*, 4(3), 187-196.
- Ginsberg, T. (2006, May 28). Donations tie drug firms and nonprofits. *Philadelphia Inquirer*, Retrieved from <http://www.philly.com>
- Gleason, M.M., Egger, H.L., Emslie, G.J., Greenhill, L.L., Kowatch, R.A., Lieberman, A.F., et al. (2007). Psychopharmacological treatment for very young children: contexts and guidelines. *Journal of the American Academy of Child & Adolescent Psychiatry*, 46(12), 1532-1572.
- Gonzalez-Heydrich, J., Raches, D., Wilens, T., Leichtner, A., & Mezzacappa, E. (2003). Retrospective study of hepatic enzyme elevations in children treated with olanzapine, divalproex, and their combination. *Journal of the American Academy of Child and Adolescent Psychiatry*, 42(10), 1227-1233.
- Goode, E. (2003, January, 14). Study finds jump in children taking psychiatric drugs. *The New York Times*. Retrieved from <http://query.nytimes.com/gst/fullpage.html?res=9906EEDE1231F937A25752C0A9659C8B63&n=Top/Reference/Times%20Topics/People/G/Goode,%20Erica>
- Goodman, R.L. (2007). Medical education and the pharmaceutical industry. *Perspectives in Biology and Medicine*, 50(1), 32-39.
- Goodwin, R., Gould, M.S., Blanco, C., & Olfson, M. (2001). Prescription of psychotropic medications to youths in office-based practice. *Psychiatric Services*, 52(8), 1081-1087.
- Goozner, M. (2004). *The \$800 million pill: The truth about the cost of new drugs*. Berkeley, CA: University of California Press.
- Gracious, B., Llana, M., & Barton, D. (1999). Lithium and geographic tongue. *Journal of the American Academy of Child and Adolescent Psychiatry*, 38(9), 1069-1070.
- Gray, M., & McDonald, C. (2006). Pursuing good practice? The limits of evidence-based practice. *Journal of Social Work*, 6(1), 7-20.
- Green, D.L., Hawkins, W., & Hawkins, M. (2005). Medication of children and youth in foster care. *Journal of Social Work in Disability & Rehabilitation*, 4(1/2), 43-55.
- Greene, J.A. (2004). Attention to 'details': Etiquette and the pharmaceutical salesman in postwar American. *Social Studies of Science*, 34(2), 271-292
- Greenhill, L. L., Vitiello, B., Abikoff, H., Levine, J., March, J. S. et al. (2003). Developing methodologies for monitoring long-term safety of psychotropic medications in children: Report on the NIMH conference, September 25, 2000. *Journal of the American Academy of Child and Adolescent Psychiatry*, 42(6), 651-655.

- Greenwald, R. (2000). A trauma-focused individual therapy approach for adolescents with conduct disorder. *International Journal of Offender Therapy and Comparative Criminology*, 44(2), 146-163.
- Gunnar, M.R., Fisher, P.A., et al. (2006). Bringing basic research on early experience and stress neurobiology to bear on preventive interventions for neglected and maltreated children. *Development and Psychopathology*, 18, 651-677.
- Hanson, K.M., & Sheridan, K. (1997). Ethics and changing mental health care: concerns and recommendations for practice. *Journal of Clinical Psychology in Medical Settings*, 4(2), 231-242.
- Harris, G. (2004, December 6). At FDA, strong drug ties and less monitoring. *The New York Times*.
- Harris, G. (2006, August 22). FDA strengthens warnings on stimulants. *The New York Times*, Retrieved from <http://www.nytimes.com/2006/08/22/health/policy/22fda.html?ex=1313899200&en=30ee5026f29179e3&ei=5088&partner=rssnyt&emc=rss>
- Harris, G. (2006, November 22). Proof is scant on psychiatric drug mix for young. *The New York Times*. Retrieved from <http://www.nytimes.com/2006/11/23/health/23kids.html?scp=1&sq=psychiatric+medicines+and+children&st=nyt>
- Harris, G. (2007, June 27). Psychiatrists top list in drug maker gifts. *New York Times*, Retrieved from <http://www.nytimes.com/2007/06/27/health/psychology/27doctors.html>
- Harris, G. (2007, June 27). Psychiatrists top list in drug maker gifts. *New York Times*. Retrieved from <http://www.nytimes.com/2007/06/27/health/psychology/27doctors.html?ref=health>
- Harris, G. (2007, June 28). Senators to push for registry of drug makers' gifts to doctors. *New York Times*. Retrieved from
- Harris, G., Carey, B., & Roberts, J. (2007, May 10). Psychiatrists, children, and drug industry's role. *New York Times*, Retrieved from <http://www.nytimes.com/2007/05/10/health/10psyche.html?ex=1183608000&en=60e51ac0bcd516cf&ei=5070>
- Haugaard, J.J. (2004). Recognizing and treating rare behavioral and emotional disorders in children and adolescents who have been maltreated: Schizophrenia. *Child Maltreatment*, 9(2), 161-168.
- Haw, C., & Stubbs, J. (2005). A survey of off-label prescribing for inpatients with mild intellectual disability and mental illness, *Journal of Intellectual Disability Research*, 49, 858-864.
- Hawthorne, F. (2005). *Inside the FDA: The business and politics behind the drugs we take and the food we eat*. New York: Wiley.
- Healy, D., & Farquhar, G. (1998). Immediate effects of droperidol. *Human*

Psychopharmacology, 13, 113-120.

- Heiby, E. M. (2002). It is time for a moratorium on legislation enabling prescription privileges for psychologists. *Clinical Psychology: Science and Practice*, 9(3), 256-258.
- Heiby, E. M. (2002). Prescription privileges for psychologists: Can differing views be reconciled? *Journal of Clinical Psychology*, 58(6), 589-597.
- Hembree-Kigin, T., & McNeill, C.B. (1995). *Parent-child interaction therapy*. New York: Plenum.
- Heres, S., Davis, J., Maino, K. et al. (2006). Why olanzapine beats risperidone, risperidone beats quetiapine, and quetiapine beats olanzapine: An exploratory analysis of head-to-head comparison studies of second-generation antipsychotics. *American Journal of Psychiatry*, 163,185–194.
- Hibbs, E.D., & Jensen, P.S. (2005). *Psychosocial treatments for child and adolescent disorders: Empirically based strategies for clinical practice* (2nd ed.). Washington, DC: American Psychological Association.
- High rates of off-label prescriptions for antidepressant, anticonvulsant and antipsychotic drugs.* (2006). Retrieved May 24, 2007, from <http://www.news-medical.net/?id=19083>
- Hoagwood, K., Burns, B.J., Kiser, L., Ringeisen, H., Schoenwald, S.K. (2001). Evidence-based practice in child and adolescent mental health services. *Psychiatric Services*, 52(9), 1179-1189.
- Hollon, M.F. (1999). Direct-to-consumer marketing of prescription drugs: Creating consumer demand. *JAMA*, 281, 382-384.
- Hubble, M.A., Duncan, B.L., & Miller, S.D. (Eds.). (1999). *The heart and soul of change: What works in therapy*. Washington, DC: American Psychological Association.
- Hundley, K. (2007, July 1). Companies run trials of drugs they invest in: Critics say a conflict of interest couldn't be more clear, but the FDA doesn't think so. *St. Petersburg Times*. Retrieved from http://www.sptimes.com/2007/07/01/news_pf/Worldandnation/Companies_run_trials_shtml
- Ingersoll, R. E. (2000). Teaching a psychopharmacology course to counselors: Justification, structure and methods. *Counselor Education and Supervision*, 40(1), 58-69.
- Ingersoll, R. E. (2002). An integral approach for teaching and practicing diagnosis. *The Journal of Transpersonal Psychology*, 34, 115-127.
- Ingersoll, R.E., Bauer, A., & Burns, L. (2004). Children and psychotropic medication: What role should advocacy counseling play? *Journal of Counseling & Development*, 82, 337-343.
- Ingersoll, R. E., & Rak, C. F. (2006). *Psychopharmacology for helping professionals: An integral approach*. Pacific Grove, CA: Brooks/Cole.

- Irwin, M. (2004). Treatment of schizophrenia without neuroleptics: psychosocial interventions versus neuroleptic treatment. *Ethical Human Psychology and Psychiatry*, 6(2), 99-110.
- Jacobs, B. (2006). *Legal strategies to challenge chemical restraint of children in foster care: A resource for child advocates in Florida*. Tampa, FL: Advocacy Center for Persons with Disabilities, Inc.
- Jacobs, D.H., & Cohen, D. (1999). What is really known about psychological alterations produced by psychiatric drugs? *International Journal of Risk and Safety in Medicine*, 12(1), 37-47.
- Jacobs, D.H., & Cohen, D. (2004). Hidden in plain sight: DSM-IV's rejection of the categorical approach to diagnosis. *Review of Existential Psychology and Psychiatry*, 26(2-3), 81-96.
- Jefferson, A., Markowitz, J., & Brewerton, T. (1998). Atypical antipsychotics [letter]. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37(12), 1243-1244.
- Jensen, P.S., Bhatara, V.S., Vitiello, B., Hoagwood, K., Feil, M., and Burke, L.B. (1999). Psychoactive medication prescribing practices for U.S. children: Gaps between research and clinical practice. *Journal of the Academy of Child and Adolescent Psychiatry*, 38(5), 557-565.
- Jensen, P.S., Youngstrom, E.A., Steiner, H., Findling, R.L., Meyer, R.E., Malone, R.P., Carlson, G.A. et al. (2007). Consensus report on impulsive aggression as a symptom across all diagnostic categories in child psychiatry: implications for medication studies. *Journal of the American Academy of Child and Adolescent Psychiatry*, 46(3), 309-322.
- Johnson, L.A. (2006, August 9). Study cites need for therapy. *The Miami Herald*, Page 7A.
- Jones Harden, B. (2004). Safety and stability for foster children: A developmental perspective. *The Future of Children*, 14(1), 30-47.
- Jordan, C., & Franklin, C. (2003). *Clinical assessment for social workers: Quantitative and qualitative methods* (2nd ed.). Chicago, IL: Lyceum Books, Inc.
- Jordan, S., Knight, J., & Pointon, D. (2004). Monitoring adverse drug reactions: Scales, profiles, and checklists. *International Nursing Review*, 51(4), 208-221.
- Jureidini, J. N., Doecke, C. J., Mansfield, P. R., Haby, M. M., Menkes, D. B., & Tonkin, A. L. (2004). Efficacy and safety of antidepressants for children and adolescents. *British Medical Journal*, 328, 879-883.
- Kalachnik, J. E. (1999). Measuring side effects of psychopharmacologic medication in individuals with mental retardation and developmental disabilities. *Mental Retardation And Developmental Disabilities Research Reviews*, 5(4), 348-359.
- Kane, J. (2006). Tardive dyskinesia circa 2006. *American Journal of Psychiatry*, 163(8), 1316-1318.

- Kassirer, J.P. (2006, February 13). How drug lobbyists influence doctors. *Boston Globe*. Retrieved from http://www.boston.com/news/globe/editorial_opinion/oped/articles
- Kazdin, A. E. (2005). *Parent management training: Treatment for oppositional, aggressive, and antisocial behavior in children and adolescents*. New York: Oxford University Press.
- Kazdin, A.E. (2000). Treatments for aggressive and antisocial children. *Child and Adolescent Psychiatric Clinics of North American*, 9(4), 841-858.
- Kazdin, A.E. (2000b). *Psychotherapy for children and adolescents: Directions for research and practice*. New York: Oxford University Press.
- Kazdin, A.E., & Weisz, J.R. (Eds.). (2003). *Evidence-based psychotherapies for children and adolescents*. New York: The Guilford Press.
- Keating, L. M., Tomishima, M. A., Foster, S., & Alessandri, M. (2002). The effects of a mentoring program on at-risk youth. *Adolescence*, 37(148), 717-734.
- Kenkel, M. B. (2006). Professional psychology: Expanding its discoveries, reach, and impact. *Professional Psychology: Research and Practice*, 37(6), 587-589.
- Kesselheim, A., & Avorn, J. (2007). The role of litigation in defining drug risks. *JAMA*, 297(3), 308-311.
- Kewley, G. (1999). Risperidone in comorbid ADHD and ODD/CD [letter]. *Journal of the American Academy of Child and Adolescent Psychiatry*, 38(11), 1327-1328.
- Kilpatrick, D.G. (2005). A special section on complex trauma and a few thoughts about the need for more rigorous research on treatment efficacy, effectiveness and safety. *Journal of Traumatic Stress*, 18(5), 379-384.
- King, J. H., & Anderson, S. M. (2004). Therapeutic implications of pharmacotherapy: Current trends and ethical issues. *Journal of Counseling & Development*, 82(3), 329-336.
- Kinniburgh, K., Blaustein, M., Spinazzola, J. & van der Kolk, B. (2005). Attachment, self-regulation and competency: A comprehensive framework for intervention with childhood complex trauma. *Psychiatric Annals*, 35(5), 424-430.
- Kirk, S., & Kutchins, H. (1994). The myth of the reliability of DSM. *Journal of Mind and Behavior*, 15(1&2), 71-86.
- Kirk, S. A., & Kutchins, H. (1992). *The selling of DSM: The rhetoric of science in psychiatry*. Chicago: Aldine de Gruyter.
- Koerner, B.I. (2002, July/August). Disorders made to order. *Mother Jones magazine*. Retrieved from <http://www.motherjones.com/news/feature/2002/07/disorders.html>
- Kowalczyk, L. (2007, February 8). Psychiatrist to suspend practice; denies wrongdoing. *Boston Globe*. Retrieved from http://www.boston.com/news/local/massachusetts/articles/2007/02/08/psychiatrist_to_suspend_practice_denies_wrongdoing?mode=PF

- Kowatch, R. A., Fristad, M., Birmaher, B., Wagner, K. D., Findling, R. L., & Hellander, M. et al. (2005). Treatment guidelines for children and adolescents with bipolar disorder: Child psychiatric workgroup on bipolar disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 44(3), 213-235.
- Kowatch, R., Suppes, T., Carmody, R., Bucci, J., Hume, J., et al. (2000). Effect size of lithium, divalproex sodium, and carbamazepine in children and adolescents with bipolar disorder, *Journal of the American Academy of Child and Adolescent Psychiatry*, 39(6), 713-720.
- Kroft, S. (Reporter). (2007, April 1) Under the influence: Drug lobbyists' role in passing bill that keeps drug prices high. In *60 Minutes* (Producer). New York: CBS News. Retrieved from <http://www.cbsnews.com/stories/2007/03/29/60minutes/printable2625305.shtml>
- Kutchins, H., & Kirk, S. A. (1998). *Making us crazy: The psychiatric bible and the creation of mental disorders*. New York: Free Press.
- Lacasse, J. & Gomory, T. (2003). Is graduate social work education promoting a critical approach to mental health practice? *Journal of Social Work Education*, 39(3), 383-408.
- Lakoff, A. (2004). The anxieties of globalization: Antidepressant sales and economic crisis in Argentina. *Social Studies of Science*, 34(2), 247-269.
- Lambert, L. (2007, February 12). When families don't work. *The Patriot Ledger*. Retrieved from <http://www.patriotledger.com/articles/2007/02/12/news/news07.txt>
- Lasser, K.E., Allen, P.D., Woolhandler, S.J., Himmelstein, D.U., Wolfe, S.M., & Bor, D.H. (2002). Timing of new black box warnings and withdrawals for prescription medications. *JAMA*, 287(17), 2215-2220.
- Lee, V., & Hoaken, P.N.S. (2007). Cognition, emotion, and neurobiological development: Mediating the relation between maltreatment and aggression. *Child Maltreatment*, 12(3), 281-298.
- Legault, L., Anawati, M., & Flynn, R. (2005). Factors favoring psychological resilience among foster young people. *Children and Youth Services Review*, 28, 1024-1038.
- Lemon Osterling, K., & Hines, A.M. (2006). Mentoring adolescent foster youth: Promoting resilience during developmental transitions. *Child and Family Social Work*, 11, 242-253.
- Leo, J. (2004). Multiple comparisons in drug efficacy studies: Scientific or marketing principles? *Ethical Human Psychology and Psychiatry*, 6(1), 3-6.
- Leslie, L.K., Weckerly, J., Landsverk, J., Hough, R.L., Hurlburt, M.S., Wood, P.A. (2003). Racial/ethnic differences in the use of psychotropic medication in high-risk children and adolescents. *Journal of the Academy of Child and Adolescent Psychiatry*, 42(1), 1433-1442.

- Levant, R. F., & Shapiro, A. E. (2002). Training psychologists in clinical psychopharmacology. *Journal of Clinical Psychology, 58*(6), 611-615.
- Lieberman, J. A., Stroup, T. S., McEvoy, J. P., Swartz, M., Rosenheck, R. A. et al. (2005). Effectiveness of antipsychotic drugs in patients with chronic schizophrenia. *New England Journal of Medicine, 353*, 1209-1223.
- Lindenmayer, J., Czoher, P., Volavka, J., Citrome, L., Sheitman, B., et al. (2003). Changes in glucose and cholesterol levels in patients with schizophrenia treated with typical or atypical antipsychotics, *American Journal of Psychiatry, 160*(2), 290-296.
- Littrell, J. (2003). Obtaining informed consent when a profession labels itself as providing treatment for mental illness. *Social Work in Mental Health, 1*(4), 107-122.
- Littrell, J., & Ashford, J. B. (1995). Is it proper for psychologists to discuss medications with clients? *Professional Psychology: Research and Practice, 26*(3), 238-244.
- Loder, E., & Biondi, D. (2004). Off-label prescribing of drugs in specialty headache practice, *Headache, 44*, 636-641.
- Long, J. E. J. (2005). Power to prescribe: The debate over prescription privileges for psychologists and the legal issues implicated. *Law & Psychology Review, 29*, 243-260.
- Lyons, J., MacIntyre, J., Lee, M., Carpinello, S., Zuber, M., & Fazio, M. (2004). Psychotropic medications prescribing patterns for children and adolescents in New York's public mental health system, *Community Mental Health Journal, 40*(2), 101-118.
- Maj, M. (2005). The aftermath of the concept of "psychiatric comorbidity." *Psychotherapy & Psychosomatics, 74*, 67-68.
- Malone, R., Sheikh, R., & Zito, J. (1999). Novel antipsychotic medications in the treatment of children and adolescents, *Psychiatric Services, 50*(2), 171-174.
- Marbin-Miller, C. (2005, January 15). 1 in 4 foster kids on risky mind medications. *The Miami Herald*, Page, 1A.
- Marbin-Miller, C. (2005, May 5). New law curbs drugging of kids. *The Miami Herald*, page 1A.
- Marbin-Miller, C. (2006, September 23). No list of kids on mood drugs. *The Miami Herald*, Page 1B.
- March, J.S. (1995). *Anxiety disorders in children and adolescents*. New York: Guilford Press.
- Marquis, A. (2008). *The integral intake: A guide to comprehensive idiographic assessment in integral psychotherapy*. New York: Routledge.
- Martin, A. & Leslie, D. (2003). Trends in psychotropic medication costs for children and adolescents, 1997-2000. *Archive of Pediatrics & Adolescent Medicine, 157*, 997-1004.

- Martin, A., Sherwin, T., Stubbe, D., Van Hoof, T., Scahill, L., & Leslie, D. (2002). Use of multiple psychotropic drugs by Medicaid-insured and privately insured children. *Psychiatric Services, 53*(12), 1508.
- Maschi, T. (2006). Trauma and violent delinquent behavior among males: the moderating role of social support. *Stress, Trauma, and Crisis, 9*, 45-72.
- Mash, E.J., & Barkley, R.A. (Eds.). (2006). *Treatment of childhood disorders* (3rd ed.) New York: The Guilford Press.
- Masters, K., & Melonas, J. (2004). Lamotrigine and informed consent [letter]. *Journal of the American Academy of Child and Adolescent Psychiatry, 43*(2), 130-131.
- Matsui, D., Kwan, C., Steer, E., & Rieder, M.J. (2003). The trials and tribulations of doing drug research in children. *Canadian Medical Association Journal, 169*(10), 1033-1034.
- McCart, M.R., Priester, P.E., Davies, W.H., & Azen, R. (2006). Differential effectiveness of behavioral parent-training and cognitive-behavioral therapy for antisocial youth: A meta-analysis. *Journal of Abnormal Child Psychology, 34*(4), 527-543
- McDonagh, M.S., Peterson, K., & Carson, S. (2006). *Drug class review on atypical antipsychotic drugs*. Center for Evidence-based Policy, Oregon Health & Science University. Update #1 Final Report. Retrieved from: <http://www.ohsu.edu/drugeffectiveness>
- McDonnell, M. G., & Dyck, D. G. (2004). Multiple-family group treatment as an effective intervention for children with psychological disorders. *Clinical Psychology Review, 24*(6), 685-706.
- McGrath, R. E. (2004). Saving our psychosocial souls. *American Psychologist, 59*(7), 644-645.
- McLeod, J.D., Pescosolido, B.A., Takeuchi, D.T., & Falkenberg White, T. (2004). Public attitudes toward the use of psychiatric medications for children. *Journal of Health and Social Behavior, 45*(3), 53-67.
- McNeill, T. (2006). Evidence-based practice in an age of relativism: Toward a model of practice. *Social Work, 51*(2), 147-156.
- Mech, E.V., Pryde, J.A., & Rycraft, J.R. (1995). Mentors for adolescents in foster care. *Child & Adolescent Social Work Journal, 12*(4), 317-328.
- Medco Health Solutions. (2004). *Medco Study Reveals Pediatric Spending Spike on Drugs to Treat Behavioral Problems*. Retrieved from: <http://medco.mediaroom.com/index.php?s=43&item=158>
- Medco Health Solutions. (2006). *2006 Drug Trend Report*. Retrieved from <http://medco.mediaroom.com/index.php?s=64&cat=5>
- Meyer, J. (2001). Novel antipsychotics and severe hyperlipidemia, *Journal of Clinical Psychopharmacology, 21*(4), 369-374.

- Meyers, L. (2007). Empty bottles: Easing clients off meds. *Monitor on Psychology*, 38(3), p. 20-21.
- Miklowitz, D. J., George, E. L., Axelson, D. A., Kim, E. Y., Birmaher, B., & Schneck, C., et al. (2004). Family-focused treatment for adolescents with bipolar disorder. *Journal of Affective Disorders*, 82(Suppl 1), S113-S128.
- Millstein, K. (2000). Confidentiality in direct social-work practice: Inevitable challenges and ethical dilemmas. *Families in Society*, 81(3), 270-282.
- Mintzes, B. (2002). Direct to consumer advertising is medicalising normal human experience. *British Medical Journal*, 324, 908-909.
- Mirowsky, J., & Ross, C. E. (1990). Subjective boundaries and combinations in psychiatric diagnosis. *Journal of Mind and Behavior*, 11(3/4), 407-424.
- Mizrahi, T., & Abramson, J. (2000). Collaboration between social workers and physicians: Perspectives on a shared case. *Social Work in Health Care*, 31(3), 1-24.
- Moffatt, B., & Elliott, C. (2007). Ghost marketing: Pharmaceutical companies and ghostwritten journal articles. *Perspectives in Biology and Medicine*, 50(1), 18-31.
- Moncrieff, J., & Cohen, D. (2005). Rethinking models of psychotropic drug action. *Psychotherapy & Psychosomatics*, 74, 145-153.
- Moore, T.J., Cohen, M.R., & Furberg, C.D. (2007). Serious adverse drug events reported to the Food and Drug Administration, 1998-2005. *Archives of Internal Medicine*, 167(16), 1752-1759.
- Moran, M. (2007). Developmental trauma merits DSM diagnosis, experts say. *Psychiatric News*, 42(3), 20. Retrieved from <http://pn.psychiatryonline.org/cgi/content/full/42/3/20?maxtoshow=&HITS=10&hits=10&RESULTFORMAT=&fulltext=Developmental+Trauma+Disorder+merits+DSM+diagnosis&searchid=1&FIRSTINDEX=0&sortspec=relevance&resourcetype=HWCIT>
- Moreno, C., Laje, G., Blanco, C., Jian, H., Schmidt, A.B., & Olfson, M. (2007). National trends in the outpatient diagnosis and treatment of bipolar disorder in youth. *Archives of General Psychiatry*, 64(9), 1032-1039.
- Morrison, A., Read, J., & Turkington, D. (2005). Trauma and psychosis: theoretical and clinical implications. *Acta Psychiatrica Scandinavica*, 112, 327-329.
- Moses, T., & Kirk, S. A. (2006). Social workers' attitudes about psychotropic drug treatment with youths. *Social work*, 51(3), 211-222.
- Moynihan, R. (2002). Drug firms hype disease as a sales ploy, industry chief claims. *British Medical Journal*, 324, 867.
- Moynihan, R., Heath, I., & Henry, D. (2002). Selling sickness: The pharmaceutical industry and disease mongering. *British Medical Journal*, 324, 886-891.
- MSNBC (2006, March 13). *States wrestle with medicating kids*. Retrieved from <http://www.msnbc.msn.com/id/17597241/>

- MSNBC (2006, May 3). Antipsychotic drug use among kids soars. Retrieved from <http://www.msnbc.msn.com/id/12616864>
- Murray, T.L. (2006). The other side of psychopharmacology: A review of the literature. *Journal of Mental Health Counseling*, 28(4), 309-337.
- Murphy, S. (2008, April 4). Doctor sued in death of girl, 4: Her psychiatrist treated her with powerful drugs. *The Boston Globe*. Retrieved on May 14, 2008 from http://www.boston.com/news/local/articles/2008/04/04/doctor_is_sued_in_death_of_girl_4/
- National Association of Social Workers. (1999). *Code of Ethics of the National Association of Social Workers*, Approved by the 1996 NASW Delegate Assembly and revised by the 1999 NASW Delegate Assembly. Washington, DC: Author. Retrieved from <http://www.socialworkers.org/pubs/code/code.asp>
- National Association of Social Workers. (2005). Social workers, medication, and scope of practice. Washington, DC: Author. Retrieved from http://www.socialworkers.org/ldf/legal_issue/200505.asp?print=1
- National Institute for Clinical Excellence. (2005). *Post-traumatic stress disorder: The management of PTSD in adults and children in primary and secondary care*. London: The Royal College of Psychiatrists & The British Psychological Society. Retrieved from <http://www.nice.org.uk/nicemedia/pdf/CG026fullguideline.pdf>
- National Institute of Mental Health. (2000). Child and Adolescent Bipolar Disorder: An update from the NIMH, (NIH Publication Number: NIH00-4778). Retrieved March 26, 2007, from <http://www.nimh.nih.gov/publicat/index.cfm>
- Ng, R. (2004). *Drugs—From discovery to approval*. New York: Wiley.
- Nilsen, W. (2007). Fostering futures: A preventive intervention program for school-age children in foster care. *Clinical Child Psychology and Psychiatry*, 12(1), 45-63.
- NJPIRG Law & Policy Center (2006). *Turning Medicine Into Snake Oil: How Pharmaceutical Marketers Put Patients at Risk*. NJPIRG Law & Policy Center [Report]: Trenton, NJ: Author. Retrieved from <http://www.njpirg.org/home/reports/report-archives/health-care/health-care/turning-medicine-into-snake-oil-how-pharmaceutical-marketers-put-patients-at-risk>
- Norcross, J.C., Beutler, L.E., & Levant, R.F. (Eds.). (2005). *Evidence-based practices in mental health: Debate and dialogue on the fundamental questions*. Washington, DC: American Psychological Association.
- Norfleet, M. A. (2002). Responding to society's needs: Prescription privileges for psychologists. *Journal of clinical psychology*, 58(6), 599-610.
- Office of the Inspector General (2003) *Medicaid's Mental Health Drug Expenditures. A Report by the Office of the Inspector General*. U.S. Department of Health and Human Services Report OEI-05-02-00080. Retrieved from <http://oig.hhs.gov/oei/reports/oei-05-02-00080.pdf>

- Ogden, P., & Minton, K. (2000). Sensorimotor psychotherapy: One method for processing traumatic memory. *Traumatology*, 6(3), 149-173.
- Ogden, P., Pain, C., & Fisher, J. (2006). A sensorimotor approach to the treatment of trauma and dissociation. *Psychiatric Clinics of North America*, 29(1), 263-279.
- Oldani, M.J. (2004). Thick prescriptions: Toward an interpretation of pharmaceutical sales practices. *Medical Anthropology Quarterly*, 18(3), 325-356.
- Olfson, M., Blanco, C., Liu, L., Moreno, C., & Laje, G. (2006). National trends in the outpatient treatment of children and adolescents with antipsychotic drugs. *Archives of General Psychiatry*, 63(6), 679-685.
- Olfson, M., Marcus, S.C., Weissman, M.M., & Jensen, P.S. (2002). National trends in the use of psychotropic medications by children. *Journal of the American Academy of Child and Adolescent Psychiatry*, 41(5), 514-21.
- Olson, J. (2007, August 27). Drug makers step up giving to Minnesota psychiatrists. *Twin Cities Pioneer Press*. Retrieved February 18, 2008 from: <http://mentalthopenews.blogspot.com/2007/08/drug-makers-step-up-giving-to-minnesota.html>
- Otis, H. G., & King, J. H. (2006). Unanticipated psychotropic medication reactions. *Journal of Mental Health Counseling*, 28(3), 218-240.
- Pappadopulos, E., Guelzow, T., Wong, C., Ortega, M., & Jensen, P. (2004). A review of the growing evidence base for pediatric psychopharmacology. *Child and Adolescent Psychiatric Clinics of North America*, 13, 817-855.
- Patel, N.C., Crismon, M.L., Hoagwood, K., & Jensen, P.S. (2005). Unanswered questions regarding antipsychotic use in aggressive children and adolescents. *Journal of Adolescent and Child Psychopharmacology*, 15(2), 270-284.
- Patel, N.C., Crismon, M.L., Hoagwood, K., Johnsrud, M.T., Rascati, K.L., Wilson, J.P. & Jensen, P.S. (2005). Trends in the use of typical and atypical antipsychotics in children and adolescents. *Journal of the Academy of Child & Adolescent Psychiatry*, 44(6), 548-556).
- Pathak, S., Arszman, S.P., Danielyan, A., Johns, E. S., Smirnov, A., Kowatch, R.A. (2004). Psychotropic utilization and psychiatric presentation of hospitalized very young children. *Journal of Child and Adolescent Psychopharmacology*, 14(3), 433-442.
- Pavuluri, M., Henry, D., Devineni, B., Carbray, J., Naylor, M., & Janicak, P. (2004). A pharmacotherapy algorithm for stabilization and maintenance of pediatric bipolar disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 43(7), 859-867.
- Pearlman, L. A., & Courtois, C. A. (2005). Clinical applications of the attachment framework: Relational treatment of complex trauma. *Journal of Traumatic Stress*, 18(5), 449-459.
- Petersen, M. (2002, November 22). Madison avenue has growing role in the business of drug research. *The New York Times*. Retrieved from:

<http://query.nytimes.com/gst/fullpage.html?sec=health&res=9F0DEEDF1239F931A15752C1A9649C8B63>

- Pharmaceutical Executive (2007) *Annual Report on the World's Top 50 Pharmaceutical Companies*. Advanstar Communications: Marietta, GA. Retrieved from <http://www.pharmexec.com/pharmexec/article/articleDetail.jsp?id=423201>
- PhRMA (2004). *Pharmaceutical marketing & promotion Q&A: Tough questions, straight answers*. Fall 2004. Pharmaceutical Researchers and Manufacturers of America, Washington, DC: Author.
- Piotrowski, C., & Doelker, R. (2001). Prescription privileges: Attitudes of clinical social workers. *Psychological reports*, 88(2), 606.
- Piotrowski, C., & Keller, J. W. (1996). Prescription privileges and training issues: A pilot study. *Psychological reports*, 78(2), 445-446.
- Pliszka, S., Crismon, L., Hughes, C., Conners, C., Emslie, G, et al. (2005). The Texas Children's Medication Algorithm Project: Revision of the algorithm for pharmacotherapy of attention-deficit/hyperactivity disorder, *Journal of the American Academy of Child and Adolescent Psychiatry*, 45(6), 642-657.
- Price, J.H. (2007, April 1). Spending on psychotropic drugs soars. *The Washington Times*. Retrieved from www.washingtontimes.com
- Pringle, E. (2006). *Kids dying from off-label use of antipsychotics*, Retrieved March, 28, 2007, from <http://www.lawyersandsettlements.com/articles/00183/antipsychotics.html>
- Pringle, E. (2007). Consequences of rampant off-label prescribing of Depakote, Retrieved March 28, 2007 from, <http://www.lawyersandsettlements.com/articles/00594/off-label-depakote.html>
- Pringle, E. (2007). *Makers of Zyprexa, Risperdal and Seroquel under fire*, Retrieved March, 28, 2007, from <http://www.lawyersandsettlements.com/articles/00660/zyprexa-medical-costs.html>
- Public Citizen (2002). *Drug advertising masquerades as education*. Press Release, February 13, 2002. Retrieved from <http://www.citizen.org/pressroom/release.cfm?ID=1022>
- Quick, J. (2001). Maintaining the integrity of the clinical evidence base. *Bulletin of the World Health Organization*, 79, 1093.
- Racusin, R., Maerlender, A.C., Sengupta, A., Isquith, P.K., & Straus, M.B. (2005). Psychosocial treatment of children in foster care: A review. *Community Mental Health Journal*, 41(2), 199-221.
- Raeburn, P. (2007, May 30). Kids on meds – Trouble ahead. Antidepressants, designed for adults, may be altering the brains of kids who take them. *Scientific American*. Retrieved from

<http://www.sciam.com/article.cfm?chanID=sa017&articleID=D901A079-E7F2-99DF-37A3A8CBB9CC8A6F&ref=rss>

- Raghavan, R., Zima, B. T., Andersen, R. M., Leibowitz, A. A., Schuster, M. A., & Landsverk, J. (2005). Psychotropic medication use in a national probability sample of children in the child welfare system. *Journal of Child and Adolescent Psychopharmacology. Special Issue on Psychopharmacoeconomics*, 15(1), 97-106.
- Ramos, A., Shytle, R., & Silver, A. (2003). Ziprasidone-induced oculogyric crisis. *Journal of the American Academy of Child and Adolescent Psychiatry*, 42(9), 1013-1014.
- Rappaport, N., & Chubinsky, P. (2000). The meaning of psychotropic medications for children, adolescents and their families. *Journal of the Academy of Child and Adolescent Psychiatry*, 39(9), 1198-1200.
- Rappley, M.D. (2006). Actual psychotropic medication use in preschool children. *Infants & Young Children*, 19(2), 154-163.
- Read, J. (2005). The bio-bio-bio model of madness. *The Psychologists*, 18(10), 596-597.
- Read, J., & Ross, C.A. (2003). Psychological trauma and psychosis: Another reason why people diagnosed schizophrenic must be offered psychological therapies. *Journal of the American Academy of Psychoanalysis and Dynamic Psychiatry*, 31(1), 248-268.
- Read, J., Mosher, L.R., & Bentall, R.P. (2004). *Models of madness: psychological, social and biological approaches to schizophrenia*. New York: Brunner Routledge.
- Read, J., van Os, J., Morrison, A.P., & Ross, C.A. (2005). Childhood trauma, psychosis and schizophrenia: A literature review with theoretical and clinical implications. *Acta Psychiatrica Scandinavica*, 112, 330-350.
- Reamer, F.G. (2001). Ethics and managed care policy. In N.W. Veeder & W. Peebles-Wilkins (Eds.). *Managed care programs: Policy, programs and research*. New York: Oxford University Press.
- Regehr, C., Stern, S., Shlonsky, A. (2007). Operationalizing evidence-based practice: the development of an institute for evidence-based social work. *Research on Social Work Practice*, 17(3), 408-416.
- Reist & VandeCreek (2004). The pharmaceutical industry's use of gifts and educational events to influence prescription practices: ethical dilemmas and implications for psychologists. *Professional Psychology: Research & Practice*, 35(4), 329-335.
- Relman (2001). Separating continuing medical education from pharmaceutical marketing. *JAMA*, 285(15), 2009-2012
- Rhodes, J.E., Haight, W.L., & Briggs, E.C. (1999). The influence of mentoring on the peer relationships of foster youth in relative and nonrelative care. *Journal of Research on Adolescence*, 9(2), 185-201.

- Rhodes, J.E., Spencer, R., Keller, T.E., Lian, B., & Noam, G. (2006). A model for the influence of mentoring relationships on youth development. *Journal of Community Psychology, 34*(6), 691-707.
- Riddle, M.A., Kastelic, E.A. & Frosch, E. (2001). Pediatric psychopharmacology. *Journal of Child Psychology & Psychiatry, 42*(1), 73-90.
- Rivara, F. P., & Cummings, P. (2002). Publication bias: The problem and some suggestions. *Archives of Pediatric and Adolescent Medicine, 156*, 424-425.
- Rivas-Vasquez, R. A., Johnson, S. L., Blais, M. A., & Rey, G. J. (1999). Selective serotonin reuptake inhibitor syndrome: Understanding, recognition, and management for psychologists. *Professional Psychology: Research and Practice, 30*, 464-469.
- Rivas-Vazquez, R. A., & Mendez, C. I. (2002). Overview of the drug approval process. *Professional Psychology: Research & Practice, 33* (5), 502-505.
- Roberts, M. (1999). Risperdal and parkinsonian tremor, *Journal of the American Academy of Child and Adolescent Psychiatry, 38*(3), 230.
- Rodriguez, E. (2005, March 9). Bill guides DCF on kids' drugs. *The Miami Herald*, Page 1B.
- Rohde, P., Clarke, G. N., Mace, D. E., Jorgensen, J. S., & Seeley, J. R. (2004). An Efficacy/Effectiveness study of cognitive-behavioral treatment for adolescents with comorbid major depression and conduct disorder. *Journal of the American Academy of Child & Adolescent Psychiatry, 43*(6), 660-668.
- Ross, J.S., Lackner, J.E., Lurie, P., Gross, C.P., Wolfe, S., & Krumholz, H.M. (2007). Pharmaceutical company payments to physicians: Early experience with disclosure laws in Vermont and Minnesota. *JAMA, 297*(11), 1216-1223.
- Roth, A., & Fonagy, P. (1996). *What works for whom? A critical review of psychotherapy research*. New York: The Guilford Press.
- Ryan, N., Bhatara, V., & Perel, J. (1999). Mood stabilizers in children and adolescents, *Journal of the American Academy of Child and Adolescent Psychiatry, 38*(5), 529-536.
- Safer, D. J. (2002). Design and reporting modifications in industry-sponsored comparative psychopharmacology trials. *Journal of Nervous and Mental Disease, 190*(9), 583-592.
- Safer, D. J., Zito, J. M., & dosReis, S. (2003). Concomitant psychotropic medication for youths. *American Journal of Psychiatry, 160*(3), 438-449.
- Sanua, V. (2003). The "science" of psychopharmacology? Report from a 5-day seminar. *Ethical Human Sciences and Services, 5*(2), 153-156.
- Saunders, B.E., Berliner, L., & Hanson, R.F. (Eds.). (2004). *Child Physical and Sexual abuse: Guidelines for Treatment (Revised Report: April 26, 2004)*. Charleston, SC: National Crime Victims Research and Treatment Center. Retrieved from <http://www.musc.edu/cvc/>

- Scahill, L., Leckman, J., Schultz, R., Katsovich, L., & Peterson, B. (2003). A placebo-controlled trial of risperidone in Tourette syndrome, *Neurology*, *60*, 1130-1135.
- Scheffer, R., Kowatch, R., Carmody, T., & Rush A. (2005). Randomized, placebo-controlled trial of mixed amphetamine salts for symptoms of comorbid ADHD in pediatric bipolar disorder after mood stabilization with divalproex sodium, *American Journal of Psychiatry*, *162*(1), 58-64.
- Schofield, G., & Beek, M. (2005). Risk and resilience in long-term foster care. *British Journal of Social Work*, *35*, 1283-1301.
- Schore, A. N. (2001). The effects of early relational trauma on right brain development, affect regulation, and infant mental health. *Infant Mental Health Journal, Special Issue: Contributions from the decade of the brain to infant mental health*, *22*(1-2), 201-269.
- Schore, A.E. (1994). *Affect regulation and the origin of the self: The neurobiology of emotional development*. Hillsdale, NJ: Lawrence Erlbaum Associates, Inc.
- Schore, A.E. (2003). *Affect dysregulation & disorders of the self*. New York: W.W. Norton & Company.
- Schwartz, T. L., Kuhles, D. J., Wade, M., & Masand, P. S. (2001). Newly admitted psychiatric patient prescriptions and pharmaceutical sales visits. *Annals of Clinical Psychiatry*, *13*(3), 159-162.
- Scovel, K. A., Christensen, O. J., & England, J. T. (2002). Mental health counselors' perceptions regarding psychopharmacological prescriptive rights. *Journal of Mental Health Counseling*, *24*(1), 36-50.
- Sharav, V. (2007, March 20). The damaging impact of PDUFA and why it should be repealed. Infomail of the Alliance for Human Research Protection. Available at: <http://www.ahrp.org/cms/content/view/498/94/>
- Sharp, B. (2007). CMPA ADHD and comorbid aggression algorithm. *Journal of the American Academy of Child and Adolescent Psychiatry*, *46*(1), 1.
- Shaw, P., & Rapoport, J. L. (2006). Decision making about children with psychotic symptoms: Using the best evidence in choosing a treatment. *Journal of the American Academy of Child and Adolescent Psychiatry*, *45*(11), 1381.
- Shojania, K. G. (2006). Safe medication prescribing and monitoring in the outpatient setting. *Canadian Medical Association journal*, *174*(9), 1257-1258.
- Siebert, A. (2000). How non-diagnostic listening led to a rapid "recovery" from paranoid schizophrenia: What is wrong with psychiatry? *Journal of Humanistic Psychology*, *40*(1), 34-58.
- Silva, R., Munoz, D., Alpert, M., Perlmutter, I., & Diaz, J. (1999). Neuroleptic malignant syndrome in children and adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, *38*(2), 187-194.

- Singh, M., DelBello, M., & Adler, C. (2007). Acute dystonia associated with aripiprazole in a child. *Journal of the American Academy of Child and Adolescent Psychiatry*, 46(3), 306-307.
- Skarpathiotakis, M., & Westreich, N. (2005). NMS after clozapine initiation. *Journal of the American Academy of Child and Adolescent Psychiatry*, 44(11), 1101-1102.
- Slavin, P. (2004). Teamwork key in managing medication: Social workers' role vital in promoting adherence. *NASW News*, 49(6) [online]. Retrieved from <http://www.socialworkers.org/pubs/news/2004/06/teamwork2.asp?back=yes>
- Smith, R. (2003). Medical journals and pharmaceutical companies: Uneasy bedfellows. *British Medical Journal*, 326, 1202-1205.
- Smith, R. (2005). Medical journals are an extension of the marketing arm of pharmaceutical companies. *PLoS Medicine*, 2(5), e138.
- Smyer, M. A., Balster, R. L., Egli, D., & Johnson, D. L. (1993). Summary of the report of the ad hoc task force on psychopharmacology of the American psychological association. *Professional Psychology: Research and Practice*, 24(4), 394-403.
- Spetie, L., & Arnold, E. (2007). Ethical issues in child psychopharmacology research and practice: Emphasis on preschoolers, *Psychopharmacology*, 191, 15-26.
- Springer, D. W. (2006). *Treating juvenile delinquents with conduct disorder, ADHD, and oppositional defiant disorder*. In Roberts, A., & Yeager, K. (Eds.), *Foundations of evidence-based social work practice*. New York: Oxford University Press.
- St. Luke's Health Initiatives (2006) *FlashPoint Issue Brief: Children, Adolescents and Psychotropic Medications*. Retrieved from http://www.slhi.org/publications/issue_briefs/pdfs/ib-2006-August.pdf
- Stagnitti, M.N. (2007). *Trends in the use and expenditures for the therapeutic class prescribed psychotherapeutic agents and all subclasses, 1997 and 2004*. Statistical Brief #163. Agency for Healthcare Research and Quality, Rockville, MD.
- Stone, M., & Jones, M. L. (2006, November 17). *Clinical review: Relationship between antidepressant drugs and suicidality in adults*. Rockville, MD: Food and Drug Administration.
- Strom, B. L. (2006). How the US drug safety system should be changed. *JAMA*, 295(17), 2072-2075.
- Strom-Gottfried, K. (1998). Informed consent meets managed care. *Health & Social Work*, 23(1), 25-33.
- Strom-Gottfried, K. (1998). Is "ethical managed care" an oxymoron? *Families in Society*, 79(3), 297-307.
- Tatz, D. & Reinert, S. (2007, February 6). Everyone failed Rebecca: A special report. *The Patriot Ledger*. Retrieved from <http://www.southofboston.net/specialreports/rebecca/pages/020607.asp>

- Taylor, M. F., & Bentley, K. J. (2005). Professional dissonance: Colliding values and job tasks in mental health practice. *Community mental health journal*, 41(4), 469-480.
- The Center for Public Integrity (2007). *Spending on lobbying thrives -- Drug and health products industries invest \$182 million to influence legislation*. Retrieved from <http://www.publicintegrity.org/rx/report.aspx?aid=823>
- The Week* (2006). The corruption of medicine, *The Week*, 09/22/2006, Retrieved from <http://www.theweekmagazine.com/news/articles/news.aspx?ArticleID=1653>
- Thomas, C. P., Conrad, P., Casler, R., & Goodman, E. (2006). Trends in the use of psychotropic medications among adolescents, 1994 to 2001. *Psychiatric Services*, 57(1), 63-69.
- Troost, P., Lahuis, B., Steenhuis, M., Ketelaars, C., Buitelaar, J., et al. (2005). Long-term effects of risperidone in children with autism spectrum disorders: A placebo discontinuation study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 44(11), 1137-1144.
- Troyen A.B., Rothman, D.J., Blank L., Blumenthal, D., Chimonas, S.C., Cohen, J.J., Goldman, J., Kassirer, J.P., Kimball, H., Naughton, J., & Smelser, N. (2006). Health industry practices that create conflicts of interest: A policy proposal for academic medical centers, *JAMA*, 295(4), 429-433
- Tucker, G.J. (1998). Putting DSM-IV in perspective. *American Journal of Psychiatry*, 152(2), 159-161.
- Tulkin, S. R., & Stock, W. (2004). A model for predoctoral psychopharmacology training: Shaping a new frontier in clinical psychology. *Professional Psychology: Research and Practice*, 35(2), 151-157.
- Turner, E. H., Matthews, A. M., Linardatos, E., Tell, R. A., & Rosenthal, R. (2008). Selective publication of antidepressant trials and its influence on apparent efficacy. *New England Journal of Medicine*, 358, 252-260.
- USA Today (2006, May 2). *For kids, oversight of prescriptions is scarce*. Retrieved from http://www.usatoday.com/news/health/2006-05-01-antipsychotics-foster-kids_x.htm
- van der Kolk, B.A. (2003). The neurobiology of childhood trauma and abuse. *Child and Adolescent Psychiatric Clinics of North America*, 12, 293-317.
- van der Kolk, B.A. (2005). Developmental trauma disorder: toward a rational diagnosis for children with complex trauma histories. *Psychiatric Annals*, 35(5), 401-408.
- van der Kolk, B.A. (2006). Clinical implications of neuroscience research in PTSD. *Annals of the New York Academy of Science*, 1071, 277-293.
- van der Kolk, B.A., & Courtois, C. A. (2005). Editorial comments: Complex developmental trauma. *Journal of Traumatic Stress*, 18(5), 385-388.
- van der Kolk, B.A., & Fisler, R. (1994). Childhood abuse and neglect and loss of self-regulation. *Bulletin of the Menninger Clinic*, 58(2), 145-168.

- van der Kolk, B.A., McFarlane, A., & Weisaeth, L. (1996). *Traumatic stress: The effects of overwhelming experience on mind, body, and society*. New York: Guilford Press.
- van der Kolk, B.A., Pelcovitz, D., Roth, S., Mandel, F.S., McFarlane, A., & Herman, J.L. (1996). Dissociation, affect dysregulation and somatization. *American Journal of Psychiatry*, 153, 83-93.
- van der Kolk, B.A., Roth, S., Pelcovitz & Spinazzola, J. (2005). Disorders of extreme stress: The empirical foundation of a complex adaptation to trauma. *Journal of Traumatic Stress*, 18(5), 389-399.
- VanderVen, K. (2004). Adults are still needed! Intergenerational and mentoring activities. *Reclaiming Children and Youth: The Journal of Strength-based Interventions*, 13(2), 94.
- Vedantam, S. (2006, April 12). Comparison of schizophrenia drugs often favors firm funding study. *Washington Post*, page A.01. Retrieved from <http://www.washingtonpost.com>
- Velting, O. N., Setzer, N. J., & Albano, A. M. (2004). Update on and advances in assessment and cognitive-behavioral treatment of anxiety disorders in children and adolescents. *Professional Psychology: Research and Practice*, 35(1), 42-54.
- Verispan (2005). Pharmaceutical marketing scrutinized: More than 30 states debate disclosure or oversight. Press Release, August 22, 2005. Retrieved from http://www.verispan.com/about/press_release_details.php?id=gt6ts2crt9
- Viale, P. H. (2003). What nurse practitioners should know about direct-to-consumer advertising of prescription medications. *Journal of the American Academy of Nurse Practitioners*, 15(7), 297-304
- Vitiello, B. (2001). Psychopharmacology for young children: Clinical needs and research opportunities. *Pediatrics*, 108(4), 983-989.
- Vitiello, B., Zuvekas, S.H., & Norquist, G.S. (2006). National estimates of antidepressant medication use among U.S. children, 1997-2002. *Journal of the American Academy of Child and Adolescent Psychiatry*, 45(3), 271-279.
- Waller, R.J., Lewellen, K., & Bresson, D. (2005). The debate surrounding psychotropic medication usage in young children. *School Social Work Journal*, 29(2), 53-61.
- Walsh, J., Farmer, R., Taylor, M. F., & Bentley, K. J. (2003). Ethical dilemmas of practicing social workers around psychiatric medication: Results of a national study. *Social Work in Mental Health*, 1(4), 91-105.
- Washington Post (2006, January 25). Distance sought between doctors and drug industry. *Washington Post*, Page A08. Retrieved from www.washingtonpost.com
- Waters, R. (2007, September 17). Hyperactivity drugs to be studied for heart risk. Bloomberg. Retrieved from <http://www.bloomberg.com/apps/news?pid=newsarchive&sid=a5ap6JbK5NDA#>

- Wazana (2000). Physicians and the pharmaceutical industry: is a gift ever just a gift? *JAMA*, 283(3), 373-380.
- Webster-Stratton, C., & Reid, M.J. (2003). The incredible years. Parents, teachers and children training series: A multifaceted treatment approach for young children with conduct problems. In Kazdin, A.E. & Weisz, J.R., (eds.), *Evidence-based psychotherapies for children and adolescents* (pp. 224-240). New York: Guilford Press.
- Weene, K. A. (2002). The psychologist's role in the collaborative process of psychopharmacology. *Journal of clinical psychology*, 58(6), 617-621.
- Wiggins, J. G., & Wedding, D. (2004). Prescribing, professional identity, and costs. *Professional Psychology: Research and Practice*, 35(2), 148-150.
- Williams, W. I. (2006). Complex trauma: Approaches to theory and treatment. *Journal of Loss & Trauma*, 11(4), 321-335.
- Wolfe, D.A., & Mash, E.J. (Eds.). (2006). *Behavioral and emotional disorders in adolescents: Nature, Assessment, and Treatments*. New York: The Guilford Press.
- Wong, I. C. K., Murray, M. L., Camilleri-Novak, D., & Stephens, P. (2004). Increased prescribing trends of paediatric psychotropic medications. *Archives of Disease in Childhood*, 89(12), 1131-1132.
- Yehya, N., Saldarini, C., Koski, M., & Davanzo, P. (2004). Valproate-induced hyperammonemic encephalopathy. *Journal of the American Academy of Child and Adolescent Psychiatry*, 43(8), 926-927.
- Zalesky, C. D. (2006). Pharmaceutical marketing practices: Balancing public health and law enforcement interests; moving beyond regulation-through-litigation. *Journal of Health Law*, 39(2), 235-264
- Zito, J. M., & Safer, D. J. (2001). Services and prevention: Pharmacoepidemiology of antidepressant use. *Biological Psychiatry*, 49(12), 1121-1127.
- Zito, J. M., Safer, D. J., dosReis, S., Gardner, J. F., Boles, M., & Lynch, F. (2000). Trends in the prescribing of psychotropic medications to preschoolers. *JAMA*, 283(8), 1025-1030.
- Zito, Safer, dosReis et al. (2003). Psychotropic practice patterns for youth: a 10-year perspective. *Archives of Pediatrics & Adolescent Medicine*, 157(1), 17-25.

Facing Foster Care in Alaska

Mental Health Services and Foster Care



Facing Foster Care in Alaska Mental Health Services


Our Mission is to improve the foster care system through sharing our experiences, supporting and educating youth and social services and implementing positive change in society as a whole.

In 2003, a group of dedicated foster care youth and alumni from across Alaska came together to share their issues and form a Youth Advisory Board. In 2004, the group adopted the name Facing Foster Care in Alaska (FFCA). FFCA is dedicated to improving the lives of children and youth in foster care through developing and sustaining a statewide organization that will continue to work towards supporting foster youth and improving the foster care system. FFCA members are dedicated to advocating for improvements in the areas of education, mental health, permanency, disproportionality, independent living, and the overall well-being of children and youth in Alaska's foster care system.

Since their inception, the members of FFCA have been speaking out about mental health and treatment services of children and youth in foster care, and offering alternative solutions. With overwhelming feedback from youth and alumni of foster care regarding the issues of mental health and treatment services, the members FFCA came together in November of 2008 to brainstorm ideas and create possible solutions or alternatives to traditional treatment. This document encompasses their ideas and gives insight into what youth and alumni across Alaska believe is the best way to deal with the issues they face when being placed in foster care.

The content of this document reflects only the opinions and ideas of Facing Foster Care in Alaska members, and should not be viewed as the opinions or ideas from any state or private agencies.

Facing Foster Care in Alaska Mental Health Services



Youth and alumni were asked to share thoughts and ideas around what came to mind when they heard the words, "Mental Health" and "Treatment Services."

Facing Foster Care in Alaska Mental Health Services

Alternatives and Possible Solutions

The decreased use of psychiatric medications

In their 2008 Policy Agenda, FFCA members called for Decreased use of Psychotropic Medication for Alaska's foster youth. Many of Alaska's youth and alumni complain about being prescribed psychotropic medications after entering the foster care system for symptoms of depression, anxiety, trauma, attachment issues, and misbehavior. The youth and alumni of FFCA feel that these are all normal symptoms of child maltreatment and dealing with all that comes out of being placed in foster care. There has been a national focus on the use of psychotropic medications being over-prescribed for children and youth in foster care. FFCA members have also complained about side-effects caused by these medications resulting in a decreased ability to focus on their education as well as function in everyday society. The youth and alumni of FFCA would like to see that the prescription of psychotropic medications for Alaska's foster children and youth is decreased and reviewed more closely.

The right to be informed

Many of Alaska's foster children and youth don't know their rights in regards to mental health and treatment services. The members of FFCA believe that service providers should inform children and youth in foster care about their rights in regards to their treatment plan. During the November 2008 FFCA retreat, one member commented that he did know his rights, but if he did refuse medication he would be placed in North Star. FFCA members would like to see that all children and youth in foster care are informed of their rights and the repercussions if they choose not to comply with their treatment plan.

Facing Foster Care in Alaska Mental Health Services

Building Relationships

The members of FFCA believe that one of the best ways to deal with the emotional issues they face when entering foster care is through building and maintaining healthy relationships with family, friends, and permanent connections with a caring adult. The bulleted list below outlines the ideas FFCA members came up with regarding relationships.

- Trust Building opportunities
- Freedom
- Personal time
- Family Relationships/connections/visits
- Mentors
- Having good listeners
- Supporting dreams/hopes
- Non-judgmental relationships
- Permanent connection
- Getting to know us
- Talking/venting
- Acknowledgment
- Praise
- Constant affective communication
- Not relishing diagnosis/medication
- Cut out the unnecessary
- Understanding
- More homes
- Preparation for the real world
- Being placed in a stable understating home
- Extra time with peers
- Keeping siblings together
- Listen to what we have to say
- Pay attention to our needs
- We need more communication w/family & friends
- Do not separate youth and children
- If meds are absolutely necessary inform us what there for and what the side effects are
- Effective communication with social workers and GAL's

Creating a plan with the client/self determination

Over the years, FFCA members have continuously complained about treatment plans being written for them rather than with them. Many of the youth and alumni speak out about how they have no idea what's in their treatment plan or case plan. The members of FFCA believe that they cannot affectively work on their treatment plan or case plan if they do not know what is in it or don't have a say in the process of creating it. FFCA would like to see that service providers are working with children and youth to develop a plan that outlines what the children and youth feel they need to work on in order to become productive members of society.

Facing Foster Care in Alaska Mental Health Services

The right to be “Normal”

Many of Alaska’s foster children and youth complain about standing out among their peers and not being able to participate in the same activities as other young people. They say they are constantly going to appointments for counseling, medication adjustments, group therapy, family therapy, ect. The members of FFCA believe that the best treatment for depression, anxiety, attachment issues, and other behaviors that are often diagnosed in foster children and youth, can be treated by giving them the opportunity to be involved with school, community, and family events. The bulleted lists below outlines the ideas that FFCA members came up with as alternatives to the various therapy and other treatment related appointments.

Extra Curricular Activities

- Sports/Clubs
- Banking/financial skills
- Therapeutic activities
- Massage
- Journaling
- Self advocacy/empowerment
- No drugs
- Foster home with a pet
- FFCA
- Service Projects

Self Expression

- Star watching
- Poetry
- Music
- Art
- Cooking
- Life skills
- Freedom/choice

Facing Foster Care in Alaska Mental Health Services

Definitions (According to FFCA)

Youth- A young person in foster care

Alumni- A person who was in foster care at some point during their life

For more information about Facing Foster Care in Alaska please visit

http://www.alaskacasa.org/facing_foster_care_in_alaska.htm

or

<http://www.myspace.com/ffca>

This document was drafted by FFCA President Amanda Metivier using the feedback from the FFCA members that attended that November 2008 FFCA retreat in Anchorage. For questions regarding the content of this document please contact Amanda Metivier at facing_fostercare@yahoo.com or call 230-8237.

James B. Gottstein, Esq.

406 G Street, Suite 206
Anchorage, Alaska 99501
Jim.Gottstein@psychrights.org
James.B.Gottstein@gottsteinlaw.com
jg@touchngo.com
907-274-7686

Curriculum Vitae

(September 12, 2008)

Experience:

1985-Present: Owner - Law Offices of James B. Gottstein
1995-Present: CEO - Touch N' Go Systems, Inc.
2002-Present: President/Chief Executive Officer (since 2005) & Board Member, Law Project for Psychiatric Rights (PsychRights®)
2004-Present: Treasurer & Board Member of the Board of Directors, National Association of Rights Protections and Advocacy (NARPA)
2005-Present: Board Member, International Center for the Study of Psychiatry and Psychology (ICSPP)
2002-Present: Board Member, Peer Properties, Inc., (currently President).
1991-Present: Board Member, The Gottstein Family Foundation
2003-2007: President & Board Member, CHOICES, Inc.
2003-2007: President & Board Member, Soteria-Alaska, Inc.
2003-2005: Board of Directors - Alaska World Affairs Council
1998-2004: Member - Alaska Mental Health Board
1998-2002: Board of Directors - Alaska Mental Health Consumer Web
1986-1996: Board of Directors - Mental Health Consumers of Alaska
1983-1985: Staff Attorney - Carr-Gottstein Inc.
1982-1983: Associate Attorney - Delaney, Wiles, Hayes, Reitman and Brubaker
1980-1982: Partner - Goldberg and Gottstein
1978-1980: Associate Attorney - Robert M. Goldberg and Associates

Bar Memberships

1994: United States Supreme Court
1978: Alaska
1978: United States District Court, District of Alaska
1978: United States Court of Appeals; Ninth Circuit

Education:

1978 (Class of '77): J.D., Harvard Law School
1974: B.S., Business Administration with Honors (Finance), University of Oregon.

Publications:

- Involuntary Commitment and Forced Psychiatric Drugging In the Trial Courts: Rights Violations as a Matter of Course, *25 Alaska Law Review* 51 (2008).


- How the Legal System Can Help Create a Recovery Culture in Mental Health Systems, contributed chapter in R. Halgin, *Clashing Views in Abnormal Psychology*, 5th Ed., McGraw Hill, New York, New York, (2008): 17-29.
- Rights and Alternatives: Enforcing legal rights as a mechanism for creating non-medical model alternatives, contributed chapter in *Beyond Psychiatry* (2007), P. Lehmann & P. Stastny, Eds., 308-17.
- Psychiatrists' Failure to Inform: Is There Substantial Financial Exposure? James B. Gottstein, Esq., JD, *Ethical Human Psychology and Psychiatry*, Volume 9, Number 2, 2007.

Presentations

- "Advocacy and the Transformation Triangle," International Network Towards Alternatives in Recovery (INTAR) conference on International Recovery Perspectives: Action on Alternatives, University of Toronto, June 6, 2008, Toronto, Canada.
- Panel member Soteria--The Proven Model for Recovery Communities, "The Development of Soteria-Alaska," INTAR conference on International Recovery Perspectives: Action on Alternatives, University of Toronto, June 5, 2008, Toronto, Canada.
- "Over-Prescription of Psychiatric Drugs: Changing an Irrational Policy," Contemporary Social Policy and Change, University Alaska Anchorage, April 2008.
- Roundtable, Program in Psychiatry and the Law @ BIDMC Psychiatry of Harvard Medical School, October 24, 2007, Boston, Massachusetts.
- "Forced Psychiatric Drugging: A Misguided Atrocity," Hampshire College, Massachusetts, October 22, 2007.
- "The Transformation Triangle," Region Ten Consumer Advisory Council 2nd Annual Conference, Charlottesville, Virginia, October 17, 2007.
- "The Psychiatric Drugging of America's Children: Legal Rights of Children and Parents," ICSPP Annual Conference, October 14, 2007.
- *Myers, Wetherhorn & More: Litigating for Consumer Driven Services*, Mental Health Consumer and Family Education & Leadership Conference, May 17, 2007.
- Psychiatric Drugs in America: Who's Crazy? Or Through the DSM Looking Glass, ANT A655 Advanced Medical Anthropology, March 26, 2008, University of Alaska Anchorage.
- Strategic Litigation to Achieve Meaningful Change: The *Myers* Case, Alaska and a National Initiative, National Association of Rights Protection and Advocacy (NARPA) annual conference, November 16, 2006, Baltimore, Maryland.
- "The Public Mental Health System, University of Alaska Anchorage, November 2, 2006.
- The Public Mental Health System: Analysis and Suggestions for Improvement, Alaska Pacific University, October 26, 2006, Anchorage, Alaska.

- A Coordinated Campaign To Successfully Change the Mental Illness System, ICSP annual conference, October 9, 2006, October 9, 2006.
- Panel member: "Free Your Mind A discussion about psychiatric rights and how we value people in our communities," David A. Clarke School of Law, Washington, DC, October 6, 2006.
- "CHOICES, Inc., Soteria-Alaska, & Peer Properties: Agents for Positive Change," Harvard & Yale Clubs, April 4, 2006, Anchorage, Alaska.
- "CHOICES, Inc., Soteria-Alaska, & Peer Properties: Agents for Positive Change," Alaska Mental Health Consumer Web, March 31, 2006, Anchorage, Alaska.
- "CHOICES, Inc., Soteria-Alaska, & Peer Properties: Agents for Positive Change," NAMI-Anchorage Annual Meeting, March 16, 2006.
- "Multi-faceted Grassroots Efforts to Bring About Meaningful Change to Alaska's Mental Health Program," National Association of Rights Protection and Advocacy (NARPA) annual conference, November 19, 2005, Hartford, Connecticut.
- "Involuntary Mental Health 'Treatment:' Utilizing Valid Scientific Information and Client Views to Win Cases," Massachusetts, Mental Health Law Unit Continuing Legal Education for the Committee for Public Counsel Services, November, 2005, Boston, Massachusetts.
- Member of Panel Discussion of Involuntary Treatment in the U.S., International Society for the Psychological Treatment of Schizophrenia and Other Psychoses, November, 2005, Boston, Massachusetts.
- "Grass Roots, Multi-Organizational Efforts in Support of Human Rights in Mental Illness," International Center for the Study of Psychiatry and Psychology, Flushing, NY, October 8, 2005.
- "Involuntary Commitment and Medication in Alaska: or Just Because I'm Paranoid Doesn't Mean They Aren't After Me," SWK 643 Human Diversity and Special Populations, University of Alaska Anchorage, September 19, 2005.
- "Human Rights in Mental Health: Let's Do It," MindFreedom Action Conference, April 30, 2005, Washington, DC.
- "Forced Psychiatric Drugging in Alaska," presented to the Health Law Section of the Alaska Bar Association, February 3, 2005.
- "PsychRights' Legal Campaign Against Forced Drugging and How You Can Participate," International Center for the Study of Psychiatry and Psychology, Flushing, New York, October 10, 2004.
- "Involuntary Commitment in Alaska and Beyond," Alaska Libertarian State Convention, April 24, 2004, Anchorage, Alaska.
- "The Law Project for Psychiatric Rights: Progress and Directions," for the National Rights Protection and Advocacy (NARPA) Conference, November 23, 2003, Austin, Texas.

- Co-presented Alaska Mental Health Board's FY 2005 Request for Recommendations to the Alaska Mental Health Trust Authority, August, 2003.
- Alaska Mental Health Board Budget Committee's "Mental Health Budget Summit," parts 1 and 2 on March 8, 2003, in Juneau Alaska and April 11-12, 2003 in Anchorage, Alaska, respectively.
- "Unwarranted Court Ordered Medication: A Call to Action," for the National Rights Protection and Advocacy (NARPA) Conference (off-agenda), November, 2002, Portland, Oregon.
- Board Training at the Family and Consumer Conference, May 2, 2002, in Anchorage, Alaska.
- "Real Estate in the Land of the Midnight Sun," presented to the International Real Estate Society's International Congress, July 27, 2001, Girdwood, Alaska.
- "Protecting Your Privacy On-Line: Privacy, Threats, and Countermeasures," co-presented with Lara Baker, September 7, 2001, in Anchorage, Alaska.
- Board Training at the Family and Consumer Conference, April 13, 2001, in Anchorage, Alaska.
- Alaska Mental Health Board's Program and Evaluation Committee's Report on the Alaska Psychiatric Institute, presented as chair of the committee to the full board, February 16, 2000, Fairbanks, Alaska.
- Internet Strategies for the Paralegal in Alaska: A paralegals guide to the information super-highway," for the Institute for Paralegal Education, November 1998, in Anchorage, Alaska.
- "Grant Writing, 1, 2, 3," various times 1998-2002, in Anchorage, Alaska.
- "Web Page Design," Alaska Mental Health Consumer Web various times, 1998-2002.
- "Lawyers and the Internet," October 23, 1995, in Anchorage, Alaska.
- Many public presentations on the Alaska Mental Health Trust Lands litigation and proposed settlements, including radio and television appearances and testimony to legislative committees, 1986-1994



RECEIVED

JUN 29 2007

Chambers of
Dana Fabe
Chief Justice

Supreme Court
State of Alaska

303 K Street
Anchorage, Alaska
99501-2083
(907) 264-0622
FAX (907) 264-0554

June 28, 2007

James Gottstein
Law Office of James B Gottstein
406 G St., Suite 206
Anchorage, AK 99501

Dear Mr. *Jm* Gottstein:

Thank you for agreeing to serve on the Probate Rules Subcommittee on Involuntary Commitments and the Involuntary Administration of Psychotropic Medication. We look forward to receiving your help and expertise in revising the procedural rules that govern these difficult and important cases. Judge Morgan Christen will be chairing the subcommittee, and her staff will be contacting you to schedule the first meeting. Again, thank you for your assistance in this important work.

Sincerely,

Dana Fabe
Chief Justice

DF:jmh

cc: Stephanie Cole, Administrative Director
Judge Morgan Christen
Doug Wooliver, Administrative Attorney

The Washington Post

Debate Over Drugs For ADHD Reignites

Long-Term Benefit For Children at Issue

By Shankar Vedantam
Washington Post Staff Writer
Friday, March 27, 2009; A01

New data from a large federal study have reignited a debate over the effectiveness of long-term drug treatment of children with hyperactivity or attention-deficit disorder, and have drawn accusations that some members of the research team have sought to play down evidence that medications do little good beyond 24 months.

The study also indicated that **long-term use of the drugs can stunt children's growth.**

The latest data paint a very different picture than the study's positive initial results, reported in 1999.

One principal scientist in the study, psychologist William Pelham, said that **the most obvious interpretation of the data is that the medications are useful in the short term but ineffective over longer periods but added that his colleagues had repeatedly sought to explain away evidence that challenged the long-term usefulness of medication.** When their explanations failed to hold up, they reached for new ones, Pelham said.

"The stance the group took in the first paper was so strong that the people are embarrassed to say they were wrong and we led the whole field astray," said Pelham, of the State University of New York at Buffalo. Pelham said the drugs, including Adderall and Concerta, are among the medications most frequently prescribed for American children, adding: **"If 5 percent of families in the country are giving a medication to their children, and they don't realize it does not have long-term benefits but might have long-term risks, why should they not be told?"**

The disagreement has produced a range of views among the researchers about how to accurately present the results to the public. One e-mail noted that an academic review of the group's work, called the Multimodal Treatment Study of Children With ADHD (MTA), asked why the researchers were "bending over backward" to play down negative implications for drug therapy.

Peter Jensen, one of Pelham's fellow researchers, responded that Pelham was biased against the use of drugs and was substituting his personal opinion for science.

Jensen said Pelham was the only member of the team of researchers who took away "the silly message" that the study raised questions about the long-term utility of drugs, but interviews and e-mails show that Pelham was not alone.

The MTA was designed to test whether children diagnosed with attention-deficit hyperactivity disorder, or ADHD, do better when treated with drugs, with drugs plus talk therapy, with talk therapy alone or with routine medical care alone. Children with the disorder have trouble paying attention, are restless and hyperactive, and are sometimes disruptive in school.

The initial 14-month analysis published in 1999 randomly assigned children to one of four treatment options and **showed clearly that those treated with medication did much better than those who got only talk therapy or routine care.** The drugs' manufacturers distributed thousands of reprints of the article to physicians at a

time when diagnoses of ADHD were spiraling upward. Because children given drugs alone appeared to do about as well as those treated with both drugs and talk therapy, the study skewed treatment in the direction of medication.

In a second phase of the study, the researchers followed the children and compared how they fared, but researchers no longer randomly assigned them to the various treatment options, making this phase less scientifically rigorous.

In August 2007, the MTA researchers reported the first follow-up data, which by then no longer showed differences in behavior between children who were medicated and those who were not. But the data did show that children who took the drugs for 36 months were about an inch shorter and six pounds lighter than those who did not.

A news release issued by the National Institute of Mental Health (NIMH) at the time, however, presented the results in a more favorable light. The release, dated July 20, 2007, was titled "Improvement Following ADHD Treatment Sustained in Most Children." The release noted that the initial advantages of drug treatment were no longer evident, but it quoted Jensen as saying this did not mean that long-term drug therapy was ineffective.

Jensen said, "We were struck by the remarkable improvement in symptoms and functioning across all treatment groups." And rather than saying the growth of children on medication was stunted, the release said children who were not on medication "grew somewhat larger."

As the MTA study continued to find smaller and smaller behavioral differences between children who were medicated and those who were not, use of the drugs soared. Pelham said most parents and doctors took away the message that the study had found drug therapy effective over the long run. In 2004, physicians wrote 28.3 million prescriptions for ADHD drugs; last year, they wrote 39.5 million, according to data provided by IMS Health.

With the MTA having followed the children for eight years, the latest data have confirmed that there are no long-term differences between children who were continuously medicated and those who were never medicated. Some of the data were published online yesterday in the Journal of the American Academy of Child and Adolescent Psychiatry.

In a telephone interview, Jensen denied that the researchers had misled the public, pointing out that some children getting the drugs did do better over the long term. Looking at overall results was not as useful as studying how particular groups of children fared, he said.

Jensen and another co-author, L. Eugene Arnold at Ohio State University, who are both psychiatrists, emphasized the importance of individualizing treatment -- and warned parents against abruptly terminating drug therapy.

The subgroup analysis found that children in homes that were socially and economically stable did the same in the long term with or without medication. Children from troubled or deprived backgrounds slid backward as soon as the intensive therapy stopped and they went back to their communities. About one-third -- those with the least impairment to begin with -- continued to improve over the long term.

Jensen and co-author Benedetto Vitiello at the NIMH said drugs may not have shown an overall long-term benefit because the quality of routine care that children received may have been inferior to the care they got during the initial part of the study. Jensen said the take-home message is that community care needs improvement.

Brooke Molina, also a co-author and a University of Pittsburgh associate professor of psychology and

psychiatry, argued in an e-mail that if the researchers wanted to draw attention to subgroups that might be helped by medication over the long run, they also should acknowledge that "long-term treatment with medication may not be efficacious" for others.

In an interview, Molina said the data do not "support that children who stay on medication longer than two years have better outcomes than children who don't." In an e-mail she shared with Pelham, she noted that academic "reviewers thought we were bending over backward (inappropriately) to dismiss the failure to find medication effects at 8 years."

James Swanson, another MTA co-author and a psychologist at the University of California at Irvine, said he believes that the researchers have been open about the diminishing benefits of medication therapy. He cited a variety of scientific publications in which he and others reported data showing that medications lost effectiveness over time and stunted growth.

"If you want something for tomorrow, medication is the best, but if you want something three years from now, it does not matter," he said. "If you take medication long-term beyond three years, I don't think there is any evidence that medication is better than no medication."

Pelham, who has conducted many drug therapy studies, said the drugs have a valuable role: They buy parents and clinicians time to teach youngsters behavioral strategies to combat inattention and hyperactivity. Over the long term, he said, parents need to rely on those skills.

A yet-to-be-published study, Pelham added, found that 95 percent of parents who were told by clinicians to first try behavioral interventions for ADHD did so. When parents were given a prescription for a drug and then told to enroll their children in behavioral intervention programs, 75 percent did not seek out the behavioral approaches.

Post a Comment

Comments that include profanity or personal attacks or other inappropriate comments or material will be removed from the site. Additionally, entries that are unsigned or contain "signatures" by someone other than the actual author will be removed. Finally, we will take steps to block users who violate any of our posting standards, terms of use or privacy policies or any other policies governing this site. Please review the [full rules](#) governing commentaries and discussions. You are fully responsible for the content that you post.

© 2009 The Washington Post Company